

# RESTRICTED RADON TRANSFORMS AND UNIONS OF HYPERPLANES

DANIEL M. OBERLIN

Department of Mathematics, Florida State University

October 2004

## §1 Introduction

If  $\Sigma^{(n-1)}$  is the unit sphere in  $\mathbb{R}^n$ , the Radon transform  $Rf$  of a suitable function  $f$  on  $\mathbb{R}^n$  is defined by

$$Rf(\sigma, t) = \int_{\sigma^\perp} f(p + t\sigma) dm_{n-1}(p) \quad \sigma \in \Sigma^{(n-1)}, t \in \mathbb{R},$$

where the integral is with respect to  $(n-1)$ -dimensional Lebesgue measure on the hyperplane  $\sigma^\perp$ . We also define, for  $0 < \delta < 1$ ,

$$R_\delta f(\sigma, t) = \delta^{-1} \int_{[\sigma^\perp \cap B(0,1)] + B(0,\delta)} f(x + t\sigma) dm_n(x).$$

The paper [5] is concerned with the mapping properties of  $R$  from  $L^p(\mathbb{R}^n)$  into mixed norm spaces defined by the norms

$$\left( \int_{\Sigma^{(n-1)}} \left[ \int_{-\infty}^{\infty} |g(\sigma, t)|^r dt \right]^{q/r} d\sigma \right)^{1/q}.$$

Here  $d\sigma$  denotes Lebesgue measure on  $\Sigma^{(n-1)}$ . The purpose of this paper is mainly to study the possibility of analogous mixed norm estimates when  $d\sigma$  is replaced by measures  $d\mu(\sigma)$  supported on compact subsets  $S \subseteq \Sigma^{(n-1)}$  having dimension  $< n-1$ . We are usually interested in the case  $r = \infty$  and will mostly settle for estimates of restricted weak type in the indices  $p$  and  $q$  and those only for  $f$  supported in a ball. The following theorem, which we regard as an estimate for a restricted Radon transform, is typical of our results here:

**Theorem 1.** *Fix  $\alpha \in (1, n-1)$ . Suppose  $\mu$  is a nonnegative and finite Borel measure on  $\Sigma^{(n-1)}$  satisfying the Frostman condition*

$$\int_{\Sigma^{(n-1)}} \int_{\Sigma^{(n-1)}} \frac{d\mu(\sigma_1)d\mu(\sigma_2)}{|\sigma_1 - \sigma_2|^\alpha} < \infty.$$

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\TeX}$

Then

$$(1) \quad \lambda \mu(\{\sigma \in \Sigma^{(n-1)} : \sup_{t \in \mathbb{R}} R\chi_E(\sigma, t) > \lambda\})^{1/\alpha} \lesssim |E|^{1/2}$$

for  $\lambda > 0$  and Borel  $E \subseteq B(0, 1)$ . That is,

$$\|R\chi_E\|_{L_\mu^{\alpha, \infty}(L^\infty)} \lesssim |E|^{1/2}.$$

Suppose that  $\alpha \in (0, n-1)$ . Say that a Borel set  $E \subseteq \mathbb{R}^n$  satisfies the (Besicovitch) condition  $B(n-1; \alpha)$  if there is a compact set  $S \subseteq \Sigma^{(n-1)}$  having Hausdorff dimension  $\alpha$  such that for each  $\sigma \in S$  there is a translate of an  $(n-1)$ -plane orthogonal to  $\sigma$  which intersects  $E$  in a set of positive  $(n-1)$ -dimensional Lebesgue measure. It is well-known that, given  $\epsilon \in (0, \alpha)$ , such an  $S$  supports a probability measure  $\mu$  satisfying the hypothesis of Theorem 1, but with  $\alpha - \epsilon$  in place of  $\alpha$ . In conjunction with Theorem 1, standard arguments imply that such an  $E$  must have positive  $n$ -dimensional Lebesgue measure. That is,  $B(n-1; \alpha)$  sets in  $\mathbb{R}^n$  have positive Lebesgue measure if  $\alpha > 1$ . As will be pointed out in §2, the next theorem implies that, for  $\alpha \in (0, 1)$  and in certain cases,  $B(n-1; \alpha)$  sets have Hausdorff dimension at least  $n-1+\alpha$ . (Here is a notational comment:  $|E|$  will usually denote the Lebesgue measure of  $E$  with the appropriate dimension being clear from the context.)

**Theorem 2.** *Suppose  $\alpha \in (0, 1)$ . Suppose  $\tilde{\mu}$  is a nonnegative measure on a compact interval  $J \subseteq \mathbb{R}$  which satisfies the condition*

$$\tilde{\mu}(I) \lesssim |I|^\alpha$$

for subintervals  $I \subseteq J$ . Let  $\mu$  be the image of  $\tilde{\mu}$  under a one-to-one and bi-Lipschitz mapping of  $J$  into  $\Sigma^{(n-1)}$ . If  $0 < \gamma < \beta < \alpha$  and

$$\frac{1}{p} = \frac{1 + \beta - \gamma}{1 + 2\beta - \gamma}, \quad \frac{1}{q} = \frac{1 + \gamma}{1 + 2\beta - \gamma}, \quad \eta = \frac{1 - \gamma}{1 + 2\beta - \gamma}$$

then there is the estimate

$$\|R_\delta \chi_E\|_{L_\mu^{q, \infty}(L^\infty)} \lesssim |E|^{1/p} \delta^{-\eta}$$

for Borel  $E \subset B(0, 1)$  and  $\delta \in (0, 1)$ .

Contrasting with Theorems 1 and 2, the next result provides a global estimate for a restricted Radon transform:

**Theorem 3.** *Suppose  $n \geq 4$ . Let  $S$  be the  $(n-2)$ -sphere*

$$\{\sigma = (\sigma_1, \dots, \sigma_n) \in \Sigma^{(n-1)} : \sum_1^{n-1} \sigma_j^2 = \sigma_n^2\}$$

and let  $\mu$  be Lebesgue measure on  $S$ . Then there is an estimate

$$\|R\chi_E\|_{L_\mu^{n-2}(L^\infty)} \lesssim |E|^{(n-1)/n}$$

for Borel  $E \subseteq \mathbb{R}^n$ .

This result is an analogue of (3) in [5] (which is a similar estimate but with  $\mu$  replaced by Lebesgue measure on  $\Sigma^{(n-1)}$  and  $q = n$ ). The proof of Theorem 3 parallels the proof in [5] but requires the  $L^2$  Fourier restriction estimates for the light cone in  $\mathbb{R}^n$  in place of an easier  $L^2$  estimate used in [5].

The remainder of this paper is organized as follows: §2 contains the proofs of Theorems 2 and 3 and the statement and proof of a similar result in the case when  $d$  is an integer strictly between 1 and  $n - 1$  and  $\mu$  is Lebesgue measure on a suitable  $d$ -manifold in  $\Sigma^{(n-1)}$ ; §3 contains the proof of Theorem 3; §4 contains some miscellaneous observations and remarks: an analogue for Kahane's notion of Fourier dimension of Theorem 2 when  $n = 2$  and examples bearing on the question of whether  $B(n - 1; 1)$  sets in  $\mathbb{R}^n$  must have positive measure or only full dimension (the answer depends on the set  $S$  of directions).

### §2 Proofs of Theorems 1 and 2

As Theorem 1 is a consequence of its analogue, uniform in  $\delta \in (0, 1)$ , for the operators  $R_\delta$ , we will restrict our attention to these operators. A standard method for obtaining restricted weak type estimates is to estimate  $|E|$  from below. We will do this with a particularly simple-minded strategy based on two observations and originally employed in [3] and [4]. The first observation is that

$$|\cup_{n=1}^N E_n| \geq \sum_{n=1}^N |E_n| - \sum_{1 \leq m < n \leq N} |E_m \cap E_n|.$$

The second is the well-known fact that if  $\sigma \in \Sigma^{(n-1)}$  and if, for  $t \in \mathbb{R}$ ,  $P_\sigma^\delta$  denotes a plate  $[\sigma^\perp \cap B(0, 1)] + B(0, \delta) + t\sigma$ , then

$$|P_{\sigma_1}^\delta \cap P_{\sigma_2}^\delta| \leq \frac{C(n)\delta^2}{|\sigma_1 - \sigma_2|}$$

(so long as  $\sigma_1$  and  $\sigma_2$  are not too far apart, an hypothesis we tacitly assume since it can be achieved by multiplying the measures  $\mu$  appearing below by an appropriate partition of unity). Thus if, for  $n = 1, \dots, N$ , we have plates  $P_{\sigma_n}^\delta$  satisfying  $|E \cap P_{\sigma_n}^\delta| \geq C_1\lambda\delta$ , it follows that

$$(2) \quad |E| \geq C_1N\lambda\delta - C(n)\delta^2 \sum_{1 \leq m < n \leq N} \frac{1}{|\sigma_m - \sigma_n|}.$$

Our strategy, then, will be to choose  $N$  and

$$\sigma_n \in \{\sigma \in \Sigma^{(n-1)} : \sup_{t \in \mathbb{R}} R_\delta \chi_E(\sigma, t) > \lambda\}$$

so that (2) gives, for example,

$$|E| \gtrsim \lambda^2 \mu(\{\sigma \in \Sigma^{(n-1)} : \sup_{t \in \mathbb{R}} R_\delta \chi_E(\sigma, t) > \lambda\})^{2/\alpha},$$

which is the analogue of (1) for the operator  $R_\delta$ . For Theorem 1 the following lemma will facilitate this choice:

**Lemma 1.** *Let  $\mu$  be as in Theorem 1. There is  $C = C(\mu)$  such that given  $n \in \mathbb{N}$  and a Borel  $S \subseteq \Sigma^{(n-1)}$  with  $\mu(S) > 0$ , one can choose  $\sigma_n \in S$ ,  $1 \leq n \leq N$ , such that*

$$\sum_{1 \leq m < n \leq N} \frac{1}{|\sigma_m - \sigma_n|} \leq \frac{CN^2}{\mu(S)^{2/\alpha}}.$$

*Proof of Lemma 1:* Suppose  $\sigma_1, \dots, \sigma_N$  are chosen independently and at random from the probability space  $(S, \frac{\mu}{\mu(S)})$ . Then, for  $1 \leq m < n \leq N$ ,

$$\begin{aligned} \mathbb{E}\left(\frac{1}{|\sigma_m - \sigma_n|}\right) &= \frac{1}{\mu(S)^2} \int_S \int_S \frac{1}{|\sigma_m - \sigma_n|} d\mu(\sigma_m) d\mu(\sigma_n) \leq \\ \frac{1}{\mu(S)^2} \left(\int_S \int_S 1 d\mu(\sigma_m) d\mu(\sigma_n)\right)^{1-1/\alpha} &\left(\int_S \int_S \frac{1}{|\sigma_m - \sigma_n|^\alpha} d\mu(\sigma_m) d\mu(\sigma_n)\right)^{1/\alpha} \leq \frac{C}{\mu(S)^{2/\alpha}}, \end{aligned}$$

by the hypothesis on  $\mu$ . Thus

$$\mathbb{E}\left(\sum_{1 \leq m < n \leq N} \frac{1}{|\sigma_m - \sigma_n|}\right) \leq \frac{CN^2}{\mu(S)^{2/\alpha}}$$

and the lemma follows.

*Proof of Theorem 1:* Let  $S$  be the set

$$\{\sigma \in \Sigma^{(n-1)} : \sup_{t \in \mathbb{R}} R_\delta \chi_E(\sigma, t) > \lambda\}$$

so that if  $\sigma \in S$  then there is  $t \in \mathbb{R}$  such that if  $P_\sigma^\delta = [\sigma^\perp \cap B(0, 1)] + B(0, \delta) + t\sigma$  then  $|E \cap P_\sigma^\delta| \geq C_1 \lambda \delta$ . The conjunction of Lemma 1 and (2) yields

$$(3) \quad |E| \geq C_1 N \lambda \delta - C_2 \delta^2 N^2 \mu(S)^{-2\alpha}.$$

We consider two cases (noting that  $N = N_0 \doteq \lambda C_1 \mu(S)^{2/\alpha} / C_2 \delta$  makes the RHS of (3) equal to 0):

*Case I:* Assume  $N_0 > 10$ .

In this case choose  $N \in \mathbb{N}$  such that

$$\frac{\lambda C_1 \mu(S)^{2/\alpha}}{2C_2 \delta} \geq N \geq \frac{\lambda C_1 \mu(S)^{2/\alpha}}{3C_2 \delta}.$$

Then it follows from (3) that

$$|E| \geq C_1 \frac{\lambda C_1 \mu(S)^{2/\alpha}}{3C_2 \delta} \lambda \delta - C_2 \delta^2 \frac{\lambda^2 C_1^2 \mu(S)^{4/\alpha}}{4C_2^2 \delta^2} \mu(S)^{-2/\alpha} = \kappa \lambda^2 \mu(S)^{2/\alpha}$$

for  $\kappa = C_1^2 / (12C_2)$ . This gives  $\lambda \mu(S)^{1/\alpha} \lesssim |E|^{1/2}$  as desired.

*Case II:* Assume  $N_0 \leq 10$ .

In this case (unless  $S$  is empty) we estimate

$$|E| \geq C_1 \lambda \delta \geq \frac{\lambda^2 C_1^2 \mu(S)^{2/\alpha}}{10C_2}$$

which again yields  $\lambda \mu(S)^{1/\alpha} \lesssim |E|^{1/2}$  and so completes the proof of Theorem 1.

The proof of Theorem 2 requires an analogue of Lemma 1:

**Lemma 2.** *Suppose  $\mu$  is as in Theorem 2. Suppose  $0 < \gamma < \beta < \alpha$ . Then there is  $C = C(\alpha, \mu, \beta, \gamma)$  such that given a Borel  $S \subseteq \Sigma^{(n-1)}$  with  $\mu(S) > 0$  and  $N \in \mathbb{N}$ , one can choose  $\sigma_n \in S$ ,  $1 \leq n \leq N$ , such that*

$$\sum_{1 \leq m < n \leq N} \frac{1}{|\sigma_m - \sigma_n|} \leq \frac{CN^{(1+2\beta-\gamma)/\beta}}{\mu(S)^{(1+\gamma)/\beta}}.$$

*Proof of Lemma 2:* It suffices to show that there exists  $C$  such that if  $F$  is a measurable subset of  $J$  with  $\tilde{\mu}(F) > 0$  and if  $N \in 2\mathbb{N}$ , then there are  $x_1, \dots, x_{N/2}$  in  $F$  such that

$$(4) \quad \sum_{1 \leq m < n \leq N/2} \frac{1}{|x_m - x_n|} \leq \frac{CN^{(1+2\beta-\gamma)/\beta}}{\tilde{\mu}(F)^{(1+\gamma)/\beta}}.$$

Note that because  $\beta < \alpha$  it follows that  $\tilde{\mu}(I) \lesssim |I|^\beta$  for subintervals  $I$  of  $J$ . Now define  $\eta$  by  $\eta^\beta = \tilde{\mu}(F)/N$  and find  $a_1 < b_1 \leq a_2 < \dots < b_N$  in  $J$  such that  $\tilde{\mu}(F \cap [a_n, b_n]) = \eta^\beta$ . Let  $I_n = [a_n + \eta/L, b_n - \eta/L]$  where  $L$  is chosen large enough to guarantee that  $\tilde{\mu}(F \cap I_n) \geq \eta^\beta/2$  and then find intervals  $\tilde{I}_n \subseteq I_n$  satisfying  $\tilde{\mu}(F \cap \tilde{I}_n) = \eta^\beta/2$ . Choose Borel mappings

$$\tau_n : [0, \eta^\beta/2] \rightarrow F \cap \tilde{I}_n$$

such that the equalities

$$\int_{F \cap \tilde{I}_n} f \, d\tilde{\mu} = \int_0^{\eta^\beta/2} f(\tau_n(s)) \, dm_1(s)$$

hold for reasonable functions  $f$  on  $F \cap \tilde{I}_n$ . Then

$$\int_0^{\eta^\beta/2} \int_0^{\eta^\beta/2} \sum_{n \neq m} \frac{dm_1(s) \, dm_1(t)}{|\tau_m(s) - \tau_n(t)|} = \sum_{n \neq m} \int_{F \cap \tilde{I}_m} \int_{F \cap \tilde{I}_n} \frac{d\tilde{\mu}(x) \, d\tilde{\mu}(y)}{|x - y|}.$$

Since  $\gamma < 1$  and  $d(\tilde{I}_m, \tilde{I}_n) \geq \eta/L$ , the last sum is

$$\begin{aligned} &\leq C\eta^{\gamma-1} \int_F \int_F \frac{d\tilde{\mu}(x) \, d\tilde{\mu}(y)}{|x - y|^\gamma} \leq \\ &C\eta^{\gamma-1} \left( \int_F \int_F \frac{d\tilde{\mu}(x) \, d\tilde{\mu}(y)}{|x - y|^\beta} \right)^{\gamma/\beta} \tilde{\mu}(F)^{2(1-\gamma/\beta)} = C\eta^{\gamma-1} \tilde{\mu}(F)^{2(1-\gamma/\beta)} \end{aligned}$$

since

$$\int_J \int_J \frac{d\tilde{\mu}(x) \, d\tilde{\mu}(y)}{|x - y|^\beta} < \infty$$

follows from the hypothesis on  $\tilde{\mu}$  and the fact that  $\beta < \alpha$ . Thus

$$\frac{1}{(\eta^\beta/2)^2} \int_0^{\eta^\beta/2} \int_0^{\eta^\beta/2} \sum_{n \neq m} \frac{dm_1(s) dm_1(t)}{|\tau_m(s) - \tau_n(t)|} \leq C\eta^{-2\beta+\gamma-1} \tilde{\mu}(F)^{2(1-\gamma/\beta)} =$$

$$C \left( \frac{\tilde{\mu}(F)}{N} \right)^{(-2\beta+\gamma-1)/\beta} \tilde{\mu}(F)^{2(1-\gamma/\beta)} = CN^{(2\beta-\gamma+1)/\beta} \tilde{\mu}(F)^{-(1+\gamma)/\beta}.$$

It follows that there are  $s, t \in [0, \eta^\beta/2]$  such, for  $m, n = 1, \dots, N$ , the points

$$x_n = \tau_n(s) \in F \cap \tilde{I}_n, \quad y_m = \tau_m(t) \in F \cap \tilde{I}_m$$

satisfy

$$\sum_{n \neq m} \frac{1}{|x_m - y_n|} \leq \frac{CN^{(2\beta-\gamma+1)/\beta}}{\tilde{\mu}(F)^{(1+\gamma)/\beta}}.$$

Now either  $x_n \leq y_n$  for at least  $N/2$   $n$ 's or  $y_n \leq x_n$  for at least  $N/2$   $n$ 's. Without loss of generality, consider the first case and let

$$\mathcal{N} = \{n = 1, \dots, N : x_n \leq y_n\}.$$

If  $n_1, n_2 \in \mathcal{N}$  and  $n_1 < n_2$  then (because  $y_{n_1} \in I_{n_1}$  and  $x_{n_2} \in I_{n_2}$ ), we have

$$x_{n_1} \leq y_{n_1} < x_{n_2} \leq y_{n_2}$$

and so

$$|x_{n_1} - x_{n_2}| > |y_{n_1} - x_{n_2}|.$$

Thus

$$\sum_{\substack{n_1 < n_2 \\ n_1, n_2 \in \mathcal{N}}} \frac{1}{|x_{n_1} - x_{n_2}|} < \sum_{\substack{n_1 < n_2 \\ n_1, n_2 \in \mathcal{N}}} \frac{1}{|y_{n_1} - x_{n_2}|} \leq$$

$$\sum_{n \neq m} \frac{1}{|x_m - y_n|} \leq \frac{CN^{(2\beta-\gamma+1)/\beta}}{\tilde{\mu}(F)^{(1+\gamma)/\beta}}.$$

Renumbering the  $x_n$  ( $n \in \mathcal{N}$ ) gives (4) and completes the proof of the lemma.

*Proof of Theorem 2:* The proof is parallel to that of Theorem 1. Using Lemma 2 instead of Lemma 1, the analogue of (3) is

$$(5) \quad |E| \geq C_1 N \lambda \delta - C_2 \delta^2 N^{(1+2\beta-\gamma)/\beta} \mu(S)^{-(1+\gamma)/\beta}.$$

The two cases are now defined by comparing

$$N_0 \doteq \left( \frac{C_1 \lambda}{C_2 \delta} \right)^{\beta/(1+\beta-\gamma)} \mu(S)^{\frac{1+\gamma}{1+\beta-\gamma}}$$

and 10. In case  $N_0 > 10$ , choosing  $N$  in (5) such that  $N_0/2 \geq N \geq N_0/3$  gives

$$|E| \geq \lambda^{\frac{1+2\beta-\gamma}{1+\beta-\gamma}} \delta^{\frac{1-\gamma}{1+\beta-\gamma}} \mu(S)^{\frac{1+\gamma}{1+\beta-\gamma}} \kappa$$

where

$$\kappa = C_1^{\frac{1+2\beta-\gamma}{1+\beta-\gamma}} C_2^{\frac{-\beta}{1+\beta-\gamma}} \left( \frac{1}{3} - \frac{1}{2(1+2\beta-\gamma)/\beta} \right) > 0.$$

This leads directly to the desired estimate  $\lambda\mu(S)^{1/q} \lesssim |E|^{1/p}\delta^{-\eta}$  if  $N_0 > 10$ . On the other hand, the inequality  $N_0 \leq 10$  gives  $\lambda\mu(S)^{(1+\gamma)/\beta} \lesssim \delta$  and so

$$(6) \quad \lambda^A \mu(S)^{A(1+\gamma)/\beta} \lesssim \delta^A$$

if  $A > 0$ . Since  $|E| \geq C_1\lambda\delta$  (unless  $S$  is empty), there is also the inequality

$$(7) \quad \lambda^{1-A} \lesssim |E|^{1-A}\delta^{A-1}$$

as long as  $0 < A < 1$ . Multiplying (6) and (7) gives  $\lambda\mu(S)^{A(1+\gamma)/\beta} \lesssim |E|^{1-A}\delta^{2A-1}$ . Then the choice  $A = \beta/(1+2\beta-\gamma)$  yields  $\lambda\mu(S)^{1/q} \lesssim |E|^{1/p}\delta^{-\eta}$  again, completing the proof of Theorem 2.

It follows from the proof of Lemma 2.15 in [1] that the estimate

$$\|R_\delta\chi_E\|_{L_\mu^{q,\infty}(L^\infty)} \lesssim |E|^{1/p}\delta^{-\eta}$$

provides a lower bound of  $n - p\eta$  for the Hausdorff dimension of a Borel set containing positive-measure sections of hyperplanes associated with each of the directions  $\sigma$  in the support of  $\mu$ . Plugging in the values for  $p$  and  $\eta$  which are given in Theorem 2 yields first the lower bound  $n - (1 - \gamma)/(1 + \beta - \gamma)$  and then, since that is valid for  $0 < \gamma < \beta < \alpha$ , the desired lower bound of  $n - 1 + \alpha$ . A subset  $S \subseteq \Sigma^{(n-1)}$  of Hausdorff dimension  $\alpha \in (0, 1)$  and located on a curve as in the hypotheses of Theorem 2, will, for each  $\epsilon \in (0, \alpha)$ , support a measure  $\mu$  satisfying the hypotheses of Theorem 2, but with  $\alpha - \epsilon$  instead of  $\alpha$ . It follows that the  $B(n - 1; \alpha)$  sets associated with such sets of directions  $S$  will all have Hausdorff dimension at least  $n - 1 + \alpha$ . Finally, note that if  $n = 2$  then the hypothesis that  $\mu$  be supported on a curve is no restriction and so all  $B(1; \alpha)$  sets in  $\mathbb{R}^2$  have dimension at least  $1 + \alpha$ .

The next result gives, in certain special situations, an improvement over Theorem 1 on the index  $q$  in the bound  $\|R\chi_E\|_{L_\mu^{q,\infty}(L^\infty)} \lesssim |E|^{1/2}$ .

**Proposition 1.** *Suppose  $d \in \mathbb{N}$ ,  $1 < d < n - 1$ . Suppose that  $\mu$  is the image of Lebesgue measure on a closed ball in  $\mathbb{R}^d$  under a bi-Lipschitz mapping of that ball into  $\Sigma^{(n-1)}$ . Then for Borel  $E \subseteq B(0, 1)$  there is the estimate*

$$\|R\chi_E\|_{L_\mu^{2d,\infty}(L^\infty)} \lesssim |E|^{1/2}.$$

*Proof of Proposition 1:* The proof is again analogous to the proof of Theorem 1. The required analogue of Lemma 1 is

**Lemma 3.** *Suppose  $\mu$  is as in Theorem 4. Then there is  $C$  such that given a Borel  $S \subseteq \Sigma^{(n-1)}$  with  $\mu(S) > 0$  and given  $N \in \mathbb{N}$ , one can choose  $\sigma_n \in S$ ,  $1 \leq n \leq N$ , such that*

$$\sum_{1 \leq m < n \leq N} \frac{1}{|\sigma_m - \sigma_n|} \leq \frac{CN^2}{\mu(S)^{1/d}}.$$

*Proof of Lemma 3:* Letting  $\eta > 0$  be defined by  $\eta^d = \mu(S)/(CN)$ , where  $C$  is sufficiently large, choose  $N$   $\eta$ -separated points  $\sigma_1, \dots, \sigma_N$  from  $S$ . Then, for fixed  $m$ ,

$$\sum_{n \neq m} \frac{1}{|\sigma_m - \sigma_n|} \lesssim \eta^{-d} \int_{\cup_n B(\sigma_n, \eta/2)} \frac{d\sigma}{|\sigma_m - \sigma|}.$$

The function  $\sigma \mapsto |\sigma_m - \sigma|^{-1}$  is in  $L^{d,\infty}(d\mu)$ . So, still for fixed  $m$ ,

$$\sum_{n \neq m} \frac{1}{|\sigma_m - \sigma_n|} \lesssim \eta^{-d} (N\eta^d)^{1-1/d}.$$

The lemma follows from the choice of  $\eta$  by summing on  $m$ .

Returning to the proof of Proposition 1, the analogue of (3) is now

$$|E| \geq C_1 N \lambda \delta - C_2 \delta^2 N^2 \mu(S)^{-1/d},$$

the choice for  $N_0$  is  $\lambda C_1 \mu(S)^{1/d} / (C_2 \delta)$ , and the remainder of the proof of Proposition 1 is completely parallel to that of Theorem 1.

### §3 Proof of Theorem 3

As previously mentioned, the proof is an adaptation of the proof of (3) in [5]. We begin by noting that

$$\widehat{Rf(\sigma, \cdot)}(y) = \int_{-\infty}^{\infty} e^{-2\pi i y t} \int_{\sigma^\perp} f(p + t\sigma) dm_{n-1}(p) dm_1(t) = \widehat{f}(y\sigma).$$

Thus

$$\|Rf\|_{L_{d\mu}^2(L^2)}^2 = \int_S \int_{-\infty}^{\infty} |\widehat{f}(y\sigma)|^2 dm_1(y) d\mu(\sigma) = \int_{\mathbb{R}^{(n-1)}} |\widehat{f}(\xi, |\xi|)|^2 \frac{d\xi}{|\xi|^{n-2}}$$

and so estimates for  $R$  as a mapping into  $L_\mu^2(L^2)$  are just Fourier restriction estimates for the light cone in  $\mathbb{R}^n$ . More generally, we have

$$\left\| \left( \frac{\partial}{\partial t} \right)^\beta Rf \right\|_{L_\mu^2(L^2)}^2 = \int_{\mathbb{R}^{(n-1)}} |\widehat{f}(\xi, |\xi|)|^2 \frac{d\xi}{|\xi|^{n-2-2\beta}}.$$

Thus the results of 5.17(b) on p. 367 in [6] give the estimate

$$(8) \quad \left\| \left( \frac{\partial}{\partial t} \right)^\beta Rf \right\|_{L_\mu^2(L^2)} \lesssim \|f\|_p$$

whenever

$$-\frac{1}{2} < \beta \leq \frac{n-3}{2} \text{ and } \frac{1}{p} = \frac{2n-2\beta-1}{2n}.$$

Estimate (8) will lead to a mixed norm estimate in which the ‘‘inside’’ norm is a Lipschitz norm. The proof of Theorem 3 is simply an interpolation of this estimate with the trivial  $L^1 \rightarrow L^\infty(L^1)$  estimate for  $R$ . The following generalization of an observation from [5] allows this interpolation.



**Lemma 4.** Fix  $\alpha > 0$  and  $m \in \mathbb{N}$  with  $m > \alpha$ . For a Borel function  $g$  on  $\mathbb{R}$  and for  $t \in \mathbb{R}$ , write  $\Delta_t$  for the usual difference operator given by  $\Delta_t g(x) = g(x+t) - g(x)$ ,  $x \in \mathbb{R}$ . Let  $\|g\|_\alpha$  be the Lipschitz norm given by

$$\|g\|_\alpha = \sup_{x \in \mathbb{R}, t \neq 0} \frac{|\Delta_t^m g(x)|}{|t|^\alpha}.$$

Then, for  $1 \leq r < \infty$ , we have

$$\|g\|_{L^\infty} \lesssim \|g\|_{L^{r,\infty}}^{\alpha r/(1+\alpha r)} \|g\|_\alpha^{1/(1+\alpha r)}.$$

*Proof of Lemma 4:* Write

$$\Delta_t^m g(x) = \sum_{j=1}^m c_j g(x+jt) \pm g(x).$$

Assume that  $|g(x)| \geq \lambda$  for some fixed  $x \in \mathbb{R}$  and some  $\lambda > 0$ . If  $|t|$  is so small that

$$|t|^\alpha \|g\|_\alpha \leq \frac{\lambda}{2}$$

then

$$\left| \sum_{j=1}^m c_j g(x+jt) \right| \geq \frac{\lambda}{2}.$$

Thus

$$\frac{\lambda}{2} \left( 2 \left( \frac{\lambda}{2 \|g\|_\alpha} \right)^{1/\alpha} \right)^{1/r} \leq \left\| \sum_{j=1}^m c_j g(x+jt) \right\|_{L_t^{r,\infty}} \lesssim \|g\|_{L^{r,\infty}}$$

and so

$$\lambda \lesssim \|g\|_{L^{r,\infty}}^{\alpha r/(1+\alpha r)} \|g\|_\alpha^{1/(1+\alpha r)}.$$

Since  $x \in \mathbb{R}$  and  $\lambda \leq |g(x)|$  were arbitrary, the desired inequality follows and the proof of Lemma 4 is complete.

For the remainder of this section, the “outside” norms  $\|\cdot\|_{L^s}$  will refer to the measure  $\mu$  on  $S$  while  $\|\cdot\|_p$  will be the norm on  $L^p(\mathbb{R}^n)$  (or on  $L^p(\mathbb{R})$ ) and  $\|\cdot\|_\alpha$  will be the Lipschitz norm of Lemma 4. Taking  $r = 1$  in Lemma 4 gives

$$(9) \quad \|Rf\|_{L^{n-2}(L^\infty)} \lesssim \|Rf\|_1^{\alpha/(1+\alpha)} \|Rf\|_\alpha^{1/(1+\alpha)} \|Rf\|_{L^{n-2}}.$$

Since

$$\|Rf(\sigma, \cdot)\|_1 \leq \|f\|_1,$$

for all  $\sigma \in \Sigma^{(n-1)}$ , (9) gives

$$(10) \quad \|Rf\|_{L^{n-2}(L^\infty)} \lesssim \|f\|_1^{\alpha/(1+\alpha)} \|Rf\|_\alpha^{1/(1+\alpha)} \|Rf\|_{L^{n-2}}.$$

To bound the second term of the RHS of (10), we note that the estimate

$$\| \|Rf\|_\alpha \|_{L^2} \lesssim \left\| \left( \frac{\partial}{\partial t} \right)^{1/2+\alpha} Rf \right\|_{L^2(L^2)}$$

follows from Lemma 1 in [5]. Thus if

$$\alpha = \frac{n-4}{2} \quad \text{and} \quad \frac{1}{p} = \frac{n-1-\alpha}{n} = \frac{n+2}{2n},$$

(8) with  $\beta = 1/2 + \alpha$  yields

$$\| \|Rf\|_{\alpha}^{1/(1+\alpha)} \|_{L^{n-2}} = \| \|Rf\|_{\alpha} \|_{L^2}^{1/(1+\alpha)} \lesssim \left\| \left( \frac{\partial}{\partial t} \right)^{1/2+\alpha} Rf \right\|_{L^2(L^2)}^{1/(1+\alpha)} \lesssim \|f\|_{2n/(n+2)}^{1/(1+\alpha)} = \|f\|_{2n/(n+2)}^{2/(n-2)}.$$

With (10), this gives

$$\| \|R\chi_E\|_{L^\infty} \|_{L^{n-2}} \lesssim |E|^{(n-1)/n},$$

which is the desired result.

#### §4 Miscellany

##### Fourier dimension

As introduced by Kahane in [2], the Fourier dimension of a compact set  $E \subseteq \mathbb{R}^n$  is twice the least upper bound of the set of nonnegative  $\beta$ 's for which  $E$  carries a Borel probability measure  $\lambda$  satisfying  $|\widehat{\lambda}(\xi)| = o(|\xi|^\beta)$  for large  $|\xi|$ . It is observed in [2] that the Hausdorff dimension of  $E$  is always at least equal to the Fourier dimension of  $E$  and is generally strictly larger, since the Hausdorff dimension of  $E \subseteq \mathbb{R}^n$  does not change if  $\mathbb{R}^n$  is embedded in  $\mathbb{R}^{n+1}$  while the Fourier dimension of  $E$  now considered as a subset of  $\mathbb{R}^{n+1}$  will be 0. The next result is an analogue for Fourier dimension of the  $n = 2$  case of Theorem 2:

**Proposition 2.** *Suppose  $\alpha \in (0, 1)$  and  $S \subseteq \Sigma^{(1)}$  has Hausdorff dimension  $\alpha$ . Suppose that  $E$  is a compact subset of  $\mathbb{R}^2$  containing a unit line segment in each of the directions  $\sigma \in S$ . Then the Fourier dimension of  $E$  is at least  $2\alpha$ .*

Since Fourier dimension is generally strictly smaller than Hausdorff dimension, it is not surprising that our lower bound  $2\alpha$  for the Fourier dimension of  $E$  is strictly smaller than the lower bound  $1 + \alpha$  for the Hausdorff dimension of  $E$  which follows from Theorem 2. Still, it follows from Proposition 2 that Kakeya sets in  $\mathbb{R}^2$  have Fourier dimension 2, providing a different proof of the well-known fact that such sets have Hausdorff dimension 2. It would be interesting to have examples, for  $\alpha \in (0, 1)$ , of sets  $E$  as in the proposition and having Fourier dimension equal to  $2\alpha$ .

*Proof of Proposition 2:* The heuristic is simple: for each  $\beta < \alpha$ ,  $S$  carries a Borel probability measure  $\mu$  satisfying

$$(11) \quad \mu(J) \leq C |J|^\beta$$

for intervals  $J \subseteq \Sigma^{(1)}$  (where  $C$  depends on  $\beta$  and  $|J|$  denotes the ‘‘length’’ of  $J$ ). For each  $\sigma \in S$  find  $x_\sigma \in \mathbb{R}^2$  such that  $x_\sigma + t\sigma \in E$  if  $|t| \leq 1/2$ . Let  $\varphi \in C_0^\infty([-1/2, 1/2])$  be a nonnegative function with integral 1 and define the measure  $\lambda$  on  $E$  by

$$(12) \quad \int_E f \, d\lambda = \int_S \int_{-1/2}^{1/2} f(x_\sigma + t\sigma) \varphi(t) \, dt \, d\mu(\sigma).$$

Then

$$(13) \quad |\widehat{\lambda}(\xi)| \leq \int_S \left| \int_{-1/2}^{1/2} e^{-2\pi i \xi \cdot (x_\sigma + t\sigma)} \varphi(t) dt \right| d\mu(\sigma) = \int_S |\widehat{\varphi}(\xi \cdot \sigma)| d\mu(\sigma).$$

For each  $p \in \mathbb{N}$  there is  $C(p)$  such that

$$|\widehat{\varphi}(\xi \cdot \sigma)| \leq \frac{C(p)}{|\xi \cdot \sigma|^p}.$$

Thus for any  $\xi \in \mathbb{R}^2$  there are two intervals  $J_1, J_2 \subset \Sigma^{(1)}$  of length  $\eta > 0$  such that for  $\sigma \in \Sigma^{(1)} - (J_1 \cup J_2)$  we have

$$|\widehat{\varphi}(\xi \cdot \sigma)| \leq \frac{C(p)}{(|\xi|\eta)^p}.$$

With (11) and (13) this leads to

$$|\widehat{\lambda}(\xi)| \lesssim \eta^\beta + \frac{1}{(|\xi|\eta)^p}.$$

Optimizing with the choice  $\eta = |\xi|^{-p/(\beta+p)}$  then gives

$$(14) \quad |\widehat{\lambda}(\xi)| \leq C(\beta, p) |\xi|^{-\beta p/(\beta+p)},$$

and this implies the lower bound  $2\beta p/(\beta+p)$  for the Fourier dimension of  $E$ . As that bound should hold for  $0 < \beta < \alpha$  and for  $p \in \mathbb{N}$ , the desired lower bound  $2\alpha$  follows.

The problem with this heuristic argument lies, of course, in the measurability of the selection  $\sigma \mapsto x_\sigma$ . A standard approximation procedure circumvents this: for each  $N \in \mathbb{N}$ , partition  $\Sigma^{(1)}$  into  $N$  intervals  $J_1, \dots, J_N$  of length  $2\pi/N$ . Choose (if possible)  $\sigma_n \in J_n \cap S$  and define

$$\mu_N = \sum_{n=1}^N \mu(J_n) \delta_{\sigma_n}.$$

Define  $\lambda_N$  as in (12) but with  $\mu$  replaced by  $\mu_N$ . Then the argument above shows that

$$|\widehat{\lambda}_N(\xi)| \leq C(\beta, p) |\xi|^{-\beta p/(\beta+p)}$$

for  $|\xi| \leq N^{1+\beta/p}$ . Thus some weak\* limit point  $\lambda$  of the sequence  $\{\lambda_N\}$  will satisfy (14). This completes the proof of Proposition 2.

### $B(n-1; 1)$ sets

Recall that  $E \subseteq \mathbb{R}^n$  is a  $B(n-1; 1)$  set if there is a compact set  $S \subseteq \Sigma^{(n-1)}$  having Hausdorff dimension 1 such that for each  $\sigma \in S$  there is a hyperplane orthogonal to  $\sigma$  which intersects  $E$  in a set of positive  $(n-1)$ -dimensional Lebesgue measure. Although we have not proved it unless  $S$  sits on a nice curve in  $\Sigma^{(n-1)}$ , one expects that  $B(n-1; 1)$  sets should have Hausdorff dimension  $n$ . Here are some examples in dimension 3:

**Example 1.** Suppose that  $\tilde{E}$  is a (Kakeya) subset of  $\mathbb{R}^2 \hookrightarrow \mathbb{R}^3$  having 2-dimensional Lebesgue measure 0 and containing a line segment in each direction. If  $E$  is the product of  $\tilde{E}$  and a line segment orthogonal to  $\mathbb{R}^2$ , then  $E$  is a measure-zero  $B(2; 1)$  set having full dimension and associated with the 1-sphere of directions

$$S_1 \doteq \{\sigma = (\sigma_1, \sigma_2, \sigma_3) \in \Sigma^{(2)} : \sigma_3 = 0\}.$$

**Example 2.** Suppose that  $S \subseteq \Sigma^{(2)}$  is a compact set of Hausdorff dimension 1 which supports a measure  $\mu$  satisfying the condition

$$\int_S \int_S \frac{d\mu(\sigma_1)d\mu(\sigma_2)}{|\sigma_1 - \sigma_2|} < \infty.$$

(It is not too difficult to construct such an  $S$  and  $\mu$  using a Cantor set with variable ratio of dissection.) The proof of Theorem 1 yields in this case the estimate

$$\|R\chi_E\|_{L_\mu^{1,\infty}(L^\infty)} \lesssim |E|^{1/2}$$

for Borel  $E \subseteq \mathbb{R}^3$ . Thus any  $B(2; 1)$  set associated with the set of directions  $S$  must have positive measure.

**Example 3.** Consider the 1-sphere of directions

$$S_2 \doteq \{\sigma = (\sigma_1, \sigma_2, \sigma_3) \in \Sigma^{(2)} : \sigma_1^2 + \sigma_2^2 = \sigma_3^2\}.$$

As with  $S_2$  in Example 1, it follows from Theorem 2 that the  $B(2; 1)$  sets associated with  $S_2$  have full dimension. A difference between  $S_1$  and  $S_2$  appears when considering the possibility of

$$(15) \quad L^p \rightarrow L_{\mu_j}^2(L^2)$$

the circle  $S_j$ ). For  $j = 2$  there will be such an estimate for  $p = 6/5$ . This follows from (8) and, as mentioned in the proof of Theorem 3, is just the Tomas-Stein restriction theorem for the light cone in  $\mathbb{R}^3$ . On the other hand, there is no estimate (15) for  $\mu_1$  (because there are no Fourier restriction theorems for hyperplanes). It would be interesting to know whether, in contrast to the situation in Example 1, the  $B(2; 1)$  sets associated with  $S_2$  must actually have positive measure.

#### REFERENCES

1. J. Bourgain, *Besicovitch type maximal operators and applications to Fourier analysis*, Geometric and Functional Analysis **1** (1991), 147–187.
2. J.-P. Kahane, *Some Random Series of Functions*, Cambridge University Press, 1993.
3. D. Oberlin,  *$L^p \rightarrow L^q$  mapping properties of the Radon transform*, Springer Lecture Notes in Mathematics **995** (1983), 95–102.
4. ———, *An estimate for a restricted X-ray transform*, Canadian Mathematical Bulletin **43** (2000), 472–476.
5. D. Oberlin and E.M. Stein, *Mapping properties of the Radon transform*, Indiana University Mathematics Journal **31** (1982), 641–650.
6. E.M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.