

EXPLICIT FORMULAS FOR THE GROTHENDIECK CLASS OF $\overline{\mathcal{M}}_{0,n}$

PAOLO ALUFFI, MATILDE MARCOLLI, AND EDUARDO NASCIMENTO

ABSTRACT. We obtain explicit expressions for the class in the Grothendieck group of varieties of the moduli space $\overline{\mathcal{M}}_{0,n}$ of genus 0 stable curves with n marked points. This information is equivalent to the Poincaré polynomial; it implies explicit expressions for the Betti numbers of the moduli space in terms of Stirling numbers or, alternatively, Bernoulli numbers.

The expressions are obtained by solving a differential equation characterizing the generating function for the Grothendieck class as shown in work of Yuri Manin from the 1990s. This differential equation is equivalent to S. Keel's recursion for the Betti numbers of $\overline{\mathcal{M}}_{0,n}$. Our proof reduces the solution to two combinatorial identities which follow from applications of Lagrange series.

We also study generating functions for the individual Betti numbers. In previous work it had been shown that these functions are determined by a set of polynomials $p_m^{(k)}(z)$, $k \geq m$, with positive rational coefficients, which are conjecturally log-concave. We verify this conjecture for many infinite families of polynomials $p_m^{(k)}(z)$, corresponding to the generating functions for the $2k$ -Betti numbers of $\overline{\mathcal{M}}_{0,n}$ for all $k \leq 100$. Further, studying the polynomials $p_m^{(k)}(z)$ allows us to prove that the generating function for the Grothendieck class of $\mathcal{M}_{0,n}$ may be written in terms of a series of rational functions in \mathbb{L} and the principal branch of the *Lambert W-function*.

We include an interpretation of the main result in terms of Stirling matrices and a discussion of the Euler characteristic of $\overline{\mathcal{M}}_{0,n}$.

1. INTRODUCTION

Let $\overline{\mathcal{M}}_{0,n}$ be the moduli space of stable n -pointed curves of genus 0, $n \geq 3$, and denote by $[\overline{\mathcal{M}}_{0,n}]$ its class in the Grothendieck ring of varieties. This class is a polynomial in the Lefschetz-Tate class $\mathbb{L} = [\mathbb{A}^1]$, and in fact

$$[\overline{\mathcal{M}}_{0,n}] = \sum_{k=0}^{n-3} \text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n}) \mathbb{L}^k,$$

see [MM16, Remark 3.2.2]. Thus, this class is a manifestation of the Poincaré polynomial of $\overline{\mathcal{M}}_{0,n}$. The following surprisingly elegant formula is, to our knowledge, new.

Theorem 1.1.

$$(1.1) \quad [\overline{\mathcal{M}}_{0,n}] = (1 - \mathbb{L})^{n-1} \sum_{j \geq 0} \sum_{k \geq 0} s(k+n-1, k+n-1-j) S(k+n-1-j, k+1) \mathbb{L}^{k+j}.$$

Here s , resp., S denote Stirling numbers of the first, resp., second kind. The result implies that the stated expression evaluates to a degree- $(n-3)$ polynomial in \mathbb{L} with positive coefficients, a fact that seems in itself nontrivial.

An explicit formula for the individual Betti numbers is an immediate consequence of Theorem 1.1.

Corollary 1.2. *For $n \geq 3$:*

$$\dim H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \sum_{j=0}^{\ell} \sum_{k=0}^{\ell-j} (-1)^{\ell-j-k} \binom{n-1}{\ell-j-k} s(k+n-1, k+n-1-j) S(k+n-1-j, k+1).$$

For instance,

$$\begin{aligned} \dim H^6(\overline{\mathcal{M}}_{0,5}) &= s(4,1)S(1,1) - 4s(4,2)S(2,1) + 6s(4,3)S(3,1) - 4s(4,4)S(4,1) \\ &\quad + s(5,3)S(3,2) - 4s(5,4)S(4,2) + 6s(5,5)S(5,2) \\ &\quad + s(6,5)S(5,3) - 4s(6,6)S(6,3) + s(7,7)S(7,4) \\ &= (-6) \cdot 1 - 4 \cdot 11 \cdot 1 + 6 \cdot (-6) \cdot 1 - 4 \cdot 1 \cdot 1 + 35 \cdot 3 \\ &\quad - 4 \cdot (-10) \cdot 7 + 6 \cdot 1 \cdot 15 + (-15) \cdot 25 - 4 \cdot 1 \cdot 90 + 1 \cdot 350 \\ &= 0 \end{aligned}$$

while

$$\begin{aligned} \dim H^6(\overline{\mathcal{M}}_{0,10}) &= s(9,6)S(6,1) - 9s(9,7)S(7,1) + 36s(9,8)S(8,1) - 84s(9,9)S(9,1) \\ &\quad + s(10,8)S(8,2) - 9s(10,9)S(9,2) + 36s(10,10)S(10,2) \\ &\quad + s(11,10)S(10,3) - 9s(11,11)S(11,3) + s(12,12)S(12,4) \\ &= (-4536) \cdot 1 - 9 \cdot 546 \cdot 1 + 36 \cdot (-36) \cdot 1 - 84 \cdot 1 \cdot 1 + 870 \cdot 127 \\ &\quad - 9 \cdot (-45) \cdot 255 + 36 \cdot 1 \cdot 511 + (-55) \cdot 9330 - 9 \cdot 1 \cdot 28501 + 1 \cdot 611501 \\ &= 63173. \end{aligned}$$

We prove Theorem 1.1 by studying the generating function

$$(1.2) \quad \widehat{M} := 1 + z + \sum_{n \geq 3} [\overline{\mathcal{M}}_{0,n}] \frac{z^{n-1}}{(n-1)!}.$$

Recursive formulas for the Betti numbers and the Poincaré polynomial of $\overline{\mathcal{M}}_{0,n}$ have been known for three decades, since S. Keel's seminal work [Kee92]. E. Getzler ([Get95]) and Y. Manin ([Man95]) obtained explicit functional and differential equations satisfied by the generating function (1.2), and this information was interpreted in terms of the Grothendieck class of $\overline{\mathcal{M}}_{0,n}$ in [MM16]. We obtain the following explicit expressions for this generating function.

Theorem 1.3.

$$\begin{aligned} \widehat{M} &= \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} \left(((1-\mathbb{L})(1+(z+1)\mathbb{L}))^{\frac{1+\ell}{\mathbb{L}}-\ell} \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) \right) \mathbb{L}^\ell \\ &= \sum_{\ell \geq 0} \left(\sum_{k \geq 0} \frac{(\ell+1)^{\ell+k}}{(\ell+1)!k!} (z-\mathbb{L}-z\mathbb{L})^k \prod_{j=0}^{\ell+k-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) \right) \mathbb{L}^\ell. \end{aligned}$$

Here and in the rest of the paper, expressions with non-integer exponents should be understood as shorthand for the corresponding exponentials: $(\dots)^A := \exp(A \log(\dots))$.

We prove Theorem 1.1 as a direct consequence of the second expression in Theorem 1.3, and the second expression follows from the first. We prove that the first expression equals \widehat{M} by showing that it is the solution of the aforementioned differential equation characterizing \widehat{M} .

The first expression may be used to obtain alternative formulas for $[\overline{\mathcal{M}}_{0,n}]$ and for the Betti numbers.

Corollary 1.4. *For all $n \geq 3$:*

$$(1.3) \quad [\overline{\mathcal{M}}_{0,n}] = \sum_{\ell \geq 0} \frac{(1 - \mathbb{L}^2)^{\frac{\ell+1}{\mathbb{L}} - \ell}}{(1 + \mathbb{L})^{n-1}} \left(\prod_{j=0}^{\ell+n-2} (\ell + 1 - j\mathbb{L}) \right) \frac{\mathbb{L}^\ell}{(\ell + 1)!}.$$

As in the case of (1.1), the right-hand side of (1.3) is, despite appearances, a polynomial in \mathbb{L} with positive integer coefficients and degree $n - 3$. In particular, for every n , only finitely many summands of (1.3) need be computed in order to determine $[\overline{\mathcal{M}}_{0,n}]$. For example, here are the series expansions of the first few summands of (1.3), for $n = 6$:

$$\begin{aligned} \ell = 0 : \quad & 1 - 16\mathbb{L} + \frac{231}{2}\mathbb{L}^2 - \frac{3109}{6}\mathbb{L}^3 + \frac{40549}{24}\mathbb{L}^4 - \frac{265223}{60}\mathbb{L}^5 + \frac{7126141}{720}\mathbb{L}^6 + \dots \\ \ell = 1 : \quad & 32\mathbb{L} - 464\mathbb{L}^2 + 3256\mathbb{L}^3 - \frac{45326}{3}\mathbb{L}^4 + \frac{158768}{3}\mathbb{L}^5 - \frac{2288308}{15}\mathbb{L}^6 + \dots \\ \ell = 2 : \quad & \frac{729}{2}\mathbb{L}^2 - \frac{10935}{2}\mathbb{L}^3 + \frac{163215}{4}\mathbb{L}^4 - \frac{410265}{2}\mathbb{L}^5 + \frac{12663747}{16}\mathbb{L}^6 + \dots \\ \ell = 3 : \quad & \frac{8192}{3}\mathbb{L}^3 - \frac{131072}{3}\mathbb{L}^4 + \frac{1057792}{3}\mathbb{L}^5 - \frac{17410048}{9}\mathbb{L}^6 + \dots \\ \ell = 4 : \quad & \frac{390625}{24}\mathbb{L}^4 - \frac{3359375}{12}\mathbb{L}^5 + \frac{117453125}{48}\mathbb{L}^6 + \dots \\ \ell = 5 : \quad & \frac{419904}{5}\mathbb{L}^5 - \frac{7768224}{5}\mathbb{L}^6 + \dots \\ \ell = 6 : \quad & \frac{282475249}{720}\mathbb{L}^6 + \dots \end{aligned}$$

and their sum:

$$1 + 16\mathbb{L} + 16\mathbb{L}^2 + \mathbb{L}^3 + 0\mathbb{L}^4 + 0\mathbb{L}^5 + 0\mathbb{L}^6 + \dots$$

The infinite sum (1.3) converges to $[\overline{\mathcal{M}}_{0,6}] = 1 + 16\mathbb{L} + 16\mathbb{L}^2 + \mathbb{L}^3$ in the evident sense. Concerning Betti numbers, we have the following alternative to Corollary 1.2.

Corollary 1.5. *For $i > 0$, let*

$$C_{nki} = \frac{(-1)^i(2ki + ni + k + n - 1) + k - i}{i(i+1)} - \frac{1}{i(k+1)^i} \sum_{j=0}^{k+n-2} j^i.$$

Then for all $\ell \geq 0$ and all $n \geq 3$ we have

$$(1.4) \quad \text{rk } H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \sum_{k=0}^{\ell} \frac{(k+1)^{k+n-1}}{(k+1)!} \sum_{m=0}^{\ell-k} \frac{1}{m!} \sum_{i_1 + \dots + i_m = \ell-k} C_{nki_1} \dots C_{nki_m}.$$

By Faulhaber's formula, the numbers C_{nki} may be written in terms of Bernoulli numbers:

$$\begin{aligned} C_{nki} = \frac{1}{i(i+1)} & \left((-1)^i(2ki + ni + k + n - 1) + k - i \right. \\ & \left. - \frac{1}{(k+1)^i} \sum_{j=0}^i \binom{i+1}{j} B_j (k+n-1)^{i-j+1} \right). \end{aligned}$$

Thus, Corollary 1.5 expresses the Betti numbers of $\overline{\mathcal{M}}_{0,n}$ as certain combinations of Bernoulli numbers, just as Corollary 1.2 expresses the same in terms of Stirling numbers. For instance,

$$\begin{aligned} \text{rk } H^6(\overline{\mathcal{M}}_{0,5}) &= \frac{271}{2} - \frac{889}{3}B_0 - \frac{277}{3}B_1 - 8B_2 - \frac{4}{3}B_3 \\ &\quad + \frac{1427}{6}B_0^2 + \frac{590}{3}B_0B_1 + 16B_0B_2 + 42B_1^2 + 8B_1B_2 \\ &\quad - \frac{256}{3}B_0^3 - 128B_0^2B_1 - 64B_0B_1^2 - \frac{32}{3}B_1^3 \\ &= 0 \end{aligned}$$

while

$$\begin{aligned} \text{rk } H^6(\overline{\mathcal{M}}_{0,10}) &= \frac{756667}{2} - \frac{5729797}{12}B_0 - 86825B_1 - \frac{1271}{2}B_2 - 3B_3 \\ &\quad + \frac{313439}{2}B_0^2 + \frac{252355}{4}B_0B_1 + \frac{729}{4}B_0B_2 + \frac{12719}{2}B_1^2 + \frac{81}{2}B_1B_2 \\ &\quad - \frac{177147}{16}B_0^3 - \frac{59049}{8}B_0^2B_1 - \frac{6561}{4}B_0B_1^2 - \frac{243}{2}B_1^3 \\ &= 63173. \end{aligned}$$

For $n = 1, 2$, the right-hand side of (1.4) equals 1 for $k = 0$ and 0 for $k > 0$. For every $n \geq 1$, (1.4) may be viewed as an infinite collection of identities involving Bernoulli numbers, equivalent to the corresponding identities involving Stirling numbers arising from Corollary 1.2. It appears to be useful to have explicit expressions of both types. For instance, Corollary 1.5 can be used to refine the result obtained in [ACM25] on asymptotic (ultra-)log-concavity of the Grothendieck class of $\overline{\mathcal{M}}_{0,n}$, see [Nas].

Keel's recurrence relation for the Betti numbers of $\overline{\mathcal{M}}_{0,n}$ can itself be viewed as a sophisticated identity of Stirling numbers (by Corollary 1.2) or Bernoulli numbers (by Corollary 1.5). Verifying such identities directly would provide a more transparent proof of these results, but the combinatorics needed for this verification seems substantially more involved than the somewhat indirect way we present in this paper.

The proof of Theorem 1.3 is presented in §2. It relies crucially on two combinatorial identities, both of which follow from ‘Lagrange inversion’. Proofs of these identities are given in an appendix. Theorem 1.1 and the other corollaries stated above are proved in §3.

In §§4–6 we study the finer structure of the formulas obtained in the first part of the paper, building upon work carried out in [ACM25]. This information is both interesting in itself and leads to further results on the Betti numbers of $\overline{\mathcal{M}}_{0,n}$ and on the generating function \widehat{M} , see Theorem 1.9 and 1.11 below. As in [ACM25], we consider the generating function

$$\alpha_k(z) = \sum_{n \geq 3} \text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n}) \frac{z^{n-1}}{(n-1)!}$$

for the coefficients of \mathbb{L}^k in $[\overline{\mathcal{M}}_{0,n}]$, i.e., the individual Betti numbers of $\overline{\mathcal{M}}_{0,n}$. While Corollary 1.2 yields an explicit expression for the coefficients of this generating function, there is an interesting structure associated with $\alpha_k(z)$ that is not immediately accessible from such an expression. Specifically, by [ACM25, Theorem 4.1] there exist polynomials $p_m^{(k)}(z) \in \mathbb{Q}[z]$, $0 \leq m \leq k$, such that

$$(1.5) \quad \alpha_k(z) = e^z \sum_{m=0}^k (-1)^m p_m^{(k)}(z) e^{(k-m)z}$$

for all $k > 0$, while $\alpha_0(z) = e^z - z - 1$. It is proved in [ACM25] that $p_m^{(k)}(z)$ has degree $2m$ and positive leading coefficient, and that

$$(1.6) \quad p_0^{(k)} = \frac{(k+1)^k}{(k+1)!},$$

and this is used to establish an asymptotic form of log-concavity of $[\overline{\mathcal{M}}_{0,n}]$. Several polynomials $p_m^{(k)}(z)$ are computed explicitly in [ACM25]; for instance,

$$p_1^{(1)}(z) = 1 + z + \frac{z^2}{2},$$

so that

$$\alpha_1(z) = e^z \left(p_0^{(1)} e^z - p_1^{(1)}(z) \right) = e^z \left(e^z - 1 - z - \frac{z^2}{2} \right) = \frac{z^3}{3!} + 5 \frac{z^4}{4!} + 16 \frac{z^5}{5!} + 42 \frac{z^6}{6!} + \dots$$

is the generating function for $\text{rk } H^2(\overline{\mathcal{M}}_{0,n})$. On the basis of extensive computations, in [ACM25] we proposed the following.

Conjecture 1. *For all $k \geq 1$, the polynomials $p_m^{(k)}$, $m = 1, \dots, k$, have positive coefficients and are log-concave with no internal zeros. All but $p_1^{(1)}$, $p_3^{(3)}$, $p_5^{(5)}$ are ultra-log-concave.*

In this paper we prove the first statement in this conjecture and provide substantial numerical evidence for the second part. For this purpose, we assemble the polynomials $p_m^{(k)}(z)$ in the generating function

$$P(z, t, u) := \sum_{m \geq 0} \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) t^\ell u^m.$$

By (1.5), $\widehat{M}(z, \mathbb{L})$ is (up to an exponential factor) a specialization of $P(z, t, u)$. This fact and Theorem 1.3 may be used to obtain an explicit expression for $P(z, t, u)$: we prove the following statement in §4.

Theorem 1.6.

$$P(z, t, u) = \sum_{\ell \geq 0} \left(\frac{(\ell+1)^\ell}{(\ell+1)!} e^{-(\ell+1)z} ((1+u)(1-u(z+1)))^{-\frac{1+\ell(u+1)}{u}} \prod_{j=0}^{\ell-1} \left(1 + \frac{ju}{\ell+1} \right) \right) t^\ell.$$

In §5 we use Theorem 1.6 to obtain more information on the polynomials $p_m^{(k)}(z)$. Denote by $c_{mj}^{(k)}$ the coefficients of $p_m^{(k)}(z)$, so that $p_m^{(k)}(z) = \sum_{j=0}^{2m} c_{mj}^{(k)} z^j$.

Theorem 1.7. *For all $m \geq 0$, $0 \leq j \leq 2m$, there exist polynomials $\Gamma_{mj}(\ell) \in \mathbb{Q}[\ell]$ of degree $2m-j$ such that*

$$(1.7) \quad c_{mj}^{(k)} = \frac{(k-m+1)^{k-2m+j}}{(k-m+1)!} \cdot \Gamma_{mj}(k-m)$$

for all $k \geq m$. Further $\Gamma_{mj}(\ell) > 0$ for all $m \geq 0$, $0 \leq j \leq 2m$, $\ell \geq 0$.

The polynomial $\Gamma_{mj}(\ell)$ are effectively computable. For example,

$$\Gamma_{4,0} = \frac{27}{128} \ell^8 + \frac{57}{32} \ell^7 + \frac{4295}{576} \ell^6 + \frac{1341}{80} \ell^5 + \frac{28867}{1152} \ell^4 + \frac{2143}{96} \ell^3 + \frac{3619}{288} \ell^2 + \frac{119}{30} \ell + \frac{13}{24}.$$

The first 400 such polynomials $\Gamma_{mj}(\ell)$ with $0 \leq m < 20$, $0 \leq j \leq 2m$ have positive coefficients and are in fact log concave. However, $\Gamma_{20,0}$ is not log concave, and the coefficient of ℓ^2 in the degree-42 polynomial $\Gamma_{21,0}$, namely, $-\frac{97330536888617758406393}{224800145555215360000}$, is *negative*, a good

reminder of how delicate these notions are and a cautionary tale about making premature conjectures.

Nevertheless, as stated in Theorem 1.7, we can prove that $\Gamma_{mj}(\ell)$ is positive for all m , $0 \leq j \leq 2m$, $\ell \geq 0$, and this has the following immediate consequence.

Corollary 1.8. *The polynomials $p_m^{(k)}(z)$ have positive coefficients.*

This proves part of Conjecture 1. Further, we obtain substantial evidence for the rest of the conjecture, dealing with log-concavity of the polynomials $p_m^{(k)}(z)$. Specifically, we reduce the proof of ultra-log-concavity of $p_m^{(k)}(z)$ for a fixed m and all $k \geq m$ to a finite computation involving the polynomials Γ_{mj} . A few hours of computing time verified the conjecture for $m = 1, \dots, 100$.

As a byproduct of these considerations, we obtain an alternative expression for the Betti numbers of $\overline{\mathcal{M}}_{0,n}$, in terms of the polynomials $\Gamma_{mj}(\ell)$.

Theorem 1.9. *For $n \geq 3$ and $0 \leq \ell \leq n-3$:*

$$\text{rk } H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \sum_{k+m=\ell} (-1)^m \frac{(k+1)^{n-2+k-m}}{k!} \sum_{j=0}^{2m} (n-1) \cdots (n-j) \Gamma_{mj}(k).$$

This formula generalizes directly the well-known formula for the second Betti number,

$$\text{rk } H^2(\overline{\mathcal{M}}_{0,n}) = \frac{1}{2} \cdot 2^n - \frac{n^2 - n + 2}{2},$$

cf. [Kee92, p. 550]. For instance,

$$\begin{aligned} \text{rk } H^6(\overline{\mathcal{M}}_{0,n}) &= \frac{2}{3} \cdot 4^n - \frac{(n+4)(n+3)}{12} \cdot 3^n + \frac{3n^4 + 14n^3 + 57n^2 + 118n + 96}{192} \cdot 2^n \\ &\quad - \frac{n^6 - 7n^5 + 35n^4 - 77n^3 + 120n^2 - 72n + 32}{48}, \end{aligned}$$

giving

$$\text{rk } H^6(\overline{\mathcal{M}}_{0,10}) = \frac{2}{3} \cdot 4^{10} - \frac{91}{6} \cdot 3^{10} + \frac{531}{2} \cdot 2^{10} - \frac{73039}{2} = 63173.$$

Studying the function $P(z, t, u)$ also reveals an intriguing connection with the *Lambert W-function*, which we explore in §6. Recall that the Lambert W-function $W(t)$ is characterized by the identity $W(t)e^{W(t)} = t$; the reader is addressed to [CGH⁺96] for a detailed treatment of this function. We consider in particular the ‘tree function’ $T(t) = -W(-t)$, where W is the principal branch of the Lambert W-function. (This function owes its name to the fact that $T(t) = \sum_{n \geq 1} \frac{T_n t^n}{n!}$, where $T_n = n^{n-1}$ is the number of rooted trees on n labelled vertices.) For a fixed m , consider the generating function

$$(1.8) \quad P_m(z, t) = \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) t^\ell,$$

that is, the coefficient of u^m in P . This is a polynomial in $\mathbb{Q}[[t]][z]$ of degree $2m$. Explicit computations show that

$$\begin{aligned} P_0 &= e^T \\ P_1 &= \frac{e^T}{(1-T)} \left(\frac{1}{2}z^2 + (1+T)z + \frac{1}{2}(2+T^2) \right) \end{aligned}$$

and

$$P_2 = \frac{e^T}{(1-T)^3} \left(\frac{1}{8} z^4 + \frac{5+2T-T^2}{6} z^3 + \frac{8+4T+T^2-2T^3}{4} z^2 + \frac{4+2T+T^2-T^4}{2} z + \frac{12+24T-12T^2+8T^3-T^4-4T^5}{24} \right)$$

where $T = T(t)$ is the tree function. (In fact, the first expression is a restatement of (1.6).) For all $m \geq 0$, the series $e^{-T} P_m(z, t)$ is a polynomial in z with coefficients in $\mathbb{Q}[[t]]$. We prove that for all $m \geq 0$ these coefficients can be expressed as *rational functions* in the tree function T , as in the examples shown above. More precisely,

Proposition 1.10. *Let $T = T(t) = -W(-t)$ be the tree function. For $m > 0$, there exist polynomials $F_m(z, \tau) \in \mathbb{Q}[z, \tau]$, of degree $2m$ in z and $< 3m$ in τ , such that*

$$(1.9) \quad P_m(z, t) = e^T \frac{F_m(z, T)}{(1-T)^{2m-1}}.$$

Setting $F_0(z, T) = \frac{1}{1-T}$ extends the validity of (1.9) to the case $m = 0$.

Since for $m > 0$ the polynomials $F_m(z, \tau)$ have degree $< 3m$ in τ , they are characterized by the congruence

$$F_m(z, \tau) \equiv \sum_{\ell=0}^{3m-1} p_m^{(m+\ell)}(z) (1-\tau)^{2m-1} e^{-(\ell+1)\tau} \tau^\ell \pmod{\tau^{3m}}.$$

Proposition 1.10 implies the following alternative expression for \widehat{M} .

Theorem 1.11. *Let $T = T(e^z \mathbb{L})$, where T is the tree function. Then with $F_m(z, \tau)$ as above we have*

$$(1.10) \quad \widehat{M} = \frac{T}{\mathbb{L}} \sum_{m \geq 0} \frac{(-1)^m F_m(z, T)}{(1-T)^{2m-1}} \mathbb{L}^m.$$

While the identities stated in Theorem 1.3 are more explicit expressions for \widehat{M} , they show no direct trace of the relation with the principal branch of the Lambert W-function displayed in Theorem 1.11. The proof of this result relies on another combinatorial identity involving Stirling numbers, whose proof is also given in the appendix.

It is essentially evident that (1.1) may be expressed in terms of a product of matrices. We formalize this remark in §7, by showing that the class $[\overline{\mathcal{M}}_{0,n}]$ may be recovered as a generalized trace of a matrix obtained in a very simple fashion from the standard matrices defined by Stirling numbers of first and second kind (Theorem 7.1). This is simply a restatement of Theorem 1.1, but it may help to relate the results of this paper with the extensive literature on Stirling numbers.

There is another connection between $\overline{\mathcal{M}}_{0,n}$ and the Lambert W-function: the generating function for the Euler characteristic of $\overline{\mathcal{M}}_{0,n}$ may be expressed in terms of the other real branch of the Lambert W-function, denoted W_{-1} in [CGH⁺96]. The various expressions obtained in this paper for the Betti numbers imply explicit formulas for the individual Euler characteristics $\chi(\overline{\mathcal{M}}_{0,n})$. In §8 we highlight one compelling appearance of $\chi(\overline{\mathcal{M}}_{0,n})$ as the leading coefficient of a polynomial determined by a sum of products of Stirling numbers, see Proposition 8.1.

Choi, Kiem, and Lee have extensively studied the S_n -representation structure of $H_*(\overline{\mathcal{M}}_{0,n})$ ([CKL23, CKL, CKL25]), recovering in particular asymptotic results from [ACM25]. Stirling numbers are closely related to the combinatorics of S_n , and it would be interesting to investigate whether the combinatorial formulas in this paper can be understood directly in terms of representations of symmetric groups.

A new proof of the Stirling formula for the Poincaré polynomial of $\overline{\mathcal{M}}_{0,n}$ was recently obtained in [EFM⁺], in the context of a general study of Chow rings of polymatroids with respect to arbitrary building sets.

We end this introduction by pointing out that another expression for $[\overline{\mathcal{M}}_{0,n}]$ may be obtained as a consequence of the result of Ezra Getzler mentioned earlier, which we reproduce as follows. For consistency with the notation used above, we interpret Getzler's formula with $\mathbb{L} = t^2$.

Theorem 1.12 (Getzler [Get95]). *Let*

$$g(x, \mathbb{L}) = x - \sum_{n=2}^{\infty} \frac{x^n}{n!} \sum_{i=0}^{n-2} (-1)^i \mathbb{L}^{(n-i-2)} \dim H_i(\mathcal{M}_{0,n+1}) = x - \frac{(1+x)^{\mathbb{L}} - (1+\mathbb{L}x)}{\mathbb{L}(\mathbb{L}-1)}.$$

Then

$$f(x, \mathbb{L}) := x + \sum_{n=2}^{\infty} \frac{x^n}{n!} \sum_{i=0}^{n-2} \mathbb{L}^i \dim H_{2i}(\overline{\mathcal{M}}_{0,n+1})$$

is the inverse of g , in the sense that $f(g(x, \mathbb{L}), \mathbb{L}) = x$.

Getzler's generating function f differs from our \widehat{M} by the absence of the constant term. Theorem 1.12 may be seen to be equivalent to the functional equation for the Poincaré polynomial mentioned earlier, also proved (with different notation) in [Man95].

We can apply Lagrange inversion (see e.g., the first formula in [Rio68, p. 146, (18)]) to the identity $f(g(x, \mathbb{L}), \mathbb{L}) = x$. Specifically, assume that

$$f(y) = \sum_{n \geq 0} a_n \frac{y^n}{n!}$$

is a power series such that

$$\alpha(x) = f(\beta(x)) = \sum_{n \geq 0} a_n \frac{\beta(x)^n}{n!}$$

for constants a_n and functions $\alpha(x)$, $\beta(x)$ such that $\beta(0) = 0$. Then Lagrange inversion states that

$$a_n = \frac{d^{n-1}}{dx^{n-1}} (\alpha'(x) \varphi(x)^n) |_{x=0},$$

where $\varphi(x) = \frac{x}{\beta(x)}$. Applying this formula with $a_n = [\overline{\mathcal{M}}_{0,n+1}]$, $\alpha(x) = x$, $\beta(x) = g(x, \mathbb{L})$ gives

$$[\overline{\mathcal{M}}_{0,n+1}] = (n-1)! \cdot \text{coefficient of } x^{n-1} \text{ in the expansion of } \left(\frac{x}{g(x, \mathbb{L})} \right)^n$$

with $g(x, \mathbb{L})$ as in Theorem 1.12. Therefore,

$$[\overline{\mathcal{M}}_{0,n}] = (n-2)! \cdot \text{coefficient of } x^{n-2} \text{ in the expansion of } \left(\frac{\mathbb{L}(\mathbb{L}-1)x}{1 + \mathbb{L}^2 x - (1+x)^{\mathbb{L}}} \right)^{n-1}.$$

This also easily implies that

$$\widehat{\chi}(\overline{\mathcal{M}}_{0,n}) = (n-2)! \cdot \text{coefficient of } x^{n-2} \text{ in the expansion of } \left(\frac{1}{2 - \frac{1+x}{x} \log(1+x)} \right)^{n-1}.$$

We do not know if alternative proofs of the results in this paper may be obtained from these expressions.

Acknowledgments. P.A. was supported in part by the Simons Foundation, collaboration grant #625561, and thanks Caltech for the hospitality. M.M. was supported by NSF grant DMS-2104330. E.N. was supported by a Summer Undergraduate Research Fellowship at Caltech funded in part by a generous gift from Marcella Bonsall.

The authors thank the referee for the careful reading of the paper and the good suggestions for improvements.

2. PROOF OF THEOREM 1.3

Throughout the paper, all functions are implicitly considered as formal power series, used as generating functions for the coefficients in their expansion.

A recursion determining $[\overline{\mathcal{M}}_{0,n}]$ or, equivalently, the Poincaré polynomial of $\overline{\mathcal{M}}_{0,n}$, is well-known: see [Kee92], [Man95], [MM16]. It is equivalent to a differential equation for the generating function

$$\widehat{M}(z, \mathbb{L}) := 1 + z + \sum_{n \geq 3} [\overline{\mathcal{M}}_{0,n}] \frac{z^{n-1}}{(n-1)!},$$

namely,

$$(2.1) \quad \frac{\partial \widehat{M}}{\partial z} = \frac{\widehat{M}}{1 + \mathbb{L}(1+z) - \mathbb{L}\widehat{M}},$$

subject to the initial condition $\widehat{M}|_{z=0} = 1$. (This is a restatement of [Man95, (0.8)].)

Denote by \overline{M} the first expression stated in Theorem 1.3:

$$(2.2) \quad \overline{M} := \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} \left(\left((1-\mathbb{L})(1+(z+1)\mathbb{L}) \right)^{\frac{\ell+1}{\mathbb{L}}-\ell} \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^\ell,$$

where we view \mathbb{L} as an indeterminate. We need to verify that $\widehat{M} = \overline{M}$. To prove this result, it suffices to verify that \overline{M} satisfies (2.1), which we rewrite as

$$(2.3) \quad (1 + \mathbb{L}(1+z)) \frac{\partial \overline{M}}{\partial z} - \overline{M} = \mathbb{L} \frac{\partial \overline{M}}{\partial z} \overline{M},$$

and the initial condition $\overline{M}|_{z=0} = 1$.

Both of these claims will be reduced to combinatorial identities. For the first, it is convenient to separate the part of \overline{M} depending on z from the rest: write

$$\overline{M} = \sum_{k=0}^{\infty} F(k) (1 + (z+1)\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}}$$

with

$$F(k) := \frac{(k+1)^k}{(k+1)!} \left(\prod_{j=0}^{k-1} \left(1 - \frac{j\mathbb{L}}{k+1} \right) \right) (1-\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}} \mathbb{L}^k.$$

We have

$$\frac{\partial \bar{M}}{\partial z} = \sum_{k=0}^{\infty} F(k)(1 + (z+1)\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}} \frac{1 + k(1-\mathbb{L})}{(1 + (z+1)\mathbb{L})},$$

implying

$$(1 + (z+1)\mathbb{L}) \frac{\partial \bar{M}}{\partial z} - \bar{M} = \sum_{k=1}^{\infty} F(k)k(1-\mathbb{L})(1 + (z+1)\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}}.$$

In order to verify (2.3), we need to verify that this expression equals

$$\begin{aligned} \mathbb{L} \frac{\partial \bar{M}}{\partial z} \bar{M} &= \mathbb{L} \sum_{\ell, m} F(\ell)(1 + (z+1)\mathbb{L})^{\frac{1+\ell(1-\mathbb{L})}{\mathbb{L}}} \frac{1 + \ell(1-\mathbb{L})}{(1 + (z+1)\mathbb{L})} F(m)(1 + (z+1)\mathbb{L})^{\frac{1+m(1-\mathbb{L})}{\mathbb{L}}} \\ &= \mathbb{L} \sum_{\ell, m} F(\ell)F(m)(1 + \ell(1-\mathbb{L}))(1 + (z+1)\mathbb{L})^{\frac{2+(\ell+m)(1-\mathbb{L})}{\mathbb{L}}-1} \\ &= \mathbb{L} \sum_{k \geq 1} \sum_{\ell+m=k-1} F(\ell)F(m)(1 + \ell(1-\mathbb{L}))(1 + (z+1)\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}}. \end{aligned}$$

Therefore, In order to prove that \bar{M} satisfies (2.3), it suffices to prove that for all $k \geq 1$

$$\mathbb{L} \sum_{\ell+m=k-1} F(\ell)F(m) \cdot (1 + \ell(1-\mathbb{L})) = F(k) \cdot k(1-\mathbb{L}).$$

To simplify this further, let

$$E(\ell) := \frac{(\ell+1)^\ell}{(\ell+1)!} \cdot \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right),$$

so that

$$F(\ell) = E(\ell) \cdot (1-\mathbb{L})^{\frac{1+\ell(1-\mathbb{L})}{\mathbb{L}}} \mathbb{L}^\ell.$$

The sought-for identity is

$$\begin{aligned} \mathbb{L} \sum_{\ell+m=k-1} E(\ell) \cdot (1-\mathbb{L})^{\frac{1+\ell(1-\mathbb{L})}{\mathbb{L}}} \mathbb{L}^\ell E(m) \cdot (1-\mathbb{L})^{\frac{1+m(1-\mathbb{L})}{\mathbb{L}}} \mathbb{L}^m \cdot (1 + \ell(1-\mathbb{L})) \\ = E(k) \cdot (1-\mathbb{L})^{\frac{1+k(1-\mathbb{L})}{\mathbb{L}}} \mathbb{L}^k \cdot k(1-\mathbb{L}). \end{aligned}$$

Clearing the common factor proves the following.

Claim 2.1. *Let*

$$E(\ell) := \frac{(\ell+1)^\ell}{(\ell+1)!} \cdot \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right).$$

Then in order to prove that \bar{M} satisfies (2.3), it suffices to prove that for all $k \geq 1$,

$$(2.4) \quad \sum_{\ell+m=k-1} E(\ell)E(m) \cdot (1 + \ell(1-\mathbb{L})) = k \cdot E(k).$$

For every k , this is an identity of polynomials in \mathbb{L} . For example, for $k=3$ it states that

$$\left(\frac{3}{2} - \frac{\mathbb{L}}{2}\right)(3-2\mathbb{L}) + (2-\mathbb{L}) + \left(\frac{3}{2} - \frac{\mathbb{L}}{2}\right) = 8 \left(1 - \frac{\mathbb{L}}{4}\right) \left(1 - \frac{\mathbb{L}}{2}\right).$$

As such, (2.4) is equivalent to the identity obtained by performing an invertible change of variables. Setting $\mathbb{L} = -\frac{1}{w-1}$,

$$\begin{aligned} E(\ell) &= \frac{1}{(\ell+1)!} \prod_{j=0}^{\ell-1} (\ell+1-j\mathbb{L}) = \frac{1}{(\ell+1)!} \prod_{j=0}^{\ell-1} \frac{(w-1)(\ell+1)+j}{w-1} \\ &= \frac{\prod_{j=0}^{\ell-1} ((w-1)(\ell+1)+j)}{(\ell+1)!(w-1)^\ell}, \end{aligned}$$

and trivial manipulations show that (2.4) is then equivalent to the identity

$$\begin{aligned} \sum_{\ell+m=k-1} \frac{\prod_{j=1}^{\ell} ((w-1)(\ell+1)+j)}{\ell!} \cdot \frac{\prod_{j=0}^{m-1} ((w-1)(m+1)+j)}{m!} \cdot \frac{1}{m+1} \\ = \frac{\prod_{j=1}^{k-1} ((w-1)(k+1)+j)}{(k-1)!} \end{aligned}$$

of polynomials in $\mathbb{Q}[w]$. In order to verify this identity, it suffices to verify that it holds when w is evaluated at infinitely many integers. We can then restate Claim 2.1 as follows.

Claim 2.2. *In order to verify that \overline{M} satisfies (2.3), it suffices to prove the following binomial identity*

$$(2.5) \quad \sum_{\ell+m=k-1} \binom{w(\ell+1)-1}{\ell} \cdot \binom{w(m+1)-2}{m} \cdot \frac{1}{m+1} = \binom{w(k+1)-2}{k-1}$$

for all positive integers k and w .

This identity indeed does hold. It may be obtained as a specialization of more general, known, identities; see e.g., [Rio68, p. 169] or [Gou56], where such identities are identified as generalizations of ‘Vandermonde’s convolution’. We include a proof of (2.5) in the appendix, Lemma A.1, and this concludes the verification that \overline{M} satisfies the differential equation (2.3).

Next, we need to verify that \overline{M} satisfies the same initial condition as \widehat{M} , i.e., $\overline{M}|_{z=0} = 1$, that is,

$$(2.6) \quad \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} (1-\mathbb{L}^2)^{\frac{1+\ell}{\mathbb{L}}-\ell} \left(\prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^\ell = 1.$$

This statement is surprisingly nontrivial.

Remark 2.3. Identity (2.6) may be viewed as an infinite collection of identities involving Bernoulli numbers. Explicitly, for $k \geq 0$ and $i \geq 1$ let

$$A_{ki} := \begin{cases} -\frac{2(k+1)}{i+1} - \frac{1}{i(i+1)(k+1)^i} \sum_{j=0}^i \binom{i+1}{j} k^{i-j+1} B_j & i \text{ odd} > 0 \\ \frac{2k}{i} - \frac{1}{i(i+1)(k+1)^i} \sum_{j=0}^i \binom{i+1}{j} k^{i-j+1} B_j & i \text{ even} > 0. \end{cases}$$

Identity (2.6) is then equivalent to the assertion that for all positive integers ℓ

$$\sum_{k=0}^{\ell} \frac{(k+1)^k}{(k+1)!} \sum_{m \geq 0} \frac{1}{m!} \sum_{i_1+\dots+i_m=\ell-k} A_{ki_1} \cdots A_{ki_m} = 0$$

where all indices i_j in the summation are positive integers. (This computation will be carried out in a more general setting in the proof of Corollary 1.5.) For increasing values of ℓ , this identity states that

$$\begin{aligned} B_1 &= -\frac{1}{2}B_0 \\ B_2 &= 4 - \frac{13}{3}B_0 - B_1 + \frac{1}{4}(B_0 + 2B_1)^2 \\ B_3 &= 12 - \frac{259}{12}B_0 - 15B_1 + \frac{1}{2}B_2 + \frac{27}{4}B_0^2 + \frac{45}{4}B_0B_1 \\ &\quad + \frac{3}{4}B_0B_2 + \frac{7}{2}B_1^2 + \frac{3}{2}B_1B_2 - \frac{1}{16}(B_0 + 2B_1)^3 \end{aligned}$$

etc. Our verification of (2.6) proves all these identities simultaneously. \square

For increasing values of ℓ , the summands in the left-hand side of (2.6) expand to

$$\begin{aligned} \ell = 0 : \quad (1 - \mathbb{L}^2)^{\frac{1}{\mathbb{L}}} &= 1 - \mathbb{L} + \frac{1}{2}\mathbb{L}^2 - \frac{2}{3}\mathbb{L}^3 + \frac{13}{24}\mathbb{L}^4 + \dots \\ \ell = 1 : \quad (1 - \mathbb{L}^2)^{\frac{2}{\mathbb{L}}-1}\mathbb{L} &= \mathbb{L} - 2\mathbb{L}^2 + 3\mathbb{L}^3 - \frac{13}{3}\mathbb{L}^4 + \dots \\ \ell = 2 : \quad (1 - \mathbb{L}^2)^{\frac{3}{\mathbb{L}}-2}\frac{3-\mathbb{L}}{2}\mathbb{L}^2 &= \frac{3}{2}\mathbb{L}^2 - 5\mathbb{L}^3 + \frac{45}{4}\mathbb{L}^4 + \dots \\ \ell = 3 : \quad (1 - \mathbb{L}^2)^{\frac{4}{\mathbb{L}}-3}\frac{4-\mathbb{L}}{3}\frac{4-2\mathbb{L}}{2}\mathbb{L}^3 &= \frac{8}{3}\mathbb{L}^3 - \frac{38}{3}\mathbb{L}^4 + \dots \end{aligned}$$

The sum of these terms is a power series, and the task is to verify that this series is the constant 1. For this, it suffices to prove that the series has a limit of 1 for $\mathbb{L} = \frac{1}{m}$ for all integers $m > 1$. Setting $\mathbb{L} = \frac{1}{m}$ with $m > 1$ an integer, the left-hand side of (2.6) may be written

$$\begin{aligned} \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} \left(1 - \frac{1}{m^2}\right)^{(\ell+1)m-\ell} \cdot \prod_{j=0}^{\ell-1} \left(1 - \frac{j}{m(\ell+1)}\right) \frac{1}{m^\ell} \\ = \left(1 - \frac{1}{m^2}\right)^m \sum_{\ell \geq 0} \left(\frac{(m^2-1)^{m-1}}{m^{2m}}\right)^\ell \binom{m(\ell+1)}{\ell} \frac{1}{\ell+1} \end{aligned}$$

after elementary manipulations. Therefore, the following assertion holds.

Claim 2.4. *In order to verify (2.6), it suffices to verify that the identity*

$$(2.7) \quad \sum_{\ell \geq 0} \frac{1}{\ell+1} \binom{m(\ell+1)}{\ell} \left(\frac{(m^2-1)^{m-1}}{m^{2m}}\right)^\ell = \left(1 - \frac{1}{m^2}\right)^{-m}$$

holds for all integers $m > 1$.

Identity (2.7) follows from Lemma A.2, which states

$$x^\alpha = \sum_{\ell \geq 0} \frac{\alpha}{\alpha + \ell\beta} \binom{\alpha + \ell\beta}{\ell} y^\ell$$

for all positive integers α, β , with $y = (x-1)x^{-\beta}$. Indeed, setting $\alpha = \beta = m$, this identity gives

$$x^m = \sum_{\ell \geq 0} \frac{1}{\ell+1} \binom{m(\ell+1)}{\ell} y^\ell,$$

and specializing to $x = (1 - \frac{1}{m^2})^{-1}$ yields (2.7) as needed. This concludes the proof of (2.6).

Summarizing, we have proved that \overline{M} and \widehat{M} satisfy the same differential equation and initial conditions. Therefore $\widehat{M} = \overline{M}$, confirming the first assertion in Theorem 1.3. \square

In order to complete the proof of Theorem 1.3, we need to verify that \overline{M} also equals the second expression in that statement:

$$(2.8) \quad \overline{M} = \sum_{\ell \geq 0} \left(\sum_{k \geq 0} \frac{(\ell+1)^{\ell+k}}{(\ell+1)!k!} (z - \mathbb{L} - z\mathbb{L})^k \prod_{j=0}^{\ell+k-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) \right) \mathbb{L}^\ell.$$

For this, use the expansion

$$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2} x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} x^3 + \dots$$

to obtain

$$\begin{aligned} ((1-\mathbb{L})(1+(z+1)\mathbb{L}))^{\frac{\ell+1}{\mathbb{L}}-\ell} &= (1+\mathbb{L}(z-\mathbb{L}-\mathbb{L}z))^{\frac{\ell+1}{\mathbb{L}}-\ell} \\ &= \sum_{k \geq 0} \frac{1}{k!} (z-\mathbb{L}-\mathbb{L}z)^k \prod_{j=0}^{k-1} ((1+\ell) - (\ell+j)\mathbb{L}) \\ &= \sum_{k \geq 0} \frac{(\ell+1)^k}{k!} (z-\mathbb{L}-\mathbb{L}z)^k \prod_{j=0}^{k-1} \left(1 - \frac{(\ell+j)\mathbb{L}}{\ell+1}\right). \end{aligned}$$

Making use of this expression in (2.2) yields

$$\overline{M} = \sum_{\ell \geq 0} \left(\sum_{k \geq 0} \frac{(\ell+1)^{\ell+k}}{(\ell+1)!k!} (z(1-\mathbb{L}) - \mathbb{L})^k \prod_{j=0}^{k-1} \left(1 - \frac{(\ell+j)\mathbb{L}}{\ell+1}\right) \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) \right) \mathbb{L}^\ell$$

and (2.8) follows. \square

Remark 2.5. An alternative proof of Theorem 1.3 can be obtained by using a different characterization of the generating function \widehat{M} . Manin ([Man95, Theorem 0.3.1, (0.7)]) and Getzler ([Get95, Theorem 5.9]) prove that \widehat{M} is the only solution of the functional equation

$$(2.9) \quad \widehat{M}^{\mathbb{L}} = \mathbb{L}^2 \widehat{M} + (1-\mathbb{L})(1+(z+1)\mathbb{L}).$$

It can be verified that the function \overline{M} satisfies this equation, and this implies $\overline{M} = \widehat{M}$. As in the proof given above, the argument ultimately relies on Lemma A.2. \square

3. GROTHENDIECK CLASS AND BETTI NUMBERS, AND PROOF OF THEOREM 1.1

Corollaries 1.4 and 1.5 follow directly from the first expression shown in Theorem 1.3.

Proof of Corollary 1.4. According to Theorem 1.3,

$$1+z+\sum_{n \geq 3} [\overline{\mathcal{M}}_{0,n}] \frac{z^{n-1}}{(n-1)!} = \sum_{\ell \geq 0} \left(\frac{(\ell+1)^\ell}{(\ell+1)!} ((1-\mathbb{L})(1+(z+1)\mathbb{L}))^{\frac{1+\ell(1-\mathbb{L})}{\mathbb{L}}} \prod_{j=0}^{\ell-1} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) \right) \mathbb{L}^\ell.$$

Thus, the class $[\overline{\mathcal{M}}_{0,n}]$ may be obtained by setting $z = 0$ in the $(n-1)$ -st derivative of this expression with respect to z . A simple induction proves that

$$\frac{\partial^{n-1}}{\partial z^{n-1}} \left((1+(z+1)\mathbb{L})^{\frac{1+\ell}{\mathbb{L}}-\ell} \right) = (1+(z+1)\mathbb{L})^{\frac{1+\ell}{\mathbb{L}}-\ell-n+1} \prod_{j=0}^{n-2} (1+\ell-(\ell+j)\mathbb{L}),$$

and hence

$$\frac{\partial^{n-1}}{\partial z^{n-1}} \left((1 + (z+1)\mathbb{L})^{\frac{1+\ell}{\mathbb{L}} - \ell} \right) \Big|_{z=0} = (1 + \mathbb{L})^{\frac{1+\ell}{\mathbb{L}} - \ell - n + 1} (\ell+1)^{n-1} \prod_{j=0}^{n-2} \left(1 - \frac{(\ell+j)\mathbb{L}}{\ell+1} \right).$$

It follows that $[\overline{\mathcal{M}}_{0,n}]$ equals

$$(3.1) \quad \sum_{\ell \geq 0} \frac{(\ell+1)^{\ell+n-1}}{(\ell+1)!} (1 - \mathbb{L})^{\frac{1+\ell}{\mathbb{L}} - \ell} (1 + \mathbb{L})^{\frac{1+\ell}{\mathbb{L}} - \ell - n + 1} \left(\prod_{j=0}^{\ell+n-2} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^\ell$$

and the expression stated in Corollary 1.4 follows. \square

Proof of Corollary 1.5. The rank of $H^{2\ell}(\overline{\mathcal{M}}_{0,n})$ is the coefficient of \mathbb{L}^ℓ in (3.1). The logarithms of the individual factors in each summand in (3.1) expand as follows.

• $\log \left((1 - \mathbb{L})^{\frac{1+k}{\mathbb{L}} - k} \right)$:

$$\begin{aligned} \left(\frac{1+k}{\mathbb{L}} - k \right) \log(1 - \mathbb{L}) &= - \sum_{i \geq 0} (1+k) \frac{\mathbb{L}^i}{i+1} + \sum_{i \geq 0} k \frac{\mathbb{L}^{i+1}}{i+1} \\ &= -(1+k) + \sum_{i \geq 1} \frac{k-i}{i(i+1)} \mathbb{L}^i. \end{aligned}$$

• $\log \left((1 + \mathbb{L})^{\frac{1+k}{\mathbb{L}} - k - n + 1} \right)$:

$$\begin{aligned} \left(\frac{1+k}{\mathbb{L}} - k - n + 1 \right) \log(1 + \mathbb{L}) &= (1+k) \sum_{i \geq 0} (-1)^i \frac{\mathbb{L}^i}{i+1} - (k+n-1) \sum_{i \geq 0} (-1)^i \frac{\mathbb{L}^{i+1}}{i+1} \\ &= (1+k) + \sum_{i \geq 1} (-1)^i \frac{2ki + ni + k + n - 1}{i(i+1)} \mathbb{L}^i. \end{aligned}$$

• $\log \left(\prod_{j=0}^{k+n-2} \left(1 - \frac{j\mathbb{L}}{k+1} \right) \right)$:

$$- \sum_{j=0}^{k+n-2} \sum_{i \geq 0} \frac{j^{i+1} \mathbb{L}^{i+1}}{(i+1)(k+1)^{i+1}} = - \sum_{i \geq 1} \left(\sum_{j=0}^{k+n-2} j^i \right) \frac{\mathbb{L}^i}{i(k+1)^i}.$$

Therefore, the product of the three factors is the exponential of $\sum_{i \geq 1} C_{nki} \mathbb{L}^i$ where for $n \geq 0$, $k \geq 0$, $i \geq 1$ we set

$$(3.2) \quad C_{nki} := \frac{(-1)^i (2ki + ni + k + n - 1) + k - i}{i(i+1)} - \frac{1}{i(k+1)^i} \sum_{j=0}^{k+n-2} j^i.$$

With this notation,

$$\begin{aligned} [\overline{\mathcal{M}}_{0,n}] &= \sum_{\ell \geq 0} \frac{(\ell+1)^{\ell+n-1}}{(\ell+1)!} \exp \left(\sum_{i \geq 1} C_{nli} \mathbb{L}^i \right) \mathbb{L}^\ell \\ &= \sum_{\ell \geq 0} \frac{(\ell+1)^{\ell+n-1}}{(\ell+1)!} \sum_{m \geq 0} \frac{1}{m!} \left(\sum_{i \geq 1} C_{nli} \mathbb{L}^i \right)^m \mathbb{L}^\ell. \end{aligned}$$

We have

$$\left(\sum_{i \geq 1} C_{nki} \mathbb{L}^i \right)^m = \sum_{r \geq 0} \left(\sum_{i_1 + \dots + i_m = r} C_{nki_1} \cdots C_{nki_m} \right) \mathbb{L}^r$$

where the indices i_j in the summation are positive integers. The stated formula (1.4)

$$\text{rk } H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \sum_{k=0}^{\ell} \frac{(k+1)^{k+n-1}}{(k+1)!} \sum_{m=0}^{\ell-k} \frac{1}{m!} \sum_{i_1 + \dots + i_m = \ell-k} C_{nki_1} \cdots C_{nki_m}$$

follows. \square

Remark 3.1. In parsing this formula, it is important to keep in mind that the indices i_j in the summation are positive. For example, when $k = \ell$, i.e., $\ell - k = 0$, there is exactly one contribution to the last \sum , that is, the empty choice of indices ($m = 0$); that summand is the empty product of the coefficients C_{nki} , that is, 1. For instance, $\text{rk } H^0(\overline{\mathcal{M}}_{0,n}) = 1$ for all n .

By contrast, if $m > \ell - k$, then the last \sum is the empty sum, so the contribution of such terms is 0. This is why we can bound m by $\ell - k$ in the range of summation. \square

The classical *Faulhaber's formula* states that for $i \geq 1$

$$\sum_{j=0}^N j^i = \frac{1}{i+1} \sum_{j=0}^i \binom{i+1}{j} B_j N^{i-j+1}$$

where B_j denote the Bernoulli numbers (with the convention $B_1 = \frac{1}{2}$). Applying Faulhaber's formula turns (3.2) into

$$\begin{aligned} C_{nki} = \frac{1}{i(i+1)} & \left((-1)^i (2ki + ni + k + n - 1) + k - i \right. \\ & \left. - \frac{1}{(k+1)^i} \sum_{j=0}^i \binom{i+1}{j} B_j (k+n-1)^{i-j+1} \right). \end{aligned}$$

as stated in the introduction.

Proof of Theorem 1.1. To prove Theorem 1.1, we use the second expression for \widehat{M} obtained in Theorem 1.3:

$$\widehat{M} = \sum_{\ell \geq 0} \left(\sum_{k \geq 0} \frac{(\ell+1)^{\ell+k}}{(\ell+1)!k!} (z - \mathbb{L} - z\mathbb{L})^k \prod_{j=0}^{\ell+k-1} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^\ell.$$

The class is obtained by setting $z = 0$ in the $(n-1)$ -st derivative with respect to z . This straightforward operation yields

$$(1 - \mathbb{L})^{n-1} \sum_{\ell \geq 0} \left(\sum_{k \geq n-1} \frac{(\ell+1)^{\ell+k}}{(\ell+1)!(k-n+1)!} (-\mathbb{L})^{k-n+1} \prod_{j=0}^{\ell+k-1} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^\ell$$

and therefore, after simple manipulations,

$$[\overline{\mathcal{M}}_{0,n}] = (1 - \mathbb{L})^{n-1} \sum_{k \geq 0} \left(\sum_{\ell=0}^k \frac{(-1)^{k-\ell} (\ell+1)^{k+n-1}}{(\ell+1)!(k-\ell)!} \prod_{j=0}^{k+n-2} \left(1 - \frac{j\mathbb{L}}{\ell+1} \right) \right) \mathbb{L}^k.$$

Now recall that the Stirling numbers of the first kind, $s(N, j)$, are defined by the identity

$$\sum_{j=0}^N s(N, j)x^j = \prod_{j=0}^{N-1} (x - j) = x(x - 1) \cdots (x - N + 1).$$

Setting $y = \frac{1}{x}$ and clearing denominators shows that

$$\prod_{j=0}^{N-1} (1 - jy) = \sum_{j=0}^N s(N, N - j)y^j.$$

We then have

$$\prod_{j=0}^{k+n-2} \left(1 - \frac{j\mathbb{L}}{\ell+1}\right) = \sum_{j=0}^{k+n-1} s(k+n-1, k+n-1-j) \left(\frac{\mathbb{L}}{\ell+1}\right)^j$$

yielding

$$[\overline{\mathcal{M}}_{0,n}] = (1 - \mathbb{L})^{n-1} \sum_{k \geq 0} \sum_{j=0}^{k+n-1} \sum_{\ell=0}^k \frac{(-1)^{k-\ell} (\ell+1)^{k+n-1-j}}{(\ell+1)!(k-\ell)!} s(k+n-1, k+n-1-j) \mathbb{L}^{k+j}.$$

Next, recall that the Stirling numbers of the second kind, $S(N, r)$, are defined by

$$(3.3) \quad S(N, r) = \sum_{i=1}^r (-1)^{r-i} \frac{i^N}{i!(r-i)!}$$

for all $N > 0$, $r \geq 0$. Therefore

$$\sum_{\ell=0}^k (-1)^{k-\ell} \frac{(\ell+1)^{k+n-1-j}}{(\ell+1)!(k-\ell)!} = \sum_{i=1}^{k+1} (-1)^{k+1-i} \frac{i^{k+n-1-j}}{i!(k+1-i)!} = S(k+n-1-j, k+1)$$

and we can conclude

$$[\overline{\mathcal{M}}_{0,n}] = (1 - \mathbb{L})^{n-1} \sum_{k \geq 0} \sum_{j=0}^{k+n-1} S(k+n-1-j, k+1) s(k+n-1, k+n-1-j) \mathbb{L}^{k+j},$$

which is the statement. \square

Remark 3.2. Since $S(N, r) = 0$ if $r > N$, nonzero summands in this expression only occur for $0 \leq j \leq n-2$. \square

Corollary 1.2 follows immediately from Theorem 1.1, as the reader may verify.

4. THE GENERATING FUNCTION P , AND PROOF OF THEOREM 1.6

We now move to the polynomials $p_m^{(k)}(z) \in \mathbb{Q}[z]$ mentioned in §1. These polynomials were introduced in [ACM25] for the purpose of studying the generating functions for individual Betti numbers of $\overline{\mathcal{M}}_{0,n}$. As proved in [ACM25], for all $m \geq 0$ and $0 \leq m \leq k$ there exist polynomials $p_m^{(k)}(z) \in \mathbb{Q}[z]$ of degree $2m$ such that

$$(4.1) \quad \sum_{n \geq 3} \text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n}) \frac{z^{n-1}}{(n-1)!} = e^z \sum_{m=0}^k (-1)^m p_m^{(k)}(z) e^{(k-m)z}$$

for all $k > 0$. Let

$$P(z, t, u) := \sum_{\ell \geq 0} \sum_{m \geq 0} p_m^{(m+\ell)}(z) u^m t^\ell$$

be the generating function for the polynomials $p_m^{(k)}$; then (4.1) states that

$$(4.2) \quad \widehat{M}(z, \mathbb{L}) = e^z P(z, e^z \mathbb{L}, -\mathbb{L}).$$

In this section we prove Theorem 1.6 from the introduction, which gives an explicit expression for the generating function P :

$$P(z, t, u) = \sum_{\ell \geq 0} \left(\frac{(\ell+1)^\ell}{(\ell+1)!} e^{-(\ell+1)z} ((1+u)(1-u(z+1)))^{-\frac{1+\ell(u+1)}{u}} \prod_{j=0}^{\ell-1} \left(1 + \frac{ju}{\ell+1} \right) \right) t^\ell.$$

This statement will follow from Theorem 1.3, but the argument is not completely trivial for the mundane reason that $\widehat{M}(z, \mathbb{L})$ is a two-variable function while $P(z, t, u)$ is a three-variable function; $\widehat{M}(z, \mathbb{L})$ is a specialization of $P(z, t, u)$, not conversely.

Proof of Theorem 1.6. We let

$$\overline{P}(z, t, u) = \sum_{\ell \geq 0} \left(\frac{(\ell+1)^\ell}{(\ell+1)!} e^{-(\ell+1)z} ((1+u)(1-u(z+1)))^{-\frac{1+\ell(u+1)}{u}} \prod_{j=0}^{\ell-1} \left(1 + \frac{ju}{\ell+1} \right) \right) t^\ell$$

and we have to prove that $\overline{P}(z, t, u) = P(z, t, u)$. Equivalently, we will prove that

$$(4.3) \quad \overline{P}(z, su, -u) = P(z, su, -u);$$

note that the right-hand side equals $\sum_{k \geq 0} \sum_{m=0}^k (-1)^m p_m^{(k)} s^{k-m} u^k$.

Claim 4.1. *There exist polynomials $\overline{p}^{(k)}(z, s) \in \mathbb{Q}[z, s]$ such that*

$$\overline{P}(z, su, -u) = \sum_{k \geq 0} \overline{p}^{(k)}(z, s) u^k.$$

This claim implies (4.3). Indeed, by Theorem 1.3 and identity (4.1) we have

$$\sum_{k \geq 0} \overline{p}^{(k)}(z, e^z) \mathbb{L}^k = \overline{P}(z, e^z \mathbb{L}, -\mathbb{L}) = e^{-z} \widehat{M}(z, \mathbb{L}) = \sum_{k \geq 0} \left(\sum_{m=0}^k (-1)^m p_m^{(k)}(z) e^{(k-m)z} \right) \mathbb{L}^k$$

and therefore

$$(4.4) \quad \overline{p}^{(k)}(z, e^z) = \sum_{m=0}^k (-1)^m p_m^{(k)}(z) e^{(k-m)z}$$

for all $k \geq 0$. Given that Claim 4.1 holds,

$$\overline{p}^{(k)}(z, s) - \sum_{m=0}^k (-1)^m p_m^{(k)}(z) s^{(k-m)}$$

is then a *polynomial* vanishing at $s = e^z$, hence it must be identically 0 since e^z is transcendental over $\mathbb{Q}(z)$. The needed identity (4.3) follows.

Thus, we are reduced to verifying Claim 4.1.

By definition,

$$\overline{P}(z, su, -u) = \sum_{\ell \geq 0} \left(\frac{(\ell+1)^\ell}{(\ell+1)!} e^{-(\ell+1)z} ((1-u)(1+u(z+1)))^{\frac{1+\ell(1-u)}{u}} \prod_{j=0}^{\ell-1} \left(1 - \frac{ju}{\ell+1} \right) \right) s^\ell u^\ell.$$

In order to prove Claim 4.1, we have to verify that for all $k \geq 0$ the coefficient of u^k in

$$\sum_{\ell=0}^k \left(\frac{(\ell+1)^\ell}{(\ell+1)!} e^{-(\ell+1)z} ((1-u)(1+u(z+1)))^{\frac{1+\ell(1-u)}{u}} \prod_{j=0}^{\ell-1} \left(1 - \frac{ju}{\ell+1} \right) \right) s^\ell u^\ell$$

is a polynomial in z and s . This is clearly a polynomial in s , and it suffices then to verify that the coefficient of u^i in the factor

$$e^{-(\ell+1)z} (1+u(z+1))^{\frac{1+\ell(1-u)}{u}}$$

is a polynomial in z for all $i \geq 0$. Simple manipulations show that this factor equals

$$\begin{aligned} & \exp \left(\frac{1+\ell(1-u)}{u} \log(1+u(z+1)) - (1+\ell)z \right) \\ &= \sum_{k \geq 0} \frac{1}{k!} \left((1+\ell) + \sum_{i \geq 0} \left(\frac{(1+\ell)(z+1)}{i+2} + \frac{\ell}{i+1} \right) (-u(z+1))^{i+1} \right)^k \end{aligned}$$

and the statement is clear from this expression. (In fact, the coefficient of u^i in this expression is a polynomial of degree $2i$ in z .) This concludes the proof of Claim 4.1, and therefore of Theorem 1.6. \square

Of course the same result may be formulated in different ways. The following expression follows from Theorem 1.6 by manipulations analogous to the corresponding manipulations in the proof of Theorem 1.3.

Corollary 4.2.

$$P(z, t, u) = \sum_{\ell \geq 0} \sum_{k \geq 0} \left(\frac{(\ell+1)^{\ell+k}}{(\ell+1)!k!} (z+u+zu)^k \prod_{j=0}^{\ell+k-1} \left(1 + \frac{ju}{\ell+1} \right) \right) e^{-(\ell+1)z} t^\ell.$$

Remark 4.3. It is straightforward to verify that the function $P(z, t, u)$ is a solution of the differential equation

$$\frac{\partial P}{\partial z} + t \frac{\partial P}{\partial t} + P = \frac{P}{1 - u(1+z) - tP}.$$

This is the analogue for P of the differential equation (2.1) satisfied by \widehat{M} .

It is also not difficult to obtain a functional equation for P ,

$$(e^z P)^{-u} + utP = (1+u)(1-u(z+1)),$$

lifting the functional equation (2.9) satisfied by \widehat{M} . \square

5. THE POLYNOMIALS $p_m^{(k)}$ AND Γ_{mj} , AND PROOF OF THEOREMS 1.7 AND 1.9

We are interested in studying more thoroughly the polynomials $p_m^{(k)}(z) \in \mathbb{Q}[z]$ defined in §4. As proved in [ACM25], $p_m^{(k)}$ has degree $2m$ and positive leading coefficient, and these polynomials determine the Betti numbers of $\overline{\mathcal{M}}_{0,n}$ in the sense that

$$\sum_{n \geq 3} \text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n}) \frac{z^{n-1}}{(n-1)!} = e^z \sum_{m=0}^k (-1)^m p_m^{(k)}(z) e^{(k-m)z}$$

for $k > 0$ (cf. (4.1)). For instance, it follows that for every $k \geq 0$ the sequence of Betti numbers $\text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n})$ as $n = 3, 4, 5, \dots$ is determined by a finite amount of information,

specifically, the $(k+1)^2$ coefficients of the polynomials $p_m^{(k)}$, $m = 0, \dots, k$. It is hoped that more information about the polynomials $p_m^{(k)}$ will help in proving conjectured facts about the integers $\text{rk } H^{2k}(\overline{\mathcal{M}}_{0,n})$, such as log-concavity properties.

The following list of the first several polynomials $p_m^{(k)}$ is reproduced from [ACM25].

$$p_0^{(0)} = 1$$

$$p_0^{(1)} = 1$$

$$p_1^{(1)} = \frac{1}{2}z^2 + z + 1$$

$$p_0^{(2)} = \frac{3}{2}$$

$$p_1^{(2)} = z^2 + 3z + 2$$

$$p_2^{(2)} = \frac{1}{8}z^4 + \frac{5}{6}z^3 + 2z^2 + 2z + \frac{1}{2}$$

$$p_0^{(3)} = \frac{8}{3}$$

$$p_1^{(3)} = \frac{9}{4}z^2 + \frac{15}{2}z + 5$$

$$p_2^{(3)} = \frac{1}{2}z^4 + \frac{11}{3}z^3 + 9z^2 + 9z + 3$$

$$p_3^{(3)} = \frac{1}{48}z^6 + \frac{7}{24}z^5 + \frac{35}{24}z^4 + \frac{7}{2}z^3 + \frac{17}{4}z^2 + \frac{5}{2}z + \frac{2}{3}$$

$$p_0^{(4)} = \frac{125}{24}$$

$$p_1^{(4)} = \frac{16}{3}z^2 + \frac{56}{3}z + \frac{38}{3}$$

$$p_2^{(4)} = \frac{27}{16}z^4 + \frac{51}{4}z^3 + \frac{129}{4}z^2 + \frac{65}{2}z + \frac{45}{4}$$

$$p_3^{(4)} = \frac{1}{6}z^6 + \frac{13}{6}z^5 + \frac{21}{2}z^4 + \frac{74}{3}z^3 + 30z^2 + 18z + \frac{13}{3}$$

$$p_4^{(4)} = \frac{1}{384}z^8 + \frac{1}{16}z^7 + \frac{5}{9}z^6 + \frac{49}{20}z^5 + \frac{289}{48}z^4 + \frac{103}{12}z^3 + \frac{85}{12}z^2 + \frac{19}{6}z + \frac{13}{24}$$

In this section we prove that all polynomials $p_m^{(k)}$ have positive coefficients and provide evidence for the assertion that most $p_m^{(k)}$ are ultra-log-concave. Both facts were conjectured in [ACM25].

We denote by $c_{mj}^{(k)} \in \mathbb{Q}$ the coefficient of z^j in $p_m^{(k)}(z)$. The following proposition introduces rational numbers $\Gamma_{mj}(\ell)$ and establishes the equality (1.7) stated in Theorem 1.7.

Proposition 5.1. *For $\ell \geq 0$ and $j \geq 0$ let $\Gamma_{mj}(\ell) \in \mathbb{Q}$ be defined by the identity*

$$\sum_{m \geq 0} \sum_{j=0}^{2m} \Gamma_{mj}(\ell) z^j u^m = e^{-z} ((1 + (\ell + 1)u)(1 - (z + \ell + 1)u))^{-\frac{1}{u} - \ell} \prod_{j=0}^{\ell-1} (1 + ju).$$

Then $\Gamma_{mj}(\ell) = 0$ for $j > 2m$ and

$$c_{mj}^{(k)} = \frac{(k-m+1)^{k-2m+j}}{(k-m+1)!} \cdot \Gamma_{mj}(k-m)$$

for all k, m, j with $0 \leq m \leq k$, $0 \leq j \leq 2m$.

Proof. This follows from an application of Theorem 1.6:

$$\begin{aligned} & \sum_{\ell \geq 0} \sum_{m \geq 0} \sum_{j \geq 0} \frac{(\ell+1)^{\ell-m+j}}{(\ell+1)!} \Gamma_{mj}(\ell) z^j u^m t^\ell \\ &= \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} \sum_{m \geq 0} \sum_{j \geq 0} \Gamma_{mj}(\ell) ((\ell+1)z)^j \left(\frac{u}{\ell+1}\right)^m t^\ell \\ &= \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} e^{(\ell+1)z} ((1+u)(1-(z+1)u))^{-\frac{\ell+1}{u}-\ell} \prod_{j=0}^{\ell-1} \left(1 + \frac{ju}{\ell+1}\right) \\ &= P(z, t, u) = \sum_{\ell \geq 0} \sum_{m \geq 0} p^{(m+\ell)}(z) u^m t^\ell \\ &= \sum_{\ell \geq 0} \sum_{m \geq 0} \sum_{j=0}^{2m} c_{mj}^{(m+\ell)} z^j u^m t^\ell. \end{aligned}$$

Comparing coefficients gives the statement. \square

In order to prove Theorem 1.7 we have to verify that $\Gamma_{mj}(\ell)$ is a polynomial in ℓ and attains positive values for $\ell \geq 0$ in the range $m \geq 0$, $0 \leq j \leq 2m$. For this purpose, we introduce another set of ancillary rational numbers $\Delta_{mj}(\ell)$, defined by the generating function

$$(5.1) \quad \sum_{m \geq 0} \sum_{j \geq 0} \Delta_{mj}(\ell) z^j u^m = e^{-z} ((1 + (\ell+1)u)(1 - u(z + \ell + 1)))^{-(\frac{1}{u} + \ell)};$$

thus,

$$(5.2) \quad \sum_{m \geq 0} \sum_{j=0}^{2m} \Gamma_{mj}(\ell) z^j u^m = \left(\sum_{m \geq 0} \sum_{j=0}^{2m} \Delta_{mj}(\ell) z^j u^m \right) \prod_{j=0}^{\ell-1} (1 + ju).$$

Proposition 5.2. *For all $m \geq 0$ and $j = 0, \dots, 2m$, $\Delta_{mj}(\ell)$ is a polynomial in ℓ of degree $2m-j$ and with positive coefficients. For $j > 2m$, $\Delta_{mj}(\ell) = 0$.*

Proof. The proof amounts to elementary but delicate calculations. Note that

$$\begin{aligned} & \log \left(e^{-z} ((1 + (\ell+1)u)(1 - u(z + \ell + 1)))^{-(\frac{1}{u} + \ell)} \right) \\ &= -z + \left(-\frac{1}{u} - \ell \right) (\log(1 + (\ell+1)u) + \log(1 - (\ell+1+z)u)) \\ &= -z + \left(-\frac{1}{u} - \ell \right) \sum_{i \geq 0} \frac{(-1)^i (\ell+1)^{i+1} - (\ell+1+z)^{i+1}}{i+1} u^{i+1}. \end{aligned}$$

The reader will verify that this expression equals

$$\begin{aligned} \sum_{i \geq 1} \frac{1}{i(i+1)} & \left((\ell+1)^i (i+2\ell i + \ell + (-1)^{i+1}(i-\ell)) \right. \\ & \left. + \sum_{j=1}^i \binom{i+1}{j} (\ell+1)^{i-j} (\ell(i-j) + i + \ell + \ell i) z^j + i z^{i+1} \right) u^i, \end{aligned}$$

that is, the coefficient of $z^j u^i$ in the expression is 0 for $i = 0$ and for $j > i+1$, and equals

$$\begin{cases} \frac{1}{i(i+1)} (\ell+1)^i (i+2\ell i + \ell + (-1)^{i+1}(i-\ell)) & \text{for } j = 0 \\ \frac{1}{i(i+1)} \binom{i+1}{j} (\ell+1)^{i-j} (\ell(i-j) + i + \ell + \ell i) & \text{for } 1 \leq j \leq i \\ \frac{1}{i+1} & \text{for } j = i+1 \end{cases}$$

for $i \geq 1$. For all $j = 0, \dots, i+1$, this is a polynomial in ℓ of degree $i+1-j$. Further, trivially $(i-j) \geq 0$ in the range $1 \leq j \leq i$ and

$$i+2\ell i + \ell + (-1)^{i+1}(i-\ell) = \begin{cases} 2i(\ell+1) > 0 & \text{if } i \text{ is odd} \\ 2\ell(i+1) > 0 & \text{if } i \text{ is even.} \end{cases}$$

Summarizing,

$$\sum_{m \geq 0} \sum_{j \geq 0} \Delta_{mj}(\ell) z^j u^m = \exp \left(\sum_{i \geq 1} \sum_{j=0}^{i+1} \delta_{ij}(\ell) z^j u^i \right)$$

with $\delta_{ij}(\ell)$ polynomials of degree $i+1-j$ with positive coefficients. Now

$$\exp(c_1 u + c_2 u^2 + c_3 u^3 + \dots) = 1 + c_1 u + \left(\frac{c_1^2}{2} + c_2 \right) u^2 + \left(c_3 + c_1 c_2 + \frac{c_1^3}{6} \right) u^3 + \dots$$

is a series whose coefficient of u^m is a linear combination of products $c_{i_1} \cdots c_{i_r}$ with $i_k > 0$ and $\sum i_k = m$. Therefore: for every m , $\sum_{j \geq 0} \Delta_{mj}(\ell) z^j$ is a linear combination of terms

$$\left(\sum_{j_1=0}^{i_1+1} \delta_{i_1 j_1}(\ell) z^{j_1} \right) \cdots \left(\sum_{j_r=0}^{i_r+1} \delta_{i_r j_r}(\ell) z^{j_r} \right) = \sum \delta_{i_1 j_1}(\ell) \cdots \delta_{i_r j_r}(\ell) z^{j_1 + \cdots + j_r}.$$

with $i_k > 0$ and $\sum i_k = m$. Each polynomial $\delta_{i_1 j_1}(\ell) \cdots \delta_{i_r j_r}(\ell)$ has positive coefficients and degree

$$\sum (i_k + 1) - \sum j_k = m + r - j$$

with $j = \sum j_k$ the exponent of z . The degree of the sum is the maximum attained by $m+r-j$, that is, $2m-j$ (for $r=m$, $i_1 = \cdots = i_m = 1$). Thus, the coefficient $\Delta_{mj}(\ell)$ has degree $2m-j$ in ℓ , as stated. The exponent j itself ranges from 0, attained for $j_r = 0$ for all r , to $\sum (i_k + 1) = m + r$, and attains the maximum $2m$, again when all i_k equal 1.

Thus $\Delta_{mj}(\ell) = 0$ for $j > 2m$, and this concludes the proof of the proposition. \square

The polynomials $\Delta_{mj}(\ell)$ can of course be computed explicitly:

$$\Delta_{00} = 1, \quad \Delta_{10} = (\ell+1)^2, \quad \Delta_{11} = 2\ell+1, \quad \Delta_{12} = \frac{1}{2}, \quad \Delta_{20} = \frac{(\ell^2+4\ell+1)(\ell+1)^2}{2},$$

etc. We have verified that all the polynomials $\Delta_{mj}(\ell)$ with $1 \leq m \leq 50$, $0 \leq j \leq 2m$, are ultra-log-concave. The first several hundred are in fact real-rooted, but $\Delta_{19,1}(\ell)$ appears not to be.

Proposition 5.2 implies another part of Theorem 1.7, thereby confirming the first statement in Conjecture 1.

Corollary 5.3. *For all $m \geq 0$, $0 \leq j \leq 2m$, and $\ell \geq 0$ we have $\Gamma_{mj}(\ell) > 0$.*

Therefore, for all $m \geq 0$, $0 \leq j \leq 2m$, and $k \geq m$, $p_m^{(k)}(z)$ is a degree $2m$ polynomial with positive coefficients.

Proof. The second part of the statement follows from the first by Proposition 5.1.

The first part is a consequence of (5.2), since the polynomials $\Delta_{mj}(\ell)$ have positive coefficients by Proposition 5.2 and so does the factor $\prod_{j=0}^{\ell-1} (1 + ju)$. \square

The following proposition will complete the proof of Theorem 1.7.

Proposition 5.4. *For all $m \geq 0$ and $0 \leq j \leq m$, the functions $\Gamma_{mj}(\ell)$ are polynomials of degree $2m - j$ in ℓ .*

Proof. Faulhaber's formula for sums of powers $\sum_{j=0}^{\ell-1} j^i$ easily implies that

$$\begin{aligned} \prod_{j=0}^{\ell-1} (1 + ju) &= \exp \left(\sum_{i \geq 1} \sum_{j=0}^i \binom{i+1}{j} B_j \ell^{i-j+1} \frac{(-1)^{i+1} u^i}{i(i+1)} \right) \\ &= 1 + \frac{1}{2} \ell(\ell-1)u + \frac{1}{24} \ell(\ell-1)(\ell-2)(3\ell-1)u^2 + \frac{1}{48} \ell^2(\ell-1)^2(\ell-2)(\ell-3)u^3 + \dots \\ &=: \sum_{m \geq 0} \beta_m(\ell) u^m \end{aligned}$$

where B_j denotes the j -th Bernoulli number. It follows that the coefficient $\beta_m(\ell)$ of u^m in the expansion of $\prod_{j=0}^{\ell-1} (1 + ju)$ is a polynomial in $\mathbb{Q}[\ell]$ of degree $2m$ (and of course $\beta_m(\ell) = 0$ for $m \geq \ell$). By (5.2) we have

$$\Gamma_{mj}(\ell) = \sum_{m_1+m_2=m} \Delta_{m_1j}(\ell) \beta_{m_2}(\ell)$$

and it follows that $\Gamma_{mj}(\ell)$ is a polynomial in $\mathbb{Q}[\ell]$ of degree $2m - j$ as stated. \square

This concludes the proof of Theorem 1.7.

The information collected so far may also be used to investigate the conjectured log-concavity properties of the polynomials $p_m^{(k)}$ (cf. Conjecture 1 in §1). With the notation introduced above, ultra-log concavity for $p_m^{(k)}$ is the following condition:

$$\forall j = 1, \dots, 2m-1: \quad \left(\frac{c_{mj}^{(k)}}{\binom{2m}{j}} \right)^2 \geq \frac{c_{m,j-1}^{(k)}}{\binom{2m}{j-1}} \cdot \frac{c_{m,j+1}^{(k)}}{\binom{2m}{j+1}}.$$

By Corollary 5.3 $c_{mj}^{(k)} > 0$ in this range, therefore these sequences automatically have no internal zeros.

Lemma 5.5. *The polynomial $p_m^{(m+\ell)}$ is ultra-log-concave if and only if*

$$\forall j = 1, \dots, 2m-1: \quad j(2m-j)\Gamma_{mj}(\ell)^2 - (j+1)(2m-j+1)\Gamma_{m,j-1}(\ell)\Gamma_{m,j+1}(\ell) \geq 0.$$

Proof. Immediate consequence of Proposition 5.1. \square

Since by Proposition 5.4 each $\Gamma_{mj}(\ell)$ is a *polynomial* in ℓ , Lemma 5.5 provides us with an effective way to test the ultra-log-concavity of all polynomials $p_m^{(k)}$, $k \geq m$, for any given m . For example, according to Lemma 5.5, in order to verify that $p_2^{(k)}$ is ultra-log-concave for all $k \geq 2$, it suffices to observe that the polynomials

$$\begin{aligned} & \frac{9}{4}\ell^6 + \frac{131}{12}\ell^5 + \frac{41}{3}\ell^4 + \frac{85}{4}\ell^3 + \frac{403}{12}\ell^2 + \frac{67}{3}\ell + 4 \\ & \frac{13}{4}\ell^4 + \frac{25}{2}\ell^3 + 9\ell^2 + \frac{5}{4}\ell + 1 \\ & \frac{1}{4}\ell^2 + \frac{3}{4}\ell + \frac{1}{12} \end{aligned}$$

are trivially positive-valued for all $\ell \geq 0$. For larger m not all the polynomials appearing in Lemma 5.5 have positive coefficients, but it is a straightforward computational process to verify whether they only attain positive values for $\ell \geq 0$.

Proposition 5.6. *The polynomials $p_m^{(k)}(z)$ are ultra-log concave for $m \leq 100$ and all $k \geq m$, with the exceptions listed in Conjecture 1.*

Proof. As in the foregoing discussion, Lemma 5.5 reduces this statement to a finite computation, which we carried out with maple. \square

As one more application of the material developed above, we have the following formula for the Betti numbers of $\overline{\mathcal{M}}_{0,n}$ in terms of the polynomials Γ_{mj} :

$$(5.3) \quad \text{rk } H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \sum_{k+m=\ell} (-1)^m \frac{(k+1)^{n-2+k-m}}{k!} \sum_{j=0}^{2m} (n-1) \cdots (n-j) \Gamma_{mj}(k).$$

(This is Theorem 1.9.)

Proof. According to [ACM25, Theorem 5.1],

$$\text{rk } H^{2\ell}(\overline{\mathcal{M}}_{0,n}) = \frac{(\ell+1)^{\ell+n-1}}{(\ell+1)!} + \sum_{m=1}^{\ell} (-1)^m \sum_{j=0}^{2m} \binom{n-1}{j} c_{mj}^{(\ell)} j! (\ell-m+1)^{n-1-j}.$$

Identity (5.3) follows then from Proposition 5.1. \square

6. THE LAMBERT W FUNCTION AND PROOF OF THEOREM 1.11

The goal of this section is the proof of Theorem 1.11, expressing the generating function $\widehat{M} = 1 + z + \sum_{n \geq 3} [\overline{\mathcal{M}}_{0,n}] \frac{z^{n-1}}{(n-1)!}$ in terms of the principal branch W of the classical *Lambert W function*, cf. [CGH⁺96]. In fact it is more notationally convenient to work with the ‘tree function’, defined by

$$T(t) = -W(-t) = \sum_{n \geq 1} \frac{n^{n-1} t^n}{n!}$$

and we recall that this function satisfies the relation

$$(6.1) \quad T(t) = t e^{T(t)}$$

([CGH⁺96, (1.5)]).

Consider the coefficient of u^m in the generating function P introduced in §4:

$$P_m(z, t) = \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) t^\ell$$

(cf. (1.8)). It is clear that this function may be expressed as a series in the tree function $T = T(t)$, since $t = e^{-T}T$ according to (6.1):

$$(6.2) \quad P_m(z, t) = \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) e^{-\ell T(t)} T(t)^\ell.$$

We are going to verify that this series is a *rational function* in T up to a factor e^T , and in fact admits the particularly simple expression (1.9):

$$P_m = e^T \frac{F_m(z, T)}{(1 - T)^{2m-1}}$$

where $F_0 = \frac{1}{1-T}$ and F_m are polynomials for $m > 0$, of degree $< 3m$. This is Proposition 1.10, which we proceed to prove. Ultimately, this fact is a consequence of Proposition 5.4.

Proof of Proposition 1.10. For $m = 0$, using (1.6):

$$P_0(z, t) = \sum_{\ell \geq 0} p_0^{(\ell)} t^\ell = \sum_{\ell \geq 0} \frac{(\ell+1)^\ell}{(\ell+1)!} t^\ell = \frac{1}{t} T(t) = e^T$$

by (6.1), and this is the statement. For $m > 0$, and treating T as an indeterminate, consider the expression

$$(6.3) \quad \begin{aligned} (1 - T)^{2m-1} \sum_{\ell \geq 0} \frac{(\ell+1)^{\ell+a}}{(\ell+1)!} e^{-(\ell+1)T} T^\ell \\ = (1 - T)^{2m-1} \sum_{\ell \geq 0} \sum_{k \geq 0} \frac{(\ell+1)^{\ell+k+a}}{(\ell+1)!k!} (-1)^k T^{k+\ell}. \end{aligned}$$

For $1 \leq a$, (6.3) can be expressed in terms of Stirling numbers of the second kind (cf. (3.3)) and we can apply Lemma A.3:

$$\begin{aligned} (1 - T)^{2m-1} \sum_{N \geq 0} \sum_{\ell \geq 0} \frac{(\ell+1)^{N+a}}{(\ell+1)!(N-\ell)!} (-1)^{N-\ell} T^N \\ = (1 - T)^{2m-1} \sum_{N \geq 0} \sum_{i \geq 1} \frac{i^{N+a}}{i!(N+1-i)!} (-1)^{N+1-i} T^N \\ = (1 - T)^{2m-1} \sum_{N \geq 0} S(N+a, N+1) T^N \\ = (1 - T)^{2(m-a)} \sigma_a(T) \end{aligned}$$

with $\sigma_a(T)$ a polynomial of degree $\leq a-1$. In particular, for $a \leq m$, (6.3) is a polynomial of degree $\leq 2m-a-1$ in this case. If $a \leq 0$ and $N \geq -a$,

$$\sum_{i \geq 1} \frac{i^{N+a}}{i!(N+1-i)!} (-1)^{N+1-i} = S(N+a, N+1) = 0$$

since $N+a \leq N+1$. Thus, (6.3) is a polynomial of degree $\leq 2m-a-1$ in this case as well.

It follows that for all $m \geq 0$, $0 \leq j \leq 2m$, and $0 \leq r \leq 2m-j$,

$$(1 - T)^{2m-1} \sum_{\ell \geq 0} \frac{(\ell+1)^{\ell-m+j}}{(\ell+1)!} (\ell+1)^r e^{-(\ell+1)T} T^\ell$$

is a polynomial of degree $\leq 3m - j - 1$. By Proposition 5.4, $\Gamma_{mj}(\ell)$ is a polynomial of degree $2m - j$. Therefore it is a linear combination of polynomials $(\ell + 1)^r$, $0 \leq r \leq 2m - j$, and we can conclude that

$$(1 - T)^{2m-1} \sum_{\ell \geq 0} \frac{(\ell + 1)^{\ell-m+j}}{(\ell + 1)!} \Gamma_{mj}(\ell) e^{-(\ell+1)T} T^\ell$$

is a polynomial of degree $\leq 3m - 1$. By Proposition 5.1, this implies that

$$F_m(z, T) := (1 - T)^{2m-1} \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) e^{-(\ell+1)T} T^\ell$$

is a polynomial of degree $\leq 3m - 1$ in T . Now set $T = T(t)$ and use (6.2):

$$P_m(z, t) = \sum_{\ell \geq 0} p_m^{(m+\ell)}(z) e^{-\ell T(t)} T(t)^\ell = e^{T(t)} \frac{F_m(z, T(t))}{(1 - T(t))^{2m-1}}$$

to verify (1.9) as needed. \square

Theorem 1.11 follows from Proposition 1.10. Indeed, recall (4.2):

$$\widehat{M}(z, \mathbb{L}) = e^z P(z, e^z \mathbb{L}, -\mathbb{L});$$

since $P(z, t, u) = \sum_{m \geq 0} P_m(z, t) u^m$, Proposition 1.10 gives

$$\widehat{M}(z, \mathbb{L}) = e^z e^T \sum_{m \geq 0} \frac{F_m(z, T)}{(1 - T)^{2m-1}} (-\mathbb{L})^m$$

where now $T = T(e^z \mathbb{L})$; by (6.1),

$$e^z e^{T(e^z \mathbb{L})} = e^z e^{-z} \mathbb{L}^{-1} T(e^z \mathbb{L}) = \frac{T}{\mathbb{L}}$$

and (1.10) follows. \square

7. STIRLING MATRICES AND THE GROTHENDIECK CLASS OF $\overline{\mathcal{M}}_{0,n}$

The material in this section is included mostly for aesthetic reasons. We recast Theorem 1.1 in terms of products involving the infinite matrices $\mathfrak{s} = (s(i, j))_{i, j \geq 1}$, resp., $\mathfrak{S} = (S(i, j))_{i, j \geq 1}$, defined by Stirling numbers of the first, resp., second kind:

$$\mathfrak{s} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \cdots \\ -1 & 1 & 0 & 0 & 0 & \cdots \\ 2 & -3 & 1 & 0 & 0 & \cdots \\ -6 & 11 & -6 & 1 & 0 & \cdots \\ 24 & -50 & 35 & -10 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad \mathfrak{S} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & 0 & \cdots \\ 1 & 3 & 1 & 0 & 0 & \cdots \\ 1 & 7 & 6 & 1 & 0 & \cdots \\ 1 & 15 & 25 & 10 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We will also use the notation

$$\mathbb{I}_\gamma := \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & \gamma & 0 & 0 & \cdots \\ 0 & 0 & \gamma^2 & 0 & \cdots \\ 0 & 0 & 0 & \gamma^3 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad \text{Sh}_k := \begin{pmatrix} \overbrace{0 \cdots 0}^k & 1 & 0 & 0 & \cdots \\ 0 & \cdots & 0 & 0 & 1 & 0 & \cdots \\ 0 & \cdots & 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The matrix Sh_k is obtained from the identity matrix by shifting entries to the right so that the nonzero entries are placed along the NW-SE diagonal starting at $(1, k+1)$. For an infinite matrix A , the ‘ k -th trace’

$$\text{tr}_k(A) := \text{tr}(\text{Sh}_k \cdot A)$$

is the sum of the entries in the k -th subdiagonal, provided of course that this sum is defined, for example as a formal power series.

It is well-known that \mathfrak{s} and \mathfrak{S} are inverses of each other. We consider the following matrix, obtained as the product of the commutator of $\mathbb{1}_{\mathbb{L}}$ and \mathfrak{s} by $\mathbb{1}_{\mathbb{L}}$:

$$\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}} = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 - \mathbb{L} & \mathbb{L} & 0 & 0 & \cdots \\ 1 - 3\mathbb{L} + 2\mathbb{L}^2 & 3\mathbb{L} - 3\mathbb{L}^2 & \mathbb{L}^2 & 0 & \cdots \\ 1 - 6\mathbb{L} + 11\mathbb{L}^2 - 6\mathbb{L}^3 & 7\mathbb{L} - 18\mathbb{L}^2 + 11\mathbb{L}^3 & 6\mathbb{L}^2 - 6\mathbb{L}^3 & \mathbb{L}^3 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

With this notation, the following is a restatement of Theorem 1.1.

Theorem 7.1.

$$[\overline{\mathcal{M}}_{0,n}] = (1 - \mathbb{L})^{n-1} \cdot \text{tr}_{n-2}(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}).$$

The trace appearing in this statement is well-defined as a series: since the i^{th} column of the matrix is a multiple of \mathbb{L}^{i-1} , only finitely many entries on every diagonal contribute to the coefficient of each power of \mathbb{L} .

Example 7.2. The entries in the fourth subdiagonal are

$$\begin{aligned} 1 - 10\mathbb{L} + 35\mathbb{L}^2 - 50\mathbb{L}^3 + 24\mathbb{L}^4 \\ 31\mathbb{L} - 225\mathbb{L}^2 + 595\mathbb{L}^3 - 675\mathbb{L}^4 + 274\mathbb{L}^5 \\ 301\mathbb{L}^2 - 1890\mathbb{L}^3 + 4375\mathbb{L}^4 - 4410\mathbb{L}^5 + 1624\mathbb{L}^6 \\ 1701\mathbb{L}^3 - 9800\mathbb{L}^4 + 20930\mathbb{L}^5 - 19600\mathbb{L}^6 + 6769\mathbb{L}^7 \\ 6951\mathbb{L}^4 - 37800\mathbb{L}^5 + 76440\mathbb{L}^6 - 68040\mathbb{L}^7 + 22449\mathbb{L}^8 \end{aligned}$$

etc., adding up to

$$\text{tr}_4(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) = 1 + 21\mathbb{L} + 111\mathbb{L}^2 + 356\mathbb{L}^3 + 875\mathbb{L}^4 + \cdots$$

and

$$(1 - \mathbb{L})^5 \cdot \text{tr}_4(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) = 1 + 16\mathbb{L} + 16\mathbb{L}^2 + \mathbb{L}^3 = [\overline{\mathcal{M}}_{0,6}]$$

as it should. \square

Proof of Theorem 7.1. The (a, b) -entry of the matrix $\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}$ is

$$(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}})_{(a,b)} = \sum_c \mathbb{L}^{a-1} s(a, c) \mathbb{L}^{-(c-1)} S(c, b) \mathbb{L}^{b-1} = \sum_c s(a, c) S(c, b) \mathbb{L}^{a+b-c-1}.$$

For $a = k+n-1$, $b = k+1$, this gives

$$\sum_{j \geq 0} s(k+n-1, k+n-1-j) S(k+n-1-j, k+1) \mathbb{L}^{k+j}.$$

By Theorem 1.1,

$$\begin{aligned} [\overline{\mathcal{M}}_{0,n}] &= (1 - \mathbb{L})^{n-1} \cdot \sum_{k \geq 0} (\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}})_{(k+n-1, k+1)} \\ &= (1 - \mathbb{L})^{n-1} \cdot \text{tr}_{n-2}(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) \end{aligned}$$

as stated. \square

The product by $(1 - \mathbb{L})^{n-1}$ can also be accounted for in terms of this matrix calculus, giving

$$[\overline{\mathcal{M}}_{0,n}] = \text{tr}_{n-2} \left((1 - \mathbb{L}) \cdot \mathbb{1}_{1-\mathbb{L}} \cdot \mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}} \cdot \mathbb{1}_{(1-\mathbb{L})^{-1}} \right).$$

8. ABOUT THE EULER CHARACTERISTIC OF $\overline{\mathcal{M}}_{0,n}$

We will end with a few comments on the Euler characteristic $\chi(\overline{\mathcal{M}}_{0,n})$ of $\overline{\mathcal{M}}_{0,n}$. The generating function

$$\widehat{\chi}(z) := 1 + z + \sum_{n \geq 3} \chi(\overline{\mathcal{M}}_{0,n}) \frac{z^{n-1}}{(n-1)!}$$


is the specialization of \widehat{M} at $\mathbb{L} = 1$, but cannot be recovered from the expressions obtained in Theorem 1.3 because of convergence issues. However, specializing the differential equation (2.1) for \widehat{M} at $\mathbb{L} = 1$ gives

$$\frac{d\widehat{\chi}}{dz} = \frac{\widehat{\chi}}{2 + z - \widehat{\chi}}$$

with initial condition $\widehat{\chi}(0) = 1$ (cf. [Man95, (0.10)], with different notation), whose solution is

$$(8.1) \quad \widehat{\chi}(z) = \frac{z+2}{-W_{-1}(-(z+2)e^{-2})} = 1 + z + \frac{z^2}{2!} + 2 \frac{z^3}{3!} + 7 \frac{z^4}{4!} + 34 \frac{z^5}{5!} + 213 \frac{z^6}{6!} + \dots$$

Here, W_{-1} is the other real branch of the Lambert W-function, cf. [CGH⁺96].



We do not know how to extract an explicit expression for $\chi(\overline{\mathcal{M}}_{0,n})$ from (8.1); but any of the formulas obtained for the Betti numbers in this note yields such an expression. For example, Corollary 1.2 implies

$$\chi(\overline{\mathcal{M}}_{0,n}) = \sum_{\ell=0}^{n-3} \sum_{j=0}^{\ell} \sum_{k=0}^{\ell-j} (-1)^{\ell-j-k} \binom{n-1}{\ell-j-k} s(k+n-1, k+n-1-j) S(k+n-1-j, k+1).$$

An alternative formulation may be obtained in terms of the shifted trace

$$\mathrm{tr}_{n-2}(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}}^{-1} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) = \sum_{j \geq 0} s(k+n-1, k+n-1-j) S(k+n-1-j, k+1) \mathbb{L}^{k+j}$$

appearing in Theorem 7.1.

Proposition 8.1.

$$\mathrm{tr}_{n-2}(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}}^{-1} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) = \sum_{k \geq 0} a_n(k) \mathbb{L}^k$$

where $a_n(k)$ is a polynomial with rational coefficients, degree $n-2$, and leading term $\frac{\chi(\overline{\mathcal{M}}_{0,n})}{(n-2)!}$.

Proof. The class $[\overline{\mathcal{M}}_{0,n}]$ is a polynomial of degree $n-3$ in \mathbb{L} , therefore there exist integers b_0, \dots, b_{n-3} such that

$$[\overline{\mathcal{M}}_{0,n}] = \sum_{\ell=0}^{n-3} b_{\ell} (1-\mathbb{L})^{\ell},$$

and with this notation $b_0 = [\overline{\mathcal{M}}_{0,n}]|_{\mathbb{L}=1} = \chi(\overline{\mathcal{M}}_{0,n})$. By Theorem 7.1,

$$\begin{aligned} \mathrm{tr}_{n-2}(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}}^{-1} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) &= \frac{[\overline{\mathcal{M}}_{0,n}]}{(1-\mathbb{L})^{n-1}} = \sum_{\ell=0}^{n-3} \frac{b_{\ell}}{(1-\mathbb{L})^{n-\ell-1}} \\ &= \sum_{k \geq 0} \left(\sum_{\ell=0}^{n-3} b_{\ell} \binom{k+n-\ell-2}{n-\ell-2} \right) \mathbb{L}^k. \end{aligned}$$

It follows that the coefficient $a_n(k)$ of \mathbb{L}^k is a polynomial as stated, and

$$a_n(k) = b_0 \frac{k^{n-2}}{(n-2)!} + \text{lower order terms},$$

concluding the proof. \square

Example 8.2. In Example 7.2 we noted

$$\mathrm{tr}_4(\mathbb{1}_{\mathbb{L}} \cdot \mathfrak{s} \cdot \mathbb{1}_{\mathbb{L}^{-1}} \cdot \mathfrak{S} \cdot \mathbb{1}_{\mathbb{L}}) = 1 + 21\mathbb{L} + 111\mathbb{L}^2 + 356\mathbb{L}^3 + 875\mathbb{L}^4 + \dots$$

Tracing the argument in the proof of Proposition 8.1 shows that these coefficients are values of the polynomial

$$a_6(k) = \frac{17}{12} k^4 + \frac{17}{3} k^3 + \frac{97}{12} k^2 + \frac{29}{6} k + 1,$$

and $\chi(\overline{\mathcal{M}}_{0,6}) = (6-2)! \cdot \frac{17}{12} = 34$. \square

APPENDIX A. COMBINATORIAL IDENTITIES

Here are collected the combinatorial statements used in this paper; proofs are included for completeness.

Lemma A.1. *For all integers $k, w \geq 1$,*

$$\sum_{\ell+m=k-1} \binom{w(\ell+1)-1}{\ell} \cdot \binom{w(m+1)-2}{m} \cdot \frac{1}{m+1} = \binom{w(k+1)-2}{k-1}.$$

Proof. We will use one form of the Lagrange series identity, [Rio68, §4.5]:

$$\frac{f(x)}{1 - y\varphi'(x)} = \sum_{r \geq 0} \frac{y^r}{r!} \left[\frac{d^r}{dx^r} (f(x)\varphi^r(x)) \right]_{x=0},$$

where $y = \frac{x}{\varphi(x)}$. Applying this identity with $f(x) = (x+1)^\alpha$, $\varphi(x) = (x+1)^\beta$ gives

$$(A.1) \quad \frac{(x+1)^{\alpha+1}}{x+1-\beta x} = \sum_{r \geq 0} \binom{\alpha + \beta r}{r} y^r.$$

Setting $\alpha = w-1$, $\beta = w$ in (A.1) gives

$$A(x) := \sum_{\ell \geq 0} \binom{w(\ell+1)-1}{\ell} y^\ell = \frac{(x+1)^w}{x+1-wx}.$$

Setting

$$B(x) := \sum_{m \geq 0} \binom{w(m+1)-2}{m} \cdot \frac{y^m}{m+1}$$

and applying (A.1) with $\alpha = w-2$, $\beta = w$ gives

$$\frac{d}{dy}(By) = \sum_{m \geq 0} \binom{w(m+1)-2}{m} y^m = \frac{(x+1)^{w-1}}{x+1-wx},$$

from which

$$By = \int \frac{(x+1)^{w-1}}{x+1-wx} dy = \int \frac{(x+1)^{w-1}}{x+1-wx} \cdot \frac{dy}{dx} dx = \int \frac{1}{(x+1)^2} dx = -\frac{1}{x+1} + C.$$

Setting $x = 0$ determines $C = 1$, and we get

$$B(x) = (x+1)^{w-1}.$$

Another application of (A.1), with $\alpha = 2w-2$, $\beta = w$, gives

$$\begin{aligned} & \sum_{\ell \geq 0} \binom{w(\ell+1)-1}{\ell} y^\ell \cdot \sum_{m \geq 0} \binom{w(m+1)-2}{m} \cdot \frac{y^m}{m+1} \\ &= A(x)B(x) = \frac{(x+1)^{2w-1}}{1-x(w-1)} \\ &= \sum_{r \geq 0} \binom{w(r+2)-2}{r} y^r. \end{aligned}$$

Extracting the coefficient of y^{k-1} verifies the stated identity. \square

Lemma A.2. *For all positive integers α, β , we have*

$$x^\alpha = \sum_{\ell \geq 0} \frac{\alpha}{\alpha + \ell\beta} \binom{\alpha + \ell\beta}{\ell} y^\ell,$$

where $y = (x-1)x^{-\beta}$.

Proof. We will use a second form of the Lagrange series identity, [Rio68, §4.5]:

$$f(x) = f(0) + \sum_{\ell=1}^{\infty} \frac{y^\ell}{\ell!} \left(\frac{d^{\ell-1}}{dx^{\ell-1}} (f'(x)\varphi^\ell(x)) \right) |_{x=0}$$

where $y = \frac{x}{\varphi(x)}$. Apply this identity with $f(x) = (x+1)^\alpha$, $\varphi(x) = (x+1)^\beta$, $y = x(x+1)^{-\beta}$. We have

$$\begin{aligned} \frac{1}{\ell!} \frac{d^{\ell-1}}{dx^{\ell-1}} (\alpha(x+1)^{\alpha+\ell\beta-1})|_{x=0} &= \frac{\alpha}{\ell!} (\alpha + \ell\beta - 1) \cdots (\alpha_\ell\beta - (\ell-1)) (x+1)^{\alpha+\ell\beta-\ell}|_{x=0} \\ &= \frac{\alpha}{\alpha + \ell\beta} \binom{\alpha + \ell\beta}{\ell} \end{aligned}$$

and hence

$$(x+1)^\alpha = \sum_{\ell \geq 0} \frac{\alpha}{\alpha + \ell\beta} \binom{\alpha + \ell\beta}{\ell} (x(x+1)^{-\beta})^\ell.$$

Substituting $x-1$ for x gives the statement. \square

Lemma A.3. *For $N, r > 0$, let $S(N, r)$ denote the Stirling number of the second kind (cf. (3.3)). Then for all integers $a \geq 2$ there exists a polynomial $\sigma_a(x)$ of degree $a-2$ such that*

$$\sum_{N \geq 0} S(N+a, N+1) x^N = \frac{\sigma_a(x)}{(1-x)^{2a-1}}.$$

For $a = 1$, the same holds with $\sigma_a(x) = 1$.

Remark A.4. The coefficients of the polynomials σ_a are the numbers denoted $B_{k,i}$ in [GS78], where it is proved ([GS78, Theorem 2.1]) that they agree with the numbers of ‘‘Stirling permutations of the multiset $\{1, 1, 2, 2, \dots, k, k\}$ with exactly i descents.’’ In particular, they are positive integers. \square

Proof. Since $S(N+1, N+1) = 1$ and $S(N+2, N+1) = \binom{N+2}{2}$ for all $N \geq 0$, the statement is immediate for $a = 1$ and $a = 2$. For $a > 2$, let

$$\text{St}_a(x) := \sum_{N \geq 0} S(N+a, N+1) x^N = \frac{\sigma_a(x)}{(1-x)^{2a-1}}$$

and recall the basic recursive identity satisfied by Stirling numbers of the second kind:

$$S(N+a, N+1) = S(N+a-1, N) + (N+1)S(N+a-1, N+1).$$

This implies the relation

$$\text{St}_a(x) = \frac{1}{1-x} \cdot \frac{d}{dx} (x \text{St}_{a-1}(x))$$

from which the statement follows easily. In fact, this identity implies the recursion

$$\sigma_a(x) = (1-x)(\sigma_{a-1}(x) + x\sigma'_{a-1}(x)) + (2a-3)x\sigma_{a-1}(x),$$

verifying that $\sigma_a(x)$ is a polynomial of degree $a-2$, with leading coefficient $(a-1)!$, if $\sigma_{a-1}(x)$ is a polynomial of degree $a-3$ with leading coefficient $(a-2)!$. \square

REFERENCES

- [ACM25] Paolo Aluffi, Stephanie Chen, and Matilde Marcolli. Log concavity of the Grothendieck class of $\overline{\mathcal{M}}_{0,n}$. *J. Algebraic Combin.*, 61(4):47, 2025.
- [CGH⁺96] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth. On the Lambert W function. *Adv. Comput. Math.*, 5(4):329–359, 1996.
- [CKL] Jinwon Choi, Young-Hoon Kiem, and Donggun Lee. Recursive algorithm and log-concavity of representations on the cohomology of $\overline{\mathcal{M}}_{0,n}$. arXiv:2408.10728.
- [CKL23] Jinwon Choi, Young-Hoon Kiem, and Donggun Lee. Representations on the cohomology of $\overline{\mathcal{M}}_{0,n}$. *Adv. Math.*, 435(part A):Paper No. 109364, 66, 2023.

- [CKL25] Jinwon Choi, Young-Hoon Kiem, and Donggun Lee. Characteristic polynomial of $\overline{\mathcal{M}}_{0,n}$ and log-concavity. *Proc. Lond. Math. Soc. (3)*, 131(5):Paper No. e70096, 2025.
- [EFM⁺] Christopher Eur, Luis Ferroni, Jacob P. Matherne, Roberto Pagaria, and Lorenzo Vecchi. Building sets, Chow rings, and their Hilbert series. arXiv:2504.16776.
- [Get95] Ezra Getzler. Operads and moduli spaces of genus 0 Riemann surfaces. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 199–230. Birkhäuser Boston, Boston, MA, 1995.
- [Gou56] H. W. Gould. Some generalizations of Vandermonde’s convolution. *Amer. Math. Monthly*, 63:84–91, 1956.
- [GS78] Ira Gessel and Richard P. Stanley. Stirling polynomials. *J. Combinatorial Theory Ser. A*, 24(1):24–33, 1978.
- [Kee92] Sean Keel. Intersection theory of moduli space of stable n -pointed curves of genus zero. *Trans. Amer. Math. Soc.*, 330(2):545–574, 1992.
- [Man95] Yuri I. Manin. Generating functions in algebraic geometry and sums over trees. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 401–417. Birkhäuser Boston, Boston, MA, 1995.
- [MM16] Yuri I. Manin and Matilde Marcolli. Moduli operad over \mathbb{F}_1 . In *Absolute arithmetic and \mathbb{F}_1 -geometry*, pages 331–361. Eur. Math. Soc., Zürich, 2016.
- [Nas] Eduardo Nascimento. Improved bounds on ultra-log concavity of the Grothendieck class of $\overline{\mathcal{M}}_{0,n}$. arXiv:2509.11805.
- [Rio68] John Riordan. *Combinatorial identities*. John Wiley & Sons, Inc., New York-London-Sydney, 1968.

MATHEMATICS DEPARTMENT, FLORIDA STATE UNIVERSITY, TALLAHASSEE FL 32306, U.S.A.

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA CA 91105, U.S.A.

E-mail address: aluffi@math.fsu.edu

E-mail address: matilde@caltech.edu

E-mail address: ehnascim@caltech.edu