

Set 9: Second Order Linear ODEs - Part 1

Kyle A. Gallivan

Department of Mathematics

Florida State University

Ordinary Differential Equations

Fall 2009

Second Order Linear ODEs

Definition 9.1. An ODE is second order and linear if it can be written

$$P(t)y'' + Q(t)y' + R(t)y = G(t)$$

An associated initial value problem also specifies

$$y(t_0) = y_0$$

$$y'(t_0) = y'_0$$

If $P(t) \neq 0$ the explicit form

$$y'' + p(t)y' + q(t)y = g(t)$$

will often be used.

Second Order Linear ODEs

Definition 9.2. Given two functions p and q , the differential operator L associated with the ODE that maps a twice differentiable function of t on an interval $\alpha < t < \beta$ to another function of t is given by

$$L[\phi] = \phi'' + p\phi' + q\phi$$

Existence and Uniqueness

Theorem 9.1 (Textbook page 146). *If p, q , and g are continuous functions on the interval $\mathcal{I} : \alpha < t < \beta$ that contains t_0 then the initial value problem*

$$y'' + p(t)y' + q(t)y = g(t)$$

$$y(t_0) = y_0$$

$$y'(t_0) = y'_0$$

has a unique solution that exist throughout \mathcal{I} .

Superposition

Theorem 9.2 (Textbook page 147). *If y_1 and y_2 are two solutions to the homogeneous ODE*

$$L[y] = y'' + p(t)y' + q(t)y = 0$$

then the linear combination $c_1y_1 + c_2y_2$ is a solution for any constants c_1 and c_2 .

Overview

- homogeneous second order linear ODEs with constant coefficients
- general solution and fundamental solution sets
- Wronskian
- general solution for nonhomogeneous problems
- Method of undetermined coefficients for nonhomogeneous constant coefficient problems
- Method of variation of parameters for nonhomogeneous constant coefficient problems
- some nonconstant coefficient problems

Constant Coefficient Problems

Definition 9.3. If a , b , c are real constants, the ODE

$$ay'' + by' + cy = 0$$

is a second order linear ODE with constant coefficients.

- How are they solved?
- What are the important properties of the solutions?

Constant Coefficient Problems

Recall the first order problem

$$y' + cy = 0 \rightarrow y = e^{-ct}$$

Hypothesis for second order problem $y = e^{rt}$.

$$y = e^{rt}$$

$$y' = re^{rt}$$

$$y'' = r^2 e^{rt}$$

$$\begin{aligned} ay'' + by' + cy &= a(r^2 e^{rt}) + b(re^{rt}) + c(e^{rt}) \\ &= [ar^2 + br + c]e^{rt} \end{aligned}$$

Constant Coefficient Problems

$$ay'' + by' + cy = [ar^2 + br + c]e^{rt}$$

$$e^{rt} \neq 0 \rightarrow ar^2 + br + c = 0$$

r generates a solution to the ODE if it is a root of $ar^2 + br + c$

Consistent with first order

$$y' + cy = 0$$

$$a = 0, \quad b = 1$$

$$br + c = 0 \leftrightarrow r = -c$$

Solutions

$$\text{ODE: } ay'' + by' + cy = 0$$

$$\text{characteristic equation: } ar^2 + br + c$$

$$b^2 - 4ac > 0 \rightarrow \text{two distinct real roots}$$

$$b^2 - 4ac < 0 \rightarrow \text{two complex conjugate roots}$$

$$b^2 - 4ac = 0 \rightarrow \text{repeated real root}$$

Distinct Real Roots

$$y'' - y = 0, \quad y(0) = 2, \quad y'(0) = -1$$

$$r^2 - 1 = 0 \rightarrow r = \pm 1$$

$$y_1 = c_1 e^t, \quad y_2 = c_2 e^{-t}$$

$$y_1(0) = 2 = c_1 e^0 = c_1$$

$$y_1'(0) = -1 \neq c_1 e^0 = 2 \times 1 = 2$$

Cannot satisfy the initial conditions with $y_1 y_1$ solves ODE but not the IVP.

Distinct Real Roots

$$y = c_1 e^t + c_2 e^{-t} \quad y' = c_1 e^t - c_2 e^{-t}$$

$$y(0) = c_1 + c_2 = 2$$

$$y'(0) = c_1 - c_2 = -1$$

$$c_1 = \frac{1}{2}, \quad c_2 = \frac{3}{2}$$

$$y = \frac{1}{2} e^t + \frac{3}{2} e^{-t}$$

To satisfy an arbitrary initial condition, we need two solutions.

Distinct Real Roots

- ODE: $ay'' + by' + cy$
- characteristic equation: $ar^2 + br + c$
- $b^2 - 4ac > 0$, two distinct real roots r_1 and r_2
- general solution: $y = c_1e^{r_1t} + c_2e^{r_2t}$
- any $y(t_0)$ and $y'(t_0)$ values can be satisfied by choosing c_1 and c_2

Distinct Real Roots

Three cases of asymptotic behavior:

- if both roots are negative then $y \rightarrow 0$
- if at least one root is positive then $y \rightarrow \pm\infty$
- if one root is 0 and the other is negative then $y \rightarrow C$

The initial condition on y' determines the initial shape and the asymptotic behavior determines the final shape.

Example

$$y'' + 5y' + 6y = 0, \quad y(0) = 2, \quad y'(0) = 3$$

$$r^2 + 5r + 6 = (r + 2)(r + 3)$$

$$y = c_1 e^{-2t} + c_2 e^{-3t}$$

$$c_1 + c_2 = 2$$

$$-2c_1 - 3c_2 = 3$$

$$c_1 = 9, \quad c_2 = -7$$

$$y = 9e^{-2t} - 7e^{-3t}$$

Complex Conjugate Roots

$$ay'' + by' + cy = 0$$

$$ar^2 + br + c = 0$$

$b^2 - 4ac < 0 \rightarrow$ two complex conjugate roots

$$r_1 = \lambda + i\mu, \quad r_2 = \lambda - i\mu$$

$$y_1 = e^{(\lambda+i\mu)t}, \quad y_2 = e^{(\lambda-i\mu)t}$$

Euler's Formula

Definition 9.4 (Euler's Formula). For any real μ

$$e^{i\mu t} = \cos \mu t + i \sin \mu t$$

We therefore have,

$$\begin{aligned} e^{(\lambda+i\mu)t} &= e^{\lambda t} e^{i\mu t} \\ &= e^{\lambda t} \cos \mu t + e^{\lambda t} i \sin \mu t \end{aligned}$$

Complex Conjugate Roots

We want a real solution \rightarrow use linear combination.

$$y_1 = e^{\lambda t} \cos \mu t + e^{\lambda t} i \sin \mu t$$

$$y_2 = e^{\lambda t} \cos \mu t - e^{\lambda t} i \sin \mu t$$

$$y_1 + y_2 = (2)e^{\lambda t} \cos \mu t$$

$$y_1 - y_2 = (2i)e^{\lambda t} \sin \mu t$$

$$y = c_1 e^{\lambda t} \cos \mu t + c_2 e^{\lambda t} \sin \mu t$$

Real initial conditions will yield real c_i

Complex Conjugate Roots

- ODE: $ay'' + by' + cy$
- characteristic equation: $ar^2 + br + c$
- $b^2 - 4ac < 0$, complex conjugate roots $\lambda \pm i\mu$
- general solution: $y = c_1e^{\lambda t} \cos \mu t + c_2e^{\lambda t} \sin \mu t$
- any $y(t_0)$ and $y'(t_0)$ values can be satisfied by choosing c_1 and c_2
- real solutions are sinusoidal oscillations with frequency μ and damping/expansion factor $e^{\lambda t}$
- $\lambda = 0$ yields bounded oscillation

Example

$$16y'' - 8y' + 145y = 0, \quad y(0) = -2, \quad y'(0) = 1$$

$$16r^2 - 8r + 145 = 0$$

$$r = \frac{1}{4} \pm 3i$$

$$y = c_1 e^{0.25t} \cos 3t + c_2 e^{0.25t} \sin 3t$$

$$c_1 \times 1 + c_2 \times 0 = -2$$

$$0.25c_1 + 3c_2 = 1$$

$$c_1 = -2 \quad c_2 = \frac{1}{2}$$

$$y = -2e^{0.25t} \cos 3t + \frac{1}{2}e^{0.25t} \sin 3t$$

Growing oscillation.

Example

$$y'' + 9y = 0$$

$$r^2 + 9 = 0$$

$$r = \pm 3i$$

$$y = c_1 e^{0t} \cos 3t + c_2 e^{0t} \sin 3t = c_1 \cos 3t + c_2 \sin 3t$$

Bounded oscillation for any real y_0, y'_0 .

Repeated Real Root

$$ay'' + by' + cy = 0$$

$$ar^2 + br + c = 0$$

$$b^2 - 4ac = 0 \rightarrow \text{repeated real root}$$

$$r_1 = r_2 = r$$

$$y_1 = e^{rt}$$

$$y_2 = te^{rt}$$

Example

$$y'' + 4y' + 4y = 0$$

$$(r + 2)^2 = 0$$

$$y_1 = e^{-2t}, \quad y_2 = te^{-2t}$$

Check y_2 in ODE

$$y_2 = te^{-2t}$$

$$y_2' = -2e^{-2t}t + e^{-2t} = e^{-2t}(1 - 2t)$$

$$y_2'' = -2e^{-2t}(1 - 2t) - 2e^{-2t} = -2e^{-2t}(2 - 2t) = -4e^{-2t}(1 - t)$$

$$\begin{aligned} y'' + 4y' + 4y &= -4e^{-2t}(1 - t) + 4e^{-2t}(1 - 2t) + 4te^{-2t} \\ &= 4e^{-2t}[-1 + t + 1 - 2t + t] = 0 \end{aligned}$$

Example

$$y'' + 4y' + 4y = 0, \quad y(0) = 1, \quad y'(0) = -1$$

$$(r + 2)^2 = 0$$

$$y_1 = e^{-2t}, \quad y_2 = te^{-2t}$$

$$y = c_1 e^{-2t} + c_2 t e^{-2t} \quad \text{and} \quad y' = -2c_1 e^{-2t} - c_2 e^{-2t}(1 - 2t)$$

$$c_1 = 1$$

$$-2c_1 - c_2 = -1$$

$$y = e^{-2t} - t e^{-2t}$$

$$y' = -2e^{-2t} + e^{-2t}(1 - 2t)$$

Summary

- $ay'' + by' + cy = 0$ constant coefficients
- characteristic equation: $ar^2 + br + c$
- distinct real roots r_1 and r_2

$$y = c_1e^{r_1t} + c_2e^{r_2t}$$

- complex conjugate roots $\lambda \pm i\mu$

$$y = c_1e^{\lambda t} \cos \mu t + c_2e^{\lambda t} \sin \mu t$$

- repeated real root r

$$y = c_1e^{rt} + c_2te^{rt}$$

- Proofs of general solution forms?