

UNIT 2 MODULE 9
MEASURES OF CENTRAL TENDENCY

EXAMPLE 2.9.1

To paraphrase Benjamin Disraeli: "There are lies, darn lies, and DAM STATISTICS."

Compute the mean, median and mode for the following DAM STATISTICS:

<i>Name of Dam</i>	<i>Height</i>
Oroville dam	756 ft.
Hoover dam	726 ft.
Glen Canyon dam	710 ft.
Don Pedro dam	568 ft.
Hungry Horse dam	564 ft.
Round Butte dam	440 ft.
Pine Flat Lake dam	440 ft.

MEASURES of CENTRAL TENDENCY

A measure of central tendency is a number that represents the typical value in a collection of numbers. Three familiar measures of central tendency are the **mean**, the **median**, and the **mode**.

We will let **n** represent the number of data points in the distribution. Then

$$\mathbf{Mean} = \frac{\text{sum of all data points}}{n}$$

(The mean is also known as the "average" or the "arithmetic average.")

Median = "middle" data point (or average of two middle data points) when the data points are arranged in numerical order.

Mode = the value that occurs most often (if there is such a value).

HACKING MATHEMATICS

In **EXAMPLE 2.9.1** the distribution has 7 data points, so $n = 7$.

$$\begin{aligned}\text{MEAN} &= (756 + 726 + 710 + 568 + 564 + 440 + 440)/7 \\ &= 4204/7 = 600.57 \text{ (this has been rounded).}\end{aligned}$$

We can say that the typical dam is 600.57 feet tall.

We can also use the **MEDIAN** to describe the typical response. In order to find the median we must first list the data points *in numerical order*:

756, 726, 710, 568, 564, 440, 440

Now we choose the number in the middle of the list.

756, 726, 710, **568**, 564, 440, 440

The median is 568.

Because the median is 568 it is also reasonable to say that on this list the typical dam is 568 feet tall.

We can also use the **MODE** to describe the typical dam height. Since the number 440 occurs more often than any of the other numbers on this list, the mode is 440.

EXAMPLE 2.9.2

Survey question: How many semester hours are you taking this semester?

Responses: 15, 12, 18, 12, 15, 15, 12, 18, 15, 16

What was the typical response?

FINDING THE "MIDDLE" OF A LIST OF NUMBERS

In the two previous examples, we found the median by first arranging the list numerically and then crossing off data points from each end of the list until we arrived at the middle. This method of “crossing off” works well as long as there are relatively few data points to work with. In cases where we are dealing with a large collection of data, however, it is not a practical method for finding the median.

If n represents the number of data points in a distribution, then:

the position of the "middle value" is $\frac{n+1}{2}$.

If the data points have been arranged numerically, we can use this fact to efficiently find the median.

EXAMPLE

For the following list, $n = 19$. Find the median.

24, 25, 28, 31, 33, 33, 36, 42, 42, 48, 51, 57, 57, 68, 75, 79, 79, 79, 85

SOLUTION

The numbers are already in numerical order. The position of the "middle of the list" is:
 $(n+1)/2 = (19+1)/2 = 20/2 = 10$

Thus, the tenth number will be the median. We count until we arrive at the tenth number.

24, 25, 28, 31, 33, 33, 36, 42, 42, **48**, 51, 57, 57, 68, 75, 79, 79, 79, 85

The median is 48.

EXAMPLE 2.9.3

Compute the mean, median, and mode for this distribution of test scores:

92, 68, 80, 68, 84

UNIT 2 MODULE 9

We can summarize the information above in the following **frequency table**:

Value	Frequency
0	27
1	96
2	58
3	54
4	18
5	7

Now this table conveys everything that was significant about the distribution of data that we presented at the beginning of this example. When working with frequency tables, recall this fundamental fact:

A frequency table is a shorthand representation of list of data.

The numbers in the "value" column indicate which numbers appear on the original list of data. The numbers in the "frequency" column tell how many times the corresponding value appears on the original list of data.

Now we find the mean, median and mode for the data in the table.

MODE

The mode, if it exists, is easiest to find. For data presented in a frequency table, the mode is the value associated with the greatest frequency (if there is a greatest frequency).

In this case, the greatest frequency is 96 and the associated value is "1," so the mode is "1." More students received 1 parking ticket than any of the other possibilities.

MEAN

To find the mean, we must have a convenient way to determine the sum of all the data points, and also a convenient way to determine n , the number of data points in the distribution. We may be tempted to merely add the six numbers in the "value" column, and divide by six, but that would be incorrect, because it would fail to take into account that facts that the distribution includes many more than just six data points, and the various values do not all occur with the same frequency.

To find n in a case like this, we find the sum of numbers in the "frequency" column. This makes sense, when we recall that the frequencies tell how many times each of the values occurs.

HACKING MATHEMATICS

$$n = \text{sum of frequencies} = 27 + 96 + 58 + 54 + 18 + 7 = 260$$

The mean will be the sum of all 260 data points, divided by 260.

Finding the sum of all 260 data points is simpler than it may at first seem, when we recall what the table represents. For example, since the value 0 has a frequency of 27, when we took the sum of all of the zeroes in the distribution, that subtotal would be $(0)(27) = 0$. Likewise, the second row in the table shows us that the value 1 appears 96 times in the distribution, so when we took the sum of all of the ones, we would get a subtotal of $(1)(96) = 96$.

Continuing in this vein, the next row of the table tells us that the value 2 appears in the distribution 58 times, so when we took the sum of all of the twos from the list, we would have a subtotal of $(2)(58) = 116$.

This indicates that in order to find the sum of all of the data points in a frequency table, we find the sum of all subtotals formed by multiplying a value times its frequency.

$$\begin{aligned}\text{Mean} &= \frac{(0)(27) + (1)(96) + (2)(58) + (3)(54) + (4)(18) + (5)(7)}{27 + 96 + 58 + 54 + 18 + 7} \\ &= \frac{481}{260} = 18.5\end{aligned}$$

Summarizing the process described above, we have the following general rule for determining the mean for data in a frequency table:

$$\text{Mean} = S/n$$

where **n**, the number of data points in the distribution, is obtained by finding the sum of all of the frequencies, and **S**, the sum of the data points, is found by adding all of the subtotals formed by multiplying a value by its associated frequency.

MEDIAN

To find the median, we need to recall that we are trying to find the middle number (or two middle numbers) in a list of 260 numbers. Recall from earlier that the **position** of the middle number is $(n+1)/2$.

In this case, the position of the middle number is $261/2$, or 130.5.

Since 130.5 is located between 130 and 131, the median will be the average of the 130th and 131st numbers in the ordered list (by the way, since the values are arranged numerically in the table as we read from top to bottom, this data has already been ordered). We need to count through the table until we find the 130th and 131st numbers. This is done as follows, by taking into account the **cumulative frequency** as the values in the various rows are "read" from the table. To make a column for **cumulative frequency**,

UNIT 2 MODULE 9

we pretend that we are reading data from a list, in numerical order. If we were to do so, the first 27 numbers on the list would all be "0." This means that after all of the 0s are read from the list, we would have read a total of 27 numbers. We say that the value 0 has a cumulative frequency of 27:

Value	Frequency	Cumulative Frequency
0	27	27
1	96	
2	58	
3	54	
4	18	
5	7	

Next we would read all of the 1s from the list. There are 96 of them. Thus, after we have read all the 0s and 1s from the list, we would have read a total of $27 + 96 = 123$ numbers.

We summarize this by saying that the value 1 has a cumulative frequency of 123.

Value	Frequency	Cumulative Frequency
0	27	27
1	96	123
2	58	
3	54	
4	18	
5	7	

Continuing this process, after having read all of the 0s and 1s from the list, we would read all of the 2s from the list. Since the value 2 appears on the list 58 times, after we read all of the 0s, 1s and 2s from the list, we will have read a total of $27 + 96 + 58 = 181$ numbers. We say that the value 2 has a cumulative frequency of 181.

Value	Frequency	Cumulative Frequency
0	27	27
1	96	123
2	58	181
3	54	
4	18	
5	7	

HACKING MATHEMATICS

At this point we will stop. Remember, the reason we started making the column for cumulative frequency was so that we could locate the 130th and 131st numbers, in order to determine the median. Now we see that the 130th and 131st numbers are both 2s (the cumulative frequency column tells us, in fact, that the 124th through 181st numbers on the list are all 2s).

The two middle numbers are both 2s, so *the median is 2*.

*From the previous example we generalize to form this rule for determining the **median for data in a frequency table** (this rule assumes that the values appear in the table in numerical order).*

Make a column for cumulative frequency as follows:

- 1. The cumulative frequency of the first row is the same as the frequency of the first row.*
- 2. For every other row, determine cumulative frequency by adding the frequency of that row to the cumulative frequency of the previous row.*

*When cumulative frequency first equals or exceeds $(n+1)/2$, stop and use the **value** in that row for the median. (If $(n+1)/2$ is exactly .5 greater than one of the cumulative frequencies, then the median will be the average of the associated values from that row and the next row).*

EXAMPLE 2.9.6

The frequency table below represents the distribution of scores on a ten-point quiz. Compute the mean, median, and mode for this distribution.

Quiz Scores

Value	Frequency
5	6
6	8
7	14
8	22
9	28
10	36

UNIT 2 MODULE 9

EXAMPLE 2.9.7

The frequency table below represents the distribution, according to age, of students in a certain class.

Value	Frequency
18	31
19	48
20	60
21	50
22	33

1. Find the mean, median, and mode for this distribution.
2. True or false: 18 students were 31 years old.

EXAMPLE 2.9.8

The relative frequency table below shows the distribution of scores on a quiz in the course Quantum Electrodynamics For Liberal Arts. Find the mean, median and mode.

Score	Relative Frequency
4	.03
5	.10
6	.04
7	.12
8	.33
9	.20
10	.18

EXAMPLE 2.9.9

A number of people invested \$1000 each in the Gomer Family of Mutual Funds. The frequency table below shows the current values of those investments. Compute the mean, median and mode.

Value	Frequency
0	48
50	42
75	31
100	28
150	22

EXAMPLE 2.9.10

A number of people invested \$1000 each in the Gomer Family of Mutual Funds. The frequency table below shows the current values of those investments after Gomer hit the trifecta at the dog track, and hit the Cash 5 jackpot. Compute the mean, median and mode.

Value	Frequency
0	48
50	42
75	31
100	28
150	22
2,876,423	1

EXTREME VALUES and THEIR EFFECTS ON MEAN, MEDIAN and MODE

If we compare the previous two examples, we see that the two distributions are nearly identical, except that the distribution in EXAMPLE 2.9.10 contains one extra number (2,876,423) that is significantly greater than any of the other numbers in the distribution. (A number that is significantly greater or significantly less than most of the other numbers in a distribution is called an **extreme value** or **outlier**.)

Notice that including this extreme value had a huge effect on the mean of the distribution (which increased from \$61.55 to \$16,784.58) but had no effect whatsoever on either the median or the mode. Also notice that in the distribution in EXAMPLE 2.1.10, the mean is

UNIT 2 MODULE 9

not a good representation of the typical value in the distribution. This illustrates an important general fact:

*Of the three measures of central tendency (mean, median, mode), the **mean** is the measure that is most likely to be distorted by the presence of extreme values.*

EXAMPLE 2.9.11

The annual earnings for employees of a certain restaurant are given below:

12 laborers earn \$8000 each.

10 laborers earn \$9000 each.

4 supervisors earn \$11000 each

The owner/manager earns \$240,000.

Of the three measures of central tendency, which will be the least accurate representation of "typical earnings?"

WORLD WIDE WEB NOTE

For more practice on the computing measures of central tendency for data in frequency tables, visit the companion website and try THE CENTRAL TENDERIZER.

HACKING MATHEMATICS

PRACTICE EXERCISES

- Find the median of the data: 5, 7, 4, 9, 5, 4, 4, 3
A. 5.125 B. 14 C. 4.5 D. 4
- Find the mean of the following data: 12, 10, 15, 10, 16, 12, 10, 15, 15, 13
A. 13 B. 12.5 C. 15 D. 12.8
- Find the mode of the following data: 20, 14, 12, 14, 26, 16, 18, 19, 14
A. 14 B. 17 C. 26 D. 16
- Find the mean of the following data: 0, 5, 2, 4, 0, 5, 0, 3, 0, 5, 0, 3
A. 0 B. 2.25 C. 2.5 D. 3.86
- Find the median of the following data: 25, 20, 30, 30, 20, 24, 24, 30, 31
A. 20 B. 26 C. 25 D. 30
- Find the median of the following data: 1, 6, 12, 19, 5, 0, 6
A. 6 B. 7 C. 19 D. 3.5
- Find the mean of the following data: 20, 24, 24, 24, 22, 22, 24, 22, 23, 25
A. 23.5 B. 23 C. 24 D.
- Find the mode of the following data: 5, 0, 5, 4, 12, 2, 14
A. 4 B. 5 C. 6 D.. 0
- Find the mean of the following data: 0, 5, 30, 25, 16, 18, 19, 26, 0, 20, 28
A. 0 B. 18 C. 19 D. 17
- Find the median of the following data: 9, 6, 12, 5, 17, 3, 9, 5, 10, 2, 8, 7
A. 6.5 B. 7.5 C. 6 D. 7.75

UNIT 2 MODULE 9

The table at right shows the distribution of scores on Quiz #6 in MGF1106 for Sections 01-08, Spring 1999. Refer to the table for exercises 11 - 14.

score	freq
6	11
8	26
10	27
12	32
14	31
16	12
18	15
20	7

- 11.** Select the statement that is correct.
 A. 6 people had scores of 11.
 B. 27 people had scores of 10.
 C. A and B are both correct.
 D. All of these are false.

- 12.** Find the median quiz score.
 A. 13.5 B. 12 C. 31.5 D. 13

- 13.** Find the mean quiz score.
 (Answers may have been rounded.)
 A. 12 B. 12.2 C. 13 D. 13.4

- 14.** Find the mode.
 A. 31 B. 7 C. 12 D. 7

HACKING MATHEMATICS

The table at right shows the distribution of ACT Math scores for a sample of students enrolled in MGF1106, Fall 1999. The data is authentic. Refer to the table for exercises 15 - 18.

ACT Math	freq
15	1
16	1
17	10
18	7
19	13
20	9
21	7
22	7
23	13
24	4
25	3
26	4
28	3
29	1
30	1

15. Select the statement that is true.

- A. 17 students had scores of 10.
- B. 42 students had scores of 13.
- C. A and B are both true.
- D. A, B and C are all false.

16. Find the mean ACT score (answers may be rounded).

- A. 21 B. 23 C. 22 D. 24

17. Find the median ACT score (answers may be rounded).

- A. 21 B. 23 C. 22 D. 24

18. Find the mode.

- A. 19 B. 23 C. 13 D. The mode is not unique.

UNIT 2 MODULE 9

The table at right shows the relative frequency (R.F.) distribution of a population of retail Store employees according to hourly wage.

Wage	R.F.
\$7.50	.49
8.00	.12
8.50	.17
9.00	.09
9.50	.07
10.00	.06

19. Find the mean wage.
 A. \$8.75 B. \$8.16 C. \$8.42 D. \$7.92
20. Find the median wage.
 A. \$7.50 B. \$8.00 C. \$8.50 D. \$7.75

ANSWERS TO LINKED EXAMPLES

EXAMPLE 2.9.2 mode = 15, mean = 14.8, median = 15 Any of these three results could be used to represent the “typical” number on the list.

EXAMPLE 2.9.3 Mean = 78.4, median = 80, mode = 68

EXAMPLE 2.9.6 Mode = 10, mean \approx 8.46, median = 9

EXAMPLE 2.9.7 1. Mode = 20, mean = 20.03, median = 20 2. False

EXAMPLE 2.9.8 Mean = 7.94 median = 8 mode = 8

EXAMPLE 2.9.9 Mode = \$0, mean = \$61.55, median = \$50

EXAMPLE 2.9.10 More = \$0, mean = \$16,784.58, median = \$50

EXAMPLE 2.9.11 The mean is the least accurate representation .

ANSWERS TO PRACTICE EXERCISES

1. C 2. D 3. A 4. B 5. C 6. A
 7. B 8. B 9. D 10. B 11. B 12. B
 13. B 14. C 15. D 16. A 17. A 18. D
 19. B 20. B