

2-stratifolds with fundamental group \mathbb{Z}

J. C. Gómez-Larrañaga* F. González-Acuña† Wolfgang Heil‡

Abstract

2-stratifolds are a generalization of 2-manifolds that occur as objects in applications such as in TDA. These spaces can be described by an associated bicoloured labelled graph. In previous papers we obtained a classification of 1-connected trivalent 2-stratifolds. In this paper we classify trivalent 2-stratifolds that have fundamental group \mathbb{Z} . This classification implicitly gives an efficient algorithm that applies to a bicoloured labelled graph to decide whether the associated 2-stratifold has fundamental group \mathbb{Z} .^{1 2}

1 Introduction

A (closed) 2-stratifold is a compact connected 2-dimensional cell complex X that can be constructed from a disjoint union of compact (connected) 2-manifolds W^2 and a disjoint union $X^{(1)}$ of circles by attaching each component of ∂W^2 to $X^{(1)}$ via a covering map $\psi : \partial W^2 \rightarrow X^{(1)}$, with $\psi^{-1}(x) > 2$ for $x \in X^{(1)}$. These 2-stratifolds form a special class of 2-dimensional stratified spaces. Matsuzaki and Ozawa [14] defined and studied a slightly more general class of 2-dimensional stratified spaces, which they call *multibranch surfaces*, by allowing boundary curves, i.e. considering covering maps $\psi : \partial W' \rightarrow X^{(1)}$, where $\partial W'$ is a sub collection of the components of ∂W^2 . A 2-stratifold X can be described by its bicoloured labelled graph Γ_X , whose white vertices are the components W^2 , the black vertices are the components of $X^{(1)}$, and an edge is a component of $\partial W^2 \cap X^{(1)}$, where the label on an edge is the degree of the attaching map $\psi : \partial W^2 \rightarrow X^{(1)}$.

2-stratifolds arise as the nerve of certain Lusternick-Schnirelman type decompositions of 3-manifolds into pieces where they determine whether the \mathcal{G} -category of the 3-manifold is 2 or 3 ([6]). They are related to *foams*, which include special spines of 3-dimensional manifolds and which have been studied by Khovanov [11] and Carter [3]. Matsuzaki and Ozawa showed that every 2-stratifold X embeds in \mathbb{R}^4 , however X embeds into some orientable closed 3-manifold if and only if the singular set $X^{(1)}$ satisfies a certain regularity condition. In [9] we showed that very few 2-stratifolds occur as spines of closed 3-manifolds.

Graphs can be thought of as being 1-stratifolds and persistent graphs occur in Topological Data Analysis [12]. The use of 2-dimensional stratified spaces in TDA has been proposed in

*Centro de Investigación en Matemáticas, A.P. 402, Guanajuato 36000, Gto. México. jcarlos@cimat.mx

†Instituto de Matemáticas, UNAM, 62210 Cuernavaca, Morelos, México and Centro de Investigación en Matemáticas, A.P. 402, Guanajuato 36000, Gto. México. fico@matem.unam.mx

‡Department of Mathematics, Florida State University, Tallahassee, FL 32306, USA. heil@math.fsu.edu

¹AMS classification numbers: 57N10, 57M20, 57M05

²Key words and phrases: 2-stratifolds, fundamental group

the literature [1], with a concrete example given in [2]. It seems likely that 2-stratifolds will be used in a similar fashion as models for applications. For example they occur in the study of the energy landscape of cyclo-octane [13], and 2-stratifolds with boundary and foams with and without boundary occur in the study of boundary singularities produced by the motion of soap films [5].

Closed 2-manifolds are classified in terms of their fundamental groups, but for a given 2-stratifold there are infinitely many non-homeomorphic 2-stratifolds with the same fundamental group. Since the group can be computed from the labelled graph one would like to obtain conditions on the type of bicoloured labelled graphs that correspond to a given fundamental group. As a first step in this direction we obtained in [7] conditions on the labelled graph Γ_X such that the associated 2-stratifold X is 1-connected and in [8] we give a complete classification of 1-connected trivalent 2-stratifolds in terms of their graphs. The term *trivalent* means that each point of $X^{(1)}$ has a neighborhood where three sheets meet. For these 2-stratifolds we developed an efficient algorithm to decide whether they are simply connected and implemented the algorithm in [10].

The ultimate goal is to obtain a classification of all 2-stratifolds in terms of their labelled graphs. In particular, with a view towards applications, one would like to have efficient algorithms for deciding whether a given 2-stratifold is of a certain type. In this paper we consider 2-stratifolds X whose fundamental group is infinite cyclic. In sections 3 and 4 we obtain necessary and sufficient conditions on X and the type of the graph Γ_X such that $\pi_1(X) = \mathbb{Z}$. However for a general 2-stratifold graph Γ_X it is difficult to obtain conditions on the labelling that guarantee that X has fundamental group \mathbb{Z} . We are able to do this in section 6 for trivalent 2-stratifolds, after obtaining a classification for an interesting type of trivalent graphs, the echinus graphs, in section 5. The main Theorem in section 6 provides a complete classification, in terms of conditions that can be read off from the labelled graph, of trivalent 2-stratifolds with fundamental group \mathbb{Z} .

2 2-stratifolds and their graphs

We first review the basic definitions and some results given in [7] and [8]. A *2-stratifold* is a compact, Hausdorff space X that contains a closed (possibly disconnected) 1-manifold $X^{(1)}$ as a closed subspace with the following property: Each point $x \in X^{(1)}$ has a neighborhood homeomorphic to $\mathbb{R} \times CF$, where CF is the open cone on F for some (finite) set F of cardinality > 2 and $X - X^{(1)}$ is a (possibly disconnected) 2-manifold.

A component $B \approx S^1$ of $X^{(1)}$ has a regular neighborhood $N(B) = N_\pi(B)$ that is homeomorphic to $(Y \times [0, 1]) / (y, 1) \sim (h(y), 0)$, where Y is the closed cone on the discrete space $\{1, 2, \dots, d\}$ and $h : Y \rightarrow Y$ is a homeomorphism whose restriction to $\{1, 2, \dots, d\}$ is the permutation $\pi : \{1, 2, \dots, d\} \rightarrow \{1, 2, \dots, d\}$. The space $N_\pi(B)$ depends only on the conjugacy class of $\pi \in S_d$ and therefore is determined by a partition of d . A component of $\partial N_\pi(B)$ corresponds to a summand of the partition determined by π . Here the neighborhoods $N(B)$ are chosen sufficiently small so that $N(B)$ is disjoint from $N(B')$ for disjoint components B and B' of $X^{(1)}$.

For a given 2-stratifold X there is an associated bipartite graph $\Gamma = \Gamma_X$ embedded in X as follows:


The white vertices w_i of the graph Γ are the components W_i of the 2-manifold $M := \overline{X - \cup_j N(B_j)}$ where B_j runs over the components of $X^{(1)}$. The black vertices b_j are the $N(B_j)$'s. An edge e_{ij} is a component E_{ij} of ∂M ; it joins a white vertex w_i with a black vertex b_j if $E_{ij} = W_i \cap N(B_j)$. Note that the number of boundary components of W_i is the number of adjacent edges of w_i .

We label the graph Γ by assigning to a white vertex w the genus g of W and by labelling an edge e by k , where k is the summand of the partition π corresponding to the component E of $\partial N_\pi(B)$. (Here we use Neumann's [15] convention of assigning negative genus g to nonorientable surfaces; for example the genus g of the projective plane or the Moebius band is -1 , the genus of the Klein bottle is -2). Note that the partition π of a black vertex is determined by the labels of the adjacent edges.

In sections 5 and 6 we obtain a classification of trivalent 2-stratifolds in terms of their labelled graphs. Here a 2-stratifold X and its labeled bicoloured graph Γ_X are defined to be *trivalent*, if each black vertex b is incident to either three edges each with label 1 or to two edges, one with label 1, the other with label 2, or b is a terminal vertex with adjacent edge of label 3. This means that a neighborhood of a point of a component B of the 1-skeleton $X^{(1)}$ has 3 sheets; the permutation $\pi : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ of the regular neighborhood $N(B) = N_\pi(B)$ has partition $1 + 1 + 1$ or $1 + 2$ or 3 .

Notation. If Γ is a bipartite labelled graph corresponding to the 2-stratifold X we let $X_\Gamma = X$ and $\Gamma_X = \Gamma$. If W is of genus 0, we do not label w .

Another description of $X = X_\Gamma$ is as a quotient space $W \cup_\psi X^{(1)}$, where $W = \cup W_i$ and where $\psi : \partial W \rightarrow X^{(1)}$ is a covering map (and $|\psi^{-1}(x)| > 2$ for every $x \in X^{(1)}$). For a component $E \approx S^1$ of ∂W the label m on the corresponding edge e then corresponds to the attaching map $\psi(z) = z^m$. An example is given in the picture below.



If Γ is a tree, then the labeled graph determines X uniquely. If Γ is not a tree then we need some additional data for Γ which consists of a cocycle $\kappa \in H^1(\Gamma; \mathbb{Z}_2) = \text{Hom}(H_1(X), \mathbb{Z}_2)$ together with a function λ assigning to each edge e of Γ value $+1$ or -1 such that, for every simple cycle c whose edges are e_1, \dots, e_n , $\lambda(e_1) \cdot \dots \cdot \lambda(e_n) = \kappa([c])$. Here we identify \mathbb{Z}_2 , the cyclic group of order 2, with $\{-1, 1\}$ under multiplication.

One may assume such an evaluation λ is constructed as follows: take a maximal tree T in Γ , and let $\lambda(e) = 1$ if e is an edge of T and $\lambda(e) = \kappa([c])$ if e is in $\Gamma - T$ and c is the cycle of $T \cup e$. With this evaluation, the labelled graph Γ together with a cocycle κ determines X_Γ uniquely.

Thus if Γ is homotopy equivalent to S^1 there is at most one (arbitrarily chosen) edge e in the simple closed cycle C of Γ with $\lambda(e) = -1$.

The fundamental group of X_Γ can be computed from the labelled graph Γ . Generators of $\pi_1(X_\Gamma)$ are the generators of $\pi_1(W^2)$ for each white vertex w , one generator for each black vertex b (also denoted by b), and one generator t for each edge outside a maximal tree T of Γ . Relations are those of $\pi_1(W^2)$, $b^m = c$, for each edge $c \in T$ between w and b with label $m \geq 1$, and $t^{-1}ct = b^{\epsilon m}$, for each edge $c \in G - T$ between w and b with label m ($\epsilon = \pm 1$).

In [7] we showed the following.

Proposition 1. *There is a retraction $\rho : X_\Gamma \rightarrow \Gamma_X$ such that $\rho^{-1}(b)$ is a singular curve $B \in X^{(1)}$ and $\rho^{-1}(w)$ is a 2-manifold W^2 .*

For simplicity, if b is a black vertex of Γ_X , we will sometimes write b instead of $\rho^{-1}(b)$. Also we will write b instead of $[b] \in \pi_1(X_\Gamma)$.

For a given labelled graph Γ we often prune away certain edges and vertices to obtain a subgraph Γ_0 such that there is an epimorphism $\pi_1(X_\Gamma) \rightarrow \pi_1(X_{\Gamma_0})$ as follows: For a subgraph Γ'_0 of Γ let $Y' = \rho^{-1}(\Gamma'_0)$. This is almost a 2-stratifold, except that Y' has possibly boundary curves corresponding to edges of $st(\Gamma'_0) - \Gamma'_0$, where $st(\Gamma'_0)$ is the star of Γ'_0 in Γ . Let Y be the quotient of X_Γ obtained by collapsing the closure of each component of $X_\Gamma - Y'$ to a point i.e. Y is obtained from Y' by attaching disks to its boundary curves. Then Y is a 2-stratifold, $Y = X_{\Gamma_0}$, where Γ_0 is the union of Γ'_0 and the labeled edges (with their vertices) of $st(\Gamma'_0) - \Gamma'_0$ which are adjacent to a black vertex of Γ'_0 . Then the quotient map $X_\Gamma \rightarrow X_{\Gamma_0}$ induces surjections of fundamental groups and first homology groups.

Using Proposition 1 and this pruning construction we show in [7] the following

Proposition 2. *If X is simply connected, then Γ_X is a tree, all white vertices of Γ_X have genus 0, and all terminal vertices are white.*

3 The homotopy type of the graph Γ_X

In this section we show that a necessary condition for $\pi_1(X_\Gamma)$ to be infinite cyclic is that the associated graph is homotopy equivalent to a circle, the labels on all white vertices are 0 and all terminal vertices are white.


In the following Lemma S_g denotes a closed surface of genus g .

Lemma 1. *Let $\Gamma = \Gamma_X$ be any graph of a 2-stratifold $X = X_\Gamma$.*

- (1) *If Γ has (at least) two black terminal vertices then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \mathbb{Z}_m * \mathbb{Z}_n$ for some $m, n \geq 3$.*
- (2) *If Γ has a black terminal vertex and contains a white vertex of genus g then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \mathbb{Z}_m * \pi_1(S_g)$ for some $m \geq 3$.*

(3) If Γ contains two white vertices of genera g_1, g_2 then there is an epimorphism $\pi_1(X_\Gamma) \rightarrow \pi_1(S_{g_1}) * \pi_1(S_{g_2})$.

Proof. (1) Let L be a linear subgraph of Γ_X with terminal black vertices b_1, b_2 and prune Γ at L . In the resulting graph construct L_{mn} by deleting all edges (together with their white vertices) from each interior black vertex b , then adjoining to b one edge with label 1 (together with a white vertex) and finally changing the labels of all white vertices to 0 (see Figure 3). (This has the effect of killing $\rho^{-1}(b)$ in $\pi(X_\Gamma)$). Then there is a surjection $\pi_1(X_\Gamma) \rightarrow \pi_1(X_{L_{mn}}) \cong \mathbb{Z}_m * \mathbb{Z}_n$.



- (2) Let L be a linear subgraph of Γ_X with terminal black vertex b and white vertex w of genus g . Construct a graph L_m as in (1) with terminal vertices b and w . Then there is a surjection $\pi_1(X_\Gamma) \rightarrow \pi_1(X_{L_m})$. If L_m is not a one edge graph, then $\pi_1(X_{L_m}) \cong \mathbb{Z}_m * \pi_1(S_g)$, if L_m is a one edge graph then $\mathbb{Z}_m * \pi_1(S_g)$ is a quotient of $\pi_1(X_{L_m})$.
- (3) In the proof of (2) replace L by a linear subgraph of Γ_X with a terminal white vertex of genus g_1 and a terminal white vertex of genus g_2 . □

Lemma 2. Let $\Gamma = \Gamma_X$ be a tree of a 2-stratifold X_Γ .

- (1) If Γ_X has at most one black terminal vertex and all white vertices are of genus 0 then $H_1(X_\Gamma)$ is finite.
- (2) If Γ_X has no black terminal vertices, contains at most one white vertex of genus -1 and all other white vertices are of genus 0, then $H_1(X_\Gamma)$ is finite.

Proof. (1)(a) Suppose Γ has no black terminal vertices. The proof is by induction on the number of white vertices. Let w be a terminal (white) vertex with edge e with label m between w and a black vertex b . If the degree of b is > 2 then $\Gamma_Y = \Gamma - (w \cup e)$ is as in the lemma and by induction $H_1(Y)$ is finite. Therefore $H_1(X) \cong H_1(Y) / \langle b^m \rangle$ is finite.

If the degree of b is 2 let $\Gamma_Y = \Gamma_X - (st(b) \cup \{w\})$. Then $H_1(Y) \cong H_1(X) / \langle b \rangle$, where b is a non-trivial element of the torsion subgroup (since $m \geq 1$). By induction $H_1(Y)$ is finite and it follows that $H_1(X)$ is finite.

(1)(b) Now suppose Γ_X has exactly one black terminal vertex b . If b has finite order n in $H_1(X_\Gamma)$, let Γ_Y be obtained from Γ_X by attaching an edge e with label n , together with a white vertex of genus 0, to b . Then Γ_Y is as in (1)(a) and so $H_1(X_{\Gamma_X}) = H_1(X_{\Gamma_Y})$, is finite.

We claim that b does not have infinite order in $H_1(X_\Gamma)$. To see this let Γ_Y be the double of Γ_X along b (i.e. Γ_Y is obtained from two disjoint copies of Γ_X by identifying their copies of b). If b has infinite order in $H_1(X_\Gamma)$, then $\langle b \rangle$ is an infinite cyclic subgroup of $H_1(\Gamma_Y) = H_1(X_\Gamma) \oplus H_1(X_\Gamma)$ amalgamated along $\langle b \rangle$. This is a contradiction, since Γ_Y is as in (1)(a).

(2) Let $\Gamma_{Y_1}, \dots, \Gamma_{Y_r}$ be the trees of $\Gamma_X - st(w)$ where w is the white vertex of genus -1 . By (1), $H_1(\Gamma_{Y_i})$ is finite ($i = 1, \dots, r$). Now $H_1(X_\Gamma) \cong H_1(Y_1) \oplus \dots \oplus H_1(Y_r) \oplus \langle a \rangle / \langle b_1^{m_1} \dots b_r^{m_r} a^2 = 1 \rangle$, where the b_i 's are the black vertices of $st(w)$ and the m_i 's are the labels of the edges incident to w . It follows that $a^2 \in H_1(Y_1) \oplus \dots \oplus H_1(Y_r)$ and so a has finite order in $H_1(X_\Gamma)$. □

Lemma 3. *Let $\Gamma = \Gamma_X$ be a graph of a 2-stratifold X_Γ that is homotopy equivalent to S^1 .*

(1) *If Γ_X has a white vertex of genus g then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \mathbb{Z} * \pi_1(S_g)$.*

(2) *If Γ_X has a black terminal vertex then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \mathbb{Z} * \mathbb{Z}_m$ for some $m \geq 3$.*

Proof. (1) If the white vertex w of genus g is a vertex of the cycle C of Γ let C^* be the labelled graph obtained from C by adjoining to each black vertex an edge (together with a white vertex) of label 1 and assigning genus 0 to all white vertices of C^* , except for w . Then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \pi(X_{C^*}) = \mathbb{Z} * \pi_1(S_g)$.

If the white vertex w of genus g is not in C connect w to C by a linear graph L in Γ such that w is a terminal vertex of L and $L \cap C$ is a vertex v . Let C^* be as above and let L^* be obtained from $C^* \cup L$ by adjoining to each black vertex of L an edge (together with a white vertex) of label 1 and assigning genus 0 to all white vertices of L^* , except for w . Then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \pi(X_{L^*}) = \mathbb{Z} * \pi_1(S_g)$.

(2) Let L be a linear subgraph of Γ_X connecting the terminal black vertex b (and with terminal edge labelled $m \geq 3$) to a vertex of C and construct L^* as in (1). Then there is an epimorphism $\pi_1(X_\Gamma) \twoheadrightarrow \pi(X_{L^*}) = \mathbb{Z} * \mathbb{Z}_m$. \square

Proposition 3. *If $\pi_1(X_\Gamma) = \mathbb{Z}$ then Γ_X is homotopy equivalent to S^1 , all white vertices of Γ are of genus 0, and all terminal vertices are white.*

Proof. The retraction $\rho : X_\Gamma \rightarrow \Gamma_X$ induces an epimorphism $\pi(X_\Gamma) \twoheadrightarrow \pi(\Gamma_X)$ and so Γ_X is a tree or homotopy equivalent to S^1 .

By Lemma 1(1) Γ_X has at most one black terminal vertex.

If Γ_X has one black terminal vertex then by Lemma 1(2) all white vertices have genus 0. Then by Lemma 2(1) Γ is not a tree.

If Γ_X has no black terminal vertices then by Lemma 1(3) all white vertices have genus 0, except possibly for one of genus -1 . Then by Lemma 2(2) Γ is not a tree.

Hence Γ is homotopy equivalent to S^1 and by Lemma 3 all white vertices have genus 0 and all terminal vertices are white. \square

4 The general case of $\pi(X_\Gamma) \cong \mathbb{Z}$

In this section we obtain the main result in the case when Γ is any labelled graph satisfying the necessary conditions of Proposition 3. By [8], the requirements in Theorem 1 below that certain stratifolds Y_Γ be 1-connected can be determined from the labelled graphs Γ_Y .

We first obtain a Lemma that applies to any labelled graph Γ :

Lemma 4. *Let Γ be a labelled graph and suppose b is a black vertex of Γ of degree $d \geq 2$ such that $\rho^{-1}(b)$ is contractible in X_Γ . Then $\pi(X_\Gamma) = \pi(X_{\Gamma_1}) * \dots * \pi(X_{\Gamma_c}) * F_r$ where $\Gamma_1, \dots, \Gamma_c$ are the components of $\Gamma_X - stb$ and F_r is the free group of rank $r = d - c$.*

Proof. Let Y be the space obtained from X_Γ by collapsing $\rho^{-1}(b)$ to a point y_0 . Then there is an obvious map $p : Y \rightarrow \Gamma_X$ such that $p^{-1}(\Gamma_X - stb) = X_{\Gamma_1} \cup \dots \cup X_{\Gamma_c}$ and $p^{-1}(b) = y_0$. Since $\rho^{-1}(b)$ is contractible, $\pi(Y) \cong \pi(X_\Gamma)$. Now let $q : \Gamma_X \rightarrow \Gamma_Y$ be the quotient map, where Γ_Y is obtained from Γ_X by collapsing each component Γ_i to a point y_i and let T be a maximal tree in Γ_Y . Then $(qp)^{-1}(T)$ is homotopy equivalent to the wedge $X_{\Gamma_1} \vee \dots \vee X_{\Gamma_c}$. Since there are $r = d - c$ edges in $\Gamma_Y - T$ it follows that $Y = (qp)^{-1}(\Gamma_Y)$ is homotopy equivalent to $X_{\Gamma_1} \vee \dots \vee X_{\Gamma_c} \vee \bigvee_r S^1$. \square

Recall from Proposition 3 that a necessary condition for $\pi_1(X_\Gamma) = \mathbb{Z}$ is that Γ_X is homotopy equivalent to S^1 , all white vertices of Γ are of genus 0, and all terminal vertices are white. The next lemma lists more necessary conditions. By the *cycle* C of Γ we mean the subgraph that is homeomorphic to S^1 .

Lemma 5. *Assume $\pi(X_\Gamma) \cong \mathbb{Z}$, then*

- (a) Γ is not homeomorphic to S^1 .
- (b) There is at least one black vertex belonging to the cycle C of Γ whose degree is > 2 .
- (c) For each black vertex b , $\rho^{-1}(b)$ is homotopic to 0.
- (d) Let B be a collection of black vertices containing at least one $b_1 \in C$. Then every component of $\Gamma - stB$ is 1-connected.

Proof. We know that Γ_X is homotopy equivalent to S^1 , all white vertices of Γ are of genus 0, and all terminal vertices are white.

(a) We show that if $\Gamma_X = C$, a cycle, then $\pi(X_\Gamma)$ is infinite but not $\cong \mathbb{Z}$.

Starting at a white vertex w_1 and reading the edges (counterclockwise) as $w_1 - b_1 - w_2 - b_2 - \dots - w_k - b_k - w_1$ with edge labels $m_1, n_1, \dots, m_k, n_k$ we obtain a presentation $\pi_1(X_\Gamma) = \langle b_1, \dots, b_k, t \mid b_i^{n_i} = b_{i+1}^{m_{i+1}}, t^{-1} b_k^{n_k} t = b_1^{\epsilon m_1}, i = 1, \dots, k-1 \rangle$, where $n_i, m_i \geq 1$ for $i = 1, \dots, k$ and $\epsilon = \pm 1$. This group is an HNN extension of the nontrivial group (an iterated free product with amalgamation) $B := \langle b_1, \dots, b_k \mid b_i^{n_i} = b_{i+1}^{m_{i+1}}, i = 1, \dots, k-1 \rangle$ and $b_1^{m_1}$ and $b_k^{n_k}$ are of infinite order in B . Hence $\pi(X_\Gamma) \not\cong \mathbb{Z}$.

(b) If each black vertex of C has degree 2 then pruning Γ at each white vertex of C yields a graph Γ_C homeomorphic to S^1 . Since $\rho_* : \pi(X_\Gamma) \rightarrow \pi(X_{\Gamma_C})$ is surjective this contradicts (a).

(c) Since \mathbb{Z} is Hopfian, the epimorphism $\rho_* : \pi(X_\Gamma) \rightarrow \pi(\Gamma)$ is an isomorphism and since $\rho(\rho^{-1}(b))$ is a point, the circle $\rho^{-1}(b)$ is nullhomotopic in X_Γ .

(d) From (c) and Lemma 4 it follows that X_Λ is 1-connected for every component Λ of $\Gamma - st(b_1)$. If Λ contains some $b \in B$, then since $\rho^{-1}(b)$ is contractible in X_Λ it follows again from Lemma 4 that every component of $\Gamma - st\{b_1, b\}$ is 1-connected. Now (d) follows from induction. \square

We remark that in (a) $H_1(X_\Gamma) = \mathbb{Z} \oplus G_d$ where G_d is a group of order $d := n_1 n_2 \dots n_k - \epsilon m_1 m_2 \dots m_k$, the determinant of the k by k matrix representing $G_d = \langle b_1, \dots, b_k, [b_i, b_j] = 1, b_i^{n_i} = b_{i+1}^{m_{i+1}}, b_k^{n_k} = b_1^{\epsilon m_1}, i = 1, \dots, k-1 \rangle$. If X_Γ is trivalent (i.e. $0 \leq m_i, n_i \leq 2, m_i + n_i = 3$) and $k > 1$ then d is not ± 1 and so $G_d \neq 0$.

The next proposition gives conditions that are necessary and sufficient.

Proposition 4. *Let $\Gamma = \Gamma_X$ be a bicoloured labelled graph homotopy equivalent but not homeomorphic to S^1 and such that all white vertices have genus 0 and all terminal vertices are white. Let B be a collection of black vertices of Γ containing a vertex b_1 of C and such that $\rho^{-1}(b)$ is homotopic to 0 in X_Γ for each $b \in B$. Then $\pi(X_\Gamma) \cong \mathbb{Z}$ if and only if all components of $X_\Gamma - stB$ are 1-connected.*

Proof. By Lemma 5(c) the condition is necessary. Considering $b_1 \in B$ we have by Lemma 4 that $\pi(X_\Gamma) = \pi(X_{\Gamma_1}) * \dots * \pi(X_{\Gamma_c}) * \mathbb{Z}$, where $\Gamma_1, \dots, \Gamma_c$ are the components of $\Gamma_X - st b_1$. If $b \in B \cap \Gamma_i$ for some $1 \leq i \leq n$, then again by Lemma 4, $\Gamma - stb$ splits $\pi(X_{\Gamma_i})$ into a free product. Doing this successively for all $b \in B$ we obtain $\pi(X_\Gamma) = (*\pi(X_{\Gamma_{ij}})) * \mathbb{Z}$, where the Γ_{ij} are the components of $X - stB$. By assumption these are 1-connected and therefore $\pi(X_\Gamma) = \mathbb{Z}$ \square

We now state the main Theorem of this section.

Theorem 1. Let Γ be a bicoloured labelled graph. Then $\pi(X_\Gamma) \cong \mathbb{Z}$ if and only if the following four conditions hold.

- (1) Γ is homotopy equivalent but not homeomorphic to S^1 .
- (2) All white vertices have genus 0 and all terminal vertices are white.
- (3) The cycle of Γ has a black vertex b of degree $d > 2$ and $\rho^{-1}(b)$ is contractible in X_Γ .
- (4) X_{Γ_i} ($i = 1, \dots, d-1$) is 1-connected where $\Gamma_1, \dots, \Gamma_{d-1}$ are the components of $\Gamma - st(b)$.

Proof. Proposition 3 and Lemma 5 show that the conditions are necessary. Suppose (1)-(4) hold. Then for a black vertex b as in (3) and (4) it follows from Lemma 4 that $\pi(X_\Gamma) = \mathbb{Z}$. \square

5 Trivalent echinus graphs

In this section we consider bicoloured trivalent graphs $\Gamma = \Gamma_X$ that are homotopy equivalent to S^1 and such that the closure of each component of $\Gamma - C$ is a linear graph, where C is the cycle of Γ .

First we list some necessary conditions for $\pi(X_\Gamma) \cong \mathbb{Z}$.


Lemma 6. Assume $\pi(X_\Gamma) \cong \mathbb{Z}$ and Γ is trivalent, then

- (a) If all terminal edges of Γ disjoint from C have label 2, then $\Gamma - C$ contains no white branch vertices.
- (b) Let v be a non-terminal vertex of Γ and let L be linear graph that is the closure of a component of $\Gamma - \{v\}$ and contains a terminal edge of Γ of label 2. If v is a white vertex then L is a linear 1212...12 graph. If v is a black branch vertex then L is a linear 11212...12 graph.
- (c) If all terminal edges of Γ have label 2 then C contains no white branch vertices.

Proof. (a) Suppose w is a white branch vertex in $\Gamma - C$ and A_1, \dots, A_k, D are the closures of the components of $\Gamma - w$, where D contains C . All terminal edges of $A_1 \cup \dots \cup A_k$ have label 2, and so $H_1(X_{A_1 \cup \dots \cup A_k})$ is not 0. The retraction $\rho : X_D \rightarrow D$ shows that $H_1(X_D)$ contains a \mathbb{Z} -summand and collapsing in X_Γ the circle $\rho^{-1}w \cap \rho^{-1}(D)$ to a point we see that $H_1(X_\Gamma)$ has $H_1(X_D) + H_1(X_{A_1 \cup \dots \cup A_k})$ as a quotient, hence $H_1(X_\Gamma) \not\cong \mathbb{Z}$.

(b) Let B be the set of black vertices of $\Gamma - L$. Then L is a component of $\Gamma - B$ and is 1-connected by Lemma 5(d). Since all edge labels of L are 1 or 2 (and in the second case the first label is 1) the claim follows from Theorem 3 of [GGH].

(c) By (a) we know that $\Gamma - C$ has no white branch vertices. If a component of $\Gamma - C$ contains a black branch vertex let b be an outermost such vertex. Then the closures of the two components that do not meet C are linear 1212...12 graphs of length m_1, m_2 with $m_2 \geq m_1$, say. Pruning away $L_2 \cup st(b)$ as in the figure below does not change the fundamental group. Hence we may assume that the closure of every component of $\Gamma - C$ is a linear graph as in (b).



Let w be a white vertex of C of degree > 2 . We will show that $H_1(X_\Gamma) \not\cong \mathbb{Z}$.

If w' is another such white vertex prune away all subgraphs other than C that meet w' . For the resulting graph Γ' there is an epimorphism $H_1(X_\Gamma) \rightarrow H_1(X_{\Gamma'})$. So may assume that C has exactly one vertex w of degree > 2 .

If $\text{deg}(w) > 3$, prune away all subgraphs other than C that meet w , except for one. Again there is an epimorphism $H_1(X_\Gamma) \rightarrow H_1(X_{\Gamma'})$. So may assume that $\text{deg}(w) = 3$.

Note that the closure of each component of $\Gamma - C$ is a linear graph as in (2.2). If $L_r = b_1 - w_1 - b_2 - w_2 - \dots - b_r - w_r$ we may prune away the part after w_2 (the effect is killing $\rho^{-1}(b_1^2)$ in H_1). So may assume that all $r = 2$. By the same argument, if $L = w - b_1 - w_1 - \dots - b_r - w_r$ we may (by pruning L at w_1) assume that $L = w - b_1 - w_1$.

If there is an edge e of $C - stw$ with label 2 let w' be the white vertex incident with e and let γ be the path on C , containing e with endpoints w and w' . Then if A is the union of γ with the components of $\Gamma - C$ whose closure intersects γ , we have $H_1(X_A) \neq 0$ since all the terminal edges of A have label 2. By killing all the loops in $X_\Gamma - X_A$ we see that $H_1(X_\Gamma)$ has as quotient $\mathbb{Z} + H_1(X_A)$ which is not \mathbb{Z} .

If all edges of $C - stw$ have label 1 we compute $H_1(X_\Gamma)$: Let e_1 and e_2 be the labels of the edges in C adjacent to w (the other edge $w - b_1$ adjacent to w has label 1). In $H_1(X_\Gamma)$ one has $2b_1 = 0$ (where b corresponds to a black vertex of C) and $(e_2 - e_1)b = b_1$. Adjoining the relation $2b = 0$ (this is only needed if there is no black branch vertex) one obtains as a quotient of $H_1(X_\Gamma)$ the abelian group $(t, b, b_1 : 2b_1 = 0, 2b = 0, (e_2 - e_1)b = b_1)$ which is $\mathbb{Z} + \mathbb{Z}_2$ (no matter what e_1 and e_2 are). \square

A p -string (of length p) is an oriented linear graph $w_0 - b_1 - w_1 - b_2 - \dots - b_k - w_k$ with all white vertices w_i of genus 0, successive edge labels $1212 \dots 12$ (starting at w_0) and with p labels of 2.

A q -string is such an oriented linear graph with successive edge labels $2121 \dots 21$ (starting at w_0) with q labels of 2.

If L is a trivalent linear graph for which X_L is 1-connected, then Theorem 3 of [6] implies that L does not contain a linear subgraph consisting of a q -string followed by a p -string for $q, p > 0$. Thus we define $L(p, q)$ to be the linear graph consisting of a p -string followed by a q -string, i.e. $L(p, q)$ is the trivalent linear graph with terminal vertices white and edge labels $12 \dots 1221 \dots 21$, but we allow p or q to be 0.

We say that a q -string is a *terminal* string of Γ if w_k is a terminal (white) vertex of Γ . Deleting from Γ_X all terminal q -strings, except for their initial vertices w_0 , does not change the fundamental group of X_Γ . For the computation of the fundamental group it therefore suffices to restrict to *pruned* graphs, that are graphs without terminal q -strings for $p > 0$.

If Γ is a trivalent pruned graph such that the closure of each component of $\Gamma - C$ is a linear graph and $\pi(X_\Gamma) = \mathbb{Z}$ then from Lemma 6(c) we know that C contains no white branch vertices and then it follows from Proposition 1 and Theorem 1 that all components of $\Gamma - stB$ are 1-connected $L(p, q)$ -graphs, where B is the set of black branch vertices of C . Furthermore, since Γ is pruned, the closure of each component of $\Gamma - C$ is a linear graph $L(r, 0)$ -graph. We call these graphs *echinus*-graphs.

Definition. The echinus graph $E = E(p_1, q_1, r_1; \dots; p_n, q_n, r_n)$ is the trivalent labelled graph Γ with the following properties:

- (1) Γ is homotopy equivalent, but not homeomorphic to S^1 .
- (2) All vertices of Γ are of degree 2, except for $n \geq 1$ black vertices of the cycle C of Γ .
- (3) If b_1, \dots, b_n are the successive black branch vertices for a fixed orientation of C and C_1, \dots, C_n

are the successive components of $C-st\{b_1, \dots, b_n\}$, then C_i is the linear graph $L(p_i, q_i)$, with $p_i, q_i \geq 0$, $i = 1, \dots, n$.

(4) The component of $\Gamma - st(b_i)$ that does not intersect C is the linear graph $L(r_i, 0)$ with $r_i \geq 0$, $i = 1, \dots, n$.



The fundamental group of X_E for $E = E(p_1, q_1, r_1; \dots; p_n, q_n, r_n)$ has a presentation

$$\pi(X_E) = \{b_1, \dots, b_n, t : b_j^{2^{r_j}} = 1, b_i^{2^{p_i}} = b_{i+1}^{2^{q_i}}, t^{-1} b_n^{2^{p_n}} t = b_1^{2^{q_1}}, j = 1, \dots, n, i = 1, \dots, n-1\}$$

and $H_1(X_E) = \mathbb{Z} + T$, where T has a relation matrix $M = \begin{pmatrix} A \\ B \end{pmatrix}$ as follows:

A is the $n \times n$ diagonal matrix with the entry in the i th row and column 2^{r_i} ; B is the $n \times n$ matrix (b_{ij}) where $b_{ii} = 2^{p_i}$, $b_{ij} = -2^{q_i}$ if $j = i + 1 \pmod n$ and all other b_{ij} 's are 0.

We first consider echinus graphs that contain no branch vertices that are of distance 1 to a terminal vertex (i.e. all $r_i > 0$) and show:

Proposition 5. *Let $E = E(p_1, q_1, r_1; \dots; p_n, q_n, r_n)$ with $r_i > 0$ for all i . Then $H_1(X_E) = \mathbb{Z}$ if and only if exactly one of $\sum_{i=1}^n p_i$, $\sum_{i=1}^n q_i$ is 0.*

Proof. Recall that $T \cong \mathbb{Z}_{d_1} \oplus \dots \oplus \mathbb{Z}_{d_n}$ with $d_i | d_{i+1}$ and presentation matrix D the diagonal matrix with d_i the entry in the i th row and column. The d_i 's can be computed using elementary ideals as in [4] (Chapt.VIII Sect.4): M is equivalent to D and therefore $(e_k) = E_k(M) = E_k(D) = (d_1, \dots, d_{n-k})$, the subgroup generated by the product $d_1 d_2 \dots d_{n-k}$ if $0 \leq k < n$, where $e_k (\geq 0)$ is the greatest common divisor of the determinants of all $(n-k) \times (n-k)$ submatrices of M . Hence $d_i = e_{n-i} / e_{n-i+1}$ and in particular $T = 0$ if and only if $e_0 = 1$.

For $M = \begin{pmatrix} A \\ B \end{pmatrix}$ as above we have $\det A = 2^{r_1+r_2+\dots+r_n}$ and one computes $\det B = 2^{p_1+p_2+\dots+p_n} - 2^{q_1+q_2+\dots+q_n}$. All the other $n \times n$ submatrices contain a row of A and therefore their determinants are (either 0 or) divisible by 2 (since by assumption each $r_i \neq 0$), and so $e_0 = 1$ if and only if 2 does not divide $2^{p_1+p_2+\dots+p_n} - 2^{q_1+q_2+\dots+q_n}$. This happens if and only if exactly one of $\sum_{i=1}^n p_i, \sum_{i=1}^n q_i$ is 0. \square

Theorem 2. *Let $E = E(p_1, q_1, r_1; p_2, q_2, r_2; \dots; p_n, q_n, r_n)$ be an echinus graph with all $r_i \neq 0$. Then the following are equivalent*

- (a) $\pi X_E = \mathbb{Z}$
- (b) $H_1(X_E) = \mathbb{Z}$
- (c) *Precisely one of $p_1 + p_2 + \dots + p_n$ or $q_1 + q_2 + \dots + q_n$ is 0.*

Proof. By Proposition 5 it suffices to show that if $q_1 + q_2 + \dots + q_n = 0$ but $p_1 + p_2 + \dots + p_n \neq 0$ then $\pi X_E = \mathbb{Z}$.

We may assume $r_1 = \min\{r_1, \dots, r_n\}$. Then eliminating b_2, \dots, b_n in the presentation of πX_E we obtain $\pi X_E = \{b, t : (b_1)^{2^{r_1}} = 1, t^{-1}b_1^{2^m}t = b_1\}$ where $m = p_1 + p_2 + \dots + p_n > 0$.

(*) If $r_1 \leq m$, then $b_1 = 1$ and so $\pi X_E = \{t : \} = \mathbb{Z}$.

If $r_1 > m$, raising the second relation to the power 2^{r_1-m} , we get

$\pi X_E = \pi X_E(p_1, q_1, r_1 - m; p_2, q_2, r_2, \dots, p_n, q_n, r_n)$.


After a number of these reductions of r_1 we obtain

$\pi X_E = \pi X_E(p_1, q_1, r'; p_2, q_2, r_2; \dots; p_n, q_n, r_n)$, where $r' \leq m$, which by (*) is \mathbb{Z} . \square

We now consider echinus graphs that contain branch vertices of distance 1 to terminal vertices. Note that if all r_i are 0, then $\pi(X_E) = 1$.

Let E be an echinus graph with cycle C and let B be the set of black branch vertices in C at distance 1 from a terminal vertex. Assume B is not empty and let A be a component of $E - stB$. If A does not consist of a single (white) vertex, A is as in the figure below and with obvious notation we write

$$A = A(p_1, q_1, r_1; \dots; p_{n-1}, q_{n-1}, r_{n-1}; p_n, q_n), \quad n \geq 1, p_i \geq 0, q_i \geq 0, r_i > 0.$$



Proposition 6. *The following are equivalent :*

- (i) X_A is 1-connected.
- (ii) $H_1(X_A) = 0$.
- (iii) *If $q_s > 0$ for some $s < n$, then $p_t = 0$ for all $t > s$.*

Proof. Clearly (i) implies (ii). We show that (ii) implies (iii). If (iii) does not hold, let $q_s > 0$ and $p_t > 0$ for some $s < t$ and let w_1 (resp. w_2) be the white starting vertex (resp. end vertex) of the q_s -string (resp. p_t -string). Consider the subgraph A' of A that is the union of the path γ in $C \cap A$

from w_1 to w_2 and the components of $E-C$ whose closures intersect γ . Then $\pi(X_{A'})$ is a quotient of $\pi(X_A)$ and $H_1(X_{A'}) \neq 0$ since all the terminal edges of A' have label 2 (recall all $r_i > 0$). This contradicts (ii). Now we show that (iii) implies (1). This is clear if all $q_s = 0$. If $q_s \neq 0$ then (iii) implies that $A = A(p_1, 0, r_1; \dots; p_{s-1}, 0, r_{s-1}; p_s, q_s, r_s; 0, q_{s+1}, r_{s+1}; \dots; 0, q_n)$. Pruning A from the left we obtain a subgraph $A' = A(0, q_s, r_s; 0, q_{s+1}, r_{s+1}; \dots; 0, q_n)$ with $\pi(X_A) = \pi(X_{A'})$ and then pruning A' from the right we obtain $A'' = w$, a single vertex, with $\pi(X_{A'}) = \pi(X_{A''}) = 1$. \square

We now obtain a characterization of 1-connected echinus graphs that contain branch vertices that of distance 1 to terminal vertices.

Theorem 3. *Let E be an echinus graph with cycle C such that the set B of black branch vertices in C at distance 1 from a terminal vertex is not empty. Then X_E is 1-connected if and only if each component of $E-stB$ is a graph of type $A(p_1, 0, r_1; \dots; p_{s-1}, 0, r_{s-1}; p_s, q_s, r_s; 0, q_{s+1}, r_{s+1}; \dots; 0, q_n)$.*

Proof. This follows from Propositions 4 and 6. \square

6 Trivalent graphs

In [8] we classified trivalent graphs that are 1-connected in terms of their graphs. We now describe necessary and sufficient conditions for $\pi(X_\Gamma)$ to be \mathbb{Z} that can be read off the graph for any trivalent graph Γ . We assume that Γ satisfies the first three conditions of Theorem 1 which can readily be detected from the graph. Recall that *pruning* a graph means removing all terminal linear subgraphs with edge labels $21 \dots 21$. This does not change the fundamental group of the associated 2-stratifolds. Thus we also assume that the graph Γ is *pruned* i.e. there are no terminal linear subgraphs with edge labels $21 \dots 21$. To summarize, we assume

- (1) Γ is homotopy equivalent but not homeomorphic to S^1 .
- (2) All white vertices have genus 0 and all terminal vertices are white.
- (3) The cycle of Γ has a black vertex b of degree $d > 2$.
- (4) Γ is pruned.

For convenience we say that a labelled graph Δ is 1-connected if X_Δ is 1-connected. We now reduce Γ to its core as follows.

If $\Gamma-C$ does not contain a black branch vertex of distance 1 to a terminal vertex, we let $\Gamma_0 = \Gamma$ and say that Γ_0 is *core-reduced*. If $\Gamma-C$ contains a black branch vertex of distance 1 to a terminal vertex we let $B_0 = \{b_{01}, \dots, b_{0k}\}$ be the set of all such outermost black branch vertices, i.e. each b_{0i} has distance 1 from a terminal vertex w_{0i} and the component T_{0i} of $\Gamma-(st(b_{0i}) \cup w_{0i})$ that does not meet C does not contain a black branch vertex of distance 1 to a terminal vertex. If some component T_{0i} is not 1-connected let $\Gamma_0 = \emptyset$. If each T_{0i} is 1-connected let $\Gamma'_0 = \Gamma-(st(B_0) \cup \bigcup\{w_{0i}\} \cup \bigcup T_{0i})$. If Γ'_0 is pruned let $\Gamma_0 = \Gamma'_0$, otherwise let Γ_0 be the pruned Γ'_0 . Note that in this case $\pi(X_\Gamma) = \pi(X_{\Gamma_0})$, since $\rho^{-1}(b_{0i})$ is homotopic to 0 in (X_Γ) .

Inductively, if $\Gamma_{n-1}-C$ contains an outermost black branch vertex of distance 1 to a terminal vertex we let $B_{n-1} = \{b_{n-1,1}, \dots, b_{n-1,k_{n-1}}\}$ be the set of all such outermost black branch vertices that each has distance 1 from a terminal vertex $w_{n-1,i}$ and let $T_{n-1,i}$ be the component of $\Gamma-(st(b_{n-1,i}) \cup w_{n-1,i})$. If some $T_{n-1,i}$ is not 1-connected we let $\Gamma_n = \emptyset$. If each $T_{n-1,i}$ is 1-connected we let Γ_n be the pruned graph $\Gamma_{n-1}-(st(B_{n-1}) \cup \bigcup\{w_{n-1,i}\} \cup \bigcup T_{n-1,i})$. We define the *core-reduced* subgraph Γ_C as follows:

$$\Gamma_C = \begin{cases} \emptyset & \text{if } \Gamma_n = \emptyset \text{ for some } n \geq 0, \text{ otherwise} \\ \Gamma_n & \text{for the smallest } n \text{ such that } \Gamma_n - C \text{ does not contain a black branch vertex} \\ & \text{of distance 1 to a terminal vertex.} \end{cases}$$

Note that if $\Gamma_C \neq \emptyset$ then $\pi(X_{\Gamma_C}) = \pi(X_\Gamma)$.

We can now state our main classification Theorem.

Theorem 4. *Let Γ be a bicoloured pruned trivalent graph. Then $\pi(X_\Gamma) = \mathbb{Z}$ if and only if*

- (a) *The underlying graph of Γ is homotopically equivalent, but not homeomorphic, to a circle*
- (b) *All terminal vertices of Γ are white and the genus of any white vertex is 0*
- (c) *The core-subgraph Γ_C is not the empty set*
- (d) *If B is the set of black vertices on the cycle C of Γ_C at distance 1 from a terminal vertex of Γ_C then either*
 - (d1) *B is nonempty and X_Λ is 1-connected for all components Λ of $\Gamma_C - st B$ or*
 - (d2) *B is empty, all branch vertices of Γ_C are black, any edge of Γ_C whose distance to C (or, equivalently, to a terminal vertex of Γ_C) is odd has label 1, and C is alternating.*

Here the cycle C of Γ is alternating if, ignoring the edges incident to branch vertices, it is a 1212...12 cycle with a positive (even) number of edges.

Proof. Assume that $\pi(X_\Gamma) = \mathbb{Z}$. Then (a),(b),(c) and (d1) follow from Theorem 1 and Proposition 4. As for (d2) we know from Lemma 6(a) and (b) that all branch vertices Γ are black. If there is an edge at odd distance d from C with label 2 and w is its white endpoint at distance d from C then, if A is the closure of the component of $\Gamma - w$, X_A is not 1-connected because all terminal edges of A have label 2. This is a contradiction since A occurs as a component as in Lemma 5(d).

By successively applying a pruning move as in the proof of of Lemma 6(c) (pruning off suitable terminal linear subgraphs that does not change the fundamental group), we may assume that the closures of all components of $\Gamma - C$ are disjoint linear subgraphs and, by further pruning, we may assume they have length 3 with edge labels 1, 1, 2. If the labelled graph is ‘‘orientable’’ (i.e. all labels are positive) then it is an echinus graph $E = E(p_1, q_1, 1; p_2, q_2, 1; \dots; p_n, q_n, 1)$ with $n > 1$, $p_i \geq 0, q_i \geq 0$ and by Theorem 2 exactly one of $p_1 + p_2 + \dots + p_n, q_1 + q_2 + \dots + q_n = 0$. This means that C is alternating.

If the labelled graph $E = E^-$ is not orientable then we may assume there is precisely one negative label in the graph and we may assume it is a -1 label of an edge of C adjacent to (and ‘‘to the left’’ of) the black branch vertex b_1 . Then $H_1(E^-)$ is presented by a $2n \times n$ matrix $\begin{pmatrix} D \\ F \end{pmatrix}$ where D is an $n \times n$ diagonal matrix with 2’s on the diagonal and F is an $n \times n$ matrix with determinant $2^{p_1 + \dots + p_n} + 2^{q_1 + \dots + q_n}$. Again, as in the proof of Proposition 5, $H_1(E^-) = \mathbb{Z}$, if and only if exactly one of $p_1 + \dots + p_n, q_1 + \dots + q_n$ is 0, that is, C is alternating.

To prove the converse, suppose (a), (b), (c) and (d) of the Theorem hold. If b is an outermost black branch vertex of Γ_C such that the closure of the two components of $\Gamma_C - b$ which do not intersect C are linear 11212...12 graphs L_1, L_2 of lengths m_1, m_2 respectively with $m_2 \geq m_1 \geq 1$, we prune off L_2 to obtain a labelled graph Γ'_C with fewer such branch vertices. If b is an outermost black branch vertex of Γ_C adjacent to a terminal white vertex w and $L \neq w$ is the closure of the component of $\Gamma_C - b$, delete $L \cup st(b) \cup w$ from Γ_C . If the resulting subgraph is not pruned, eliminate the terminal linear subgraph with edge labels $21 \dots 21$ to get Γ' . In either case $\pi(X_{\Gamma_C}) = \pi(X_{\Gamma'})$. Repeating this pruning process we obtain an echinus graph E with $\pi(X_\Gamma) = \pi(X_{\Gamma_C}) = \pi(X_E)$ and

such that E satisfies (a), (b), (c) and (d) of the Theorem. Now $\pi(X_E) = \mathbb{Z}$ by Theorems 2 and 3. \square

Acknowledgments: J. C. Gómez-Larrañaga would like to thank INRIA Saclay, Francia and CNRS UMI 2001, Laboratorio Solomon Lefschetz, Mexico, for financial support.

References

- [1] P. Bendich , B. Wang , S. Mukherjee, Local homology transfer and stratification learning, In Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms, 1355-1370 (2012).
- [2] P. Bendich, E. Gasparovicy, C.J. Traliez, J. Harer, Scaffoldings and Spines: Organizing High-Dimensional Data Using Cover Trees, Local Principal Component Analysis, and Persistent Homology, arXiv:1602.06245v2 [cs.CG] 27 Feb 2016.
- [3] J.S. Carter, Reidemeister/Roseman-type moves to embedded foams in 4-dimensional space, Series on Knots and Everything 56, New Ideas in Low Dimensional Topology, 1-30 (2015).
- [4] R. Crowell, R. Fox, Introduction to Knot Theory, Dover Publications (2008).
- [5] R.E. Goldstein, J. McTavish, H.K. Moffatt, A.I. Pesci, Boundary singularities produced by the motion of soap films, Proc. National Academy Sciences 111 (2014), 8339-8344.
- [6] J.C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil, Categorical group invariants of 3-manifolds, manuscripta math. 145 (2014), 433-448.
- [7] J.C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil, 2-stratifolds, in “A Mathematical Tribute to José María Montesinos Amilibia”, Universidad Complutense de Madrid, 395-405 (2016).
- [8] J.C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil, Classification of Simply-connected Trivalent 2-dimensional Stratifolds, Top. Proc. (2018)
- [9] J.C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil, 2-stratifold spines of closed 3-manifolds, arXiv:1707.05663 (2017).
- [10] J.C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil, Y.A. Hernández-Esparza, Models of Simply-connected Trivalent 2-dimensional Stratifolds With an Implementation Code, arXiv:1805.06302 (2018)
- [11] M. Khovanov, $sl(3)$ link homology. Algebr. and Geom. Topol. 4, 1045-1081 (2004).
- [12] V. Kurlin, A one-dimensional homologically persistent skeleton of an unstructured point cloud in any metric space, Computer Graphics Forum, 34 (2018). 253 - 262.
- [13] S. Martin, A. Thompson, E.A. Coutsiyas, J.P. Watson, Topology of cyclo-octane energy landscape, Journal Chemical Physics, 132 (2010) 234115.
- [14] S. Matsuzaki and M. Ozawa, Genera and minors of multibranching surfaces, Topology and its Applications 230, 621-638 (2017).

- [15] W. Neumann, A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves, *Trans. Amer. Math. Soc.* 268, 299-344 (1981).