

**PARTIALLY HYPERBOLIC DIFFEOMORPHISMS HOMOTOPIC
TO THE IDENTITY IN DIMENSION 3
PART I: THE DYNAMICALLY COHERENT CASE**

THOMAS BARTHELMÉ, SERGIO R. FENLEY, STEVEN FRANKEL,
AND RAFAEL POTRIE

ABSTRACT. We study 3-dimensional dynamically coherent partially hyperbolic diffeomorphisms that are homotopic to the identity, focusing on the transverse geometry and topology of the center stable and center unstable foliations, and the dynamics within their leaves. We find a structural dichotomy for these foliations, which we use to show that every such diffeomorphism on a hyperbolic or Seifert fibered 3-manifold is leaf conjugate to the time one map of a (topological) Anosov flow. This proves a classification conjecture of Hertz-Hertz-Ures in hyperbolic 3-manifolds and in the homotopy class of the identity of Seifert manifolds.

Nous étudions, en dimension trois, les difféomorphismes partiellement hyperbolique qui sont dynamiquement cohérent et homotope à l'identité. Nous nous focalisons particulièrement sur la géométrie et la topologie de leurs feuilletage central stable et central instable, ainsi que sur leur dynamique dans les feuilles. Ainsi, nous obtenons que la structure de ces feuilletages doit satisfaire à une dichotomie. Usant de cette dichotomie, nous montrons que, lorsque la 3-variété est hyperbolique ou de Seifert, les difféomorphismes étudiés ont leur feuilletage central conjugué à celui du temps un d'un flot d'Anosov topologique. Ceci prouve une conjecture de Hertz-Hertz-Ures pour les variétés hyperbolique et dans la classe d'homotopie de l'identité pour les variétés de Seifert.

Keywords: Partial hyperbolicity, 3-manifold topology, foliations, classification.

MSC 2010: 37D30,57R30,37C15,57M50,37D20.

CONTENTS

1. Introduction	2
1.1. Homotopy, integrability, and conjugacy	3
1.2. Results	3
1.3. Remarks and references	5
1.4. Acknowledgments	5
2. Outline and discussion	5
2.1. Setup	6
2.2. Outline of the paper	7
3. Foliation and good lifts	10
3.1. General homeomorphisms	10
3.2. Consequences for partially hyperbolic systems	15
3.3. Gromov hyperbolicity of leaves	17
3.4. Summary	18
4. Center dynamics in fixed leaves	19
4.1. Perfect fits	19
4.2. Finding fixed center leaves	20

5. Mixed behavior	25
6. Double invariance	25
6.1. Fixing center leaves	26
6.2. Showing that the map is a discretized Anosov flow	29
7. Proof of Theorem A with dynamical coherence	30
8. Coarse dynamics of translations	32
9. Double translations in hyperbolic manifolds	40
Appendix A. Some 3-manifold topology	41
Appendix B. Taut foliations in 3-manifolds	42
Appendix C. Uniformization of leaves	43
Appendix D. Uniform foliations and transverse pseudo-Anosov flows	44
Appendix E. Axes	44
Appendix F. On partial hyperbolicity	46
Appendix G. Discretized Anosov flows	47
Appendix H. The graph transform argument	49
References	51

1. INTRODUCTION

A diffeomorphism f of a 3-manifold M is *partially hyperbolic* if it preserves a splitting of the tangent bundle TM into three 1-dimensional sub-bundles

$$TM = E^s \oplus E^c \oplus E^u,$$

where the *stable bundle* E^s is eventually contracted, the *unstable bundle* E^u is eventually expanded, and the *center bundle* E^c is distorted less than the stable and unstable bundles at each point.

From a dynamical perspective, the interest in partial hyperbolicity stems from its appearance as a generic consequence of certain dynamical conditions, such as stable ergodicity and robust transitivity. For example, a diffeomorphism is *transitive* if it has a dense orbit, and *robustly transitive* if this behavior persists under C^1 -small deformations. Every robustly transitive diffeomorphism on a 3-manifold is either Anosov or “weakly” partially hyperbolic [DPU99]. Analogous results hold for stable ergodicity and in higher dimensions [BDV05].

From a geometric perspective, one can think of partial hyperbolicity as a generalization of the discrete behavior of Anosov flows, which feature prominently in the theory of 3-manifolds. Recall that a flow Φ on a 3-manifold M is *Anosov* if it preserves a splitting of the unit tangent bundle TM into three 1-dimensional sub-bundles

$$TM = E^s \oplus T\Phi \oplus E^u,$$

where $T\Phi$ is the tangent direction to the flow, E^s is eventually exponentially contracted, and E^u is eventually exponentially expanded. After flowing for a fixed time, an Anosov flow generates a partially hyperbolic diffeomorphism of a particularly simple type, where the stable and unstable bundles are contracted uniformly, and the center direction, which corresponds to $T\Phi$, is left undistorted. More generally, one can construct partially hyperbolic diffeomorphisms of the form $f(x) = \Phi_{\tau(x)}(x)$ where Φ is a (topological) Anosov flow and $\tau: M \rightarrow \mathbb{R}_{>0}$ is a positive continuous function; the partially hyperbolic diffeomorphisms obtained in this way are called *discretized Anosov flows*¹.

¹Note that a discretized Anosov flow is not in general the time-1 map of a reparametrization of the Anosov flow, see Appendix G

In this article and its sequel [BFFPa] we show that large classes of partially hyperbolic diffeomorphisms can be identified with discretized Anosov flows. This confirms a large part of the well-known conjecture by Pujals [BW05], which attempts to classify 3-dimensional partially hyperbolic diffeomorphisms by asserting that they are all either discretized Anosov flows or deformations of certain kinds of algebraic examples.

1.1. Homotopy, integrability, and conjugacy. There are two important obstructions to identifying a partially hyperbolic diffeomorphism with a discretized Anosov flow. The first comes from the fact that the latter are homotopic to the identity, while the former may be homotopically nontrivial. Homotopically nontrivial examples include Anosov diffeomorphisms on the 3-torus with distinct eigenvalues, “skew products,” and the counterexamples to Pujals’ conjecture constructed in [BPP16, BGP16, BZ17, BGHP17].

The second major obstruction comes from the integrability of the bundles in a partially hyperbolic splitting. In the context of an Anosov flow Φ , the stable and unstable bundles E^s and E^u integrate uniquely into a pair of 1-dimensional foliations, the *stable* foliation \mathcal{W}^s and *unstable* foliation \mathcal{W}^u . In fact, even the *weak stable* and *weak unstable* bundles $E^s \oplus T\Phi$ and $E^u \oplus T\Phi$ integrate uniquely into a transverse pair of Φ -invariant 2-dimensional foliations, the *weak stable* foliation \mathcal{W}^{ws} and *weak unstable* foliation \mathcal{W}^{wu} .

In the context of a partially hyperbolic diffeomorphism f , the stable and unstable bundles still integrate uniquely into stable and unstable foliations, \mathcal{W}^s and \mathcal{W}^u [HP18]. However, the *center stable* and *center unstable* bundles $E^c \oplus E^s$ and $E^c \oplus E^u$ may fail to be uniquely integrable. In fact, there are examples where it is impossible to find *any* f -invariant 2-dimensional foliation tangent to the center stable or center unstable bundle [RHRHU16, BGHP17].

If one can find a pair of f -invariant foliations tangent to the center stable and center unstable bundles then f is said to be *dynamically coherent*. This condition is certainly satisfied if f is a discretized Anosov flow (cf. Appendix G).

We take dynamical coherence as an assumption in the present article; Part II [BFFPa] works without this.

1.2. Results. Most of the existing progress towards classifying partially hyperbolic diffeomorphisms takes an outside-in approach, restricting attention to particular classes of manifolds, and comparing to an *a priori* known model partially hyperbolic (see [CRRU15, HP18] for recent surveys). In particular, partially hyperbolic diffeomorphisms have been completely classified in manifolds with solvable or virtually solvable fundamental group [HP14, HP15].

Ours is an inside-out approach, using the theory of foliations to understand the way the local structure that defines partial hyperbolicity is pieced together into a global picture. We then relate the dynamics of these foliations to the large-scale structure of the ambient manifold. An advantage of this method is that, since it does not rely on a model partially hyperbolic to compare to, we can consider any manifold, not just one where an Anosov flow is known to exist.

The following two theorems are the main consequences of our work, applied to two of the major classes of 3-manifolds. Note that the classification of partially hyperbolic diffeomorphisms is always considered up to finite lifts and iterates, since one can easily build infinitely many different but uninteresting examples by taking finite covers.

Theorem A (Seifert manifolds). *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a closed Seifert fibered 3-manifold. If f is homotopic to the identity, then some iterate is a discretized Anosov flow.*

We eliminate the assumption of dynamical coherence in [BFFPa]; this resolves the Pujals' Conjecture for Seifert fibered manifolds². Note that Theorem A does not use the classification of Anosov flows on Seifert fibered 3-manifolds [Ghy84, Bar96].

Theorem B (Hyperbolic manifolds). *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a closed hyperbolic 3-manifold. Then some iterate of f is a discretized Anosov flow.*

This resolves a classification conjecture in [CRRU15] for hyperbolic 3-manifolds.

Note that this theorem does not assume that f is homotopic to the identity, since any homeomorphism on a closed hyperbolic 3-manifold has a finite power that is homotopic to the identity. It does, however, assume dynamical coherence.

Theorems A and B strengthen a more general statement which works in every 3-manifold and which requires some knowledge of taut foliations. See Appendix B for the relevant definitions.

Let $f: M \rightarrow M$ be a partially hyperbolic diffeomorphism on a closed 3-manifold M . When f is homotopic to the identity, we denote by \tilde{f} a lift to the universal cover \tilde{M} that is obtained by lifting such a homotopy. For a dynamically coherent partially hyperbolic diffeomorphism, we denote the center stable and center unstable foliations by \mathcal{W}^{cs} and \mathcal{W}^{cu} , and their lifts to \tilde{M} by $\tilde{\mathcal{W}}^{cs}$ and $\tilde{\mathcal{W}}^{cu}$.

Theorem C. *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a closed 3-manifold M that is homotopic to the identity. If \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal, then either*

- (i) f is a discretized Anosov flow, or
- (ii) \mathcal{W}^{cs} and \mathcal{W}^{cu} are \mathbb{R} -covered and uniform, and \tilde{f} acts as a translation on the leaf spaces of $\tilde{\mathcal{W}}^{cs}$ and $\tilde{\mathcal{W}}^{cu}$.

Here, f -minimal means that the only closed sets that are both f -invariant and saturated by the foliation are the empty set and the whole manifold M . If f is transitive or volume preserving, then it is already known that \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal [BW05]. We will show that this holds as well when M is hyperbolic or Seifert and the lift \tilde{f} fixes a leaf in the universal cover (see Proposition 3.15). We show that (ii) cannot occur in a hyperbolic manifold, and Theorem B follows.

It is likely that Theorem B could be shown in the setting of 3-manifolds with atoroidal pieces in their JSJ decomposition, but we have not pursued this here as it would require proving results similar to [Thu, Cal00, Fen02] in this setting.

The technique to eliminate the possibility (ii) in Theorem C is more widely applicable: In a companion article [BFFP20], we use the same ideas to show that a partially hyperbolic diffeomorphism on a Seifert manifold which acts as a pseudo-Anosov on (part of) the base is not dynamically coherent.

For Seifert manifolds, it is possible to show that, after taking an iterate, there is another lift \tilde{f} that is still a bounded distance from the identity and which

²The conjecture is true for Seifert manifolds with fundamental group with polynomial growth [HP14] and false in Seifert fibered manifolds when the isotopy class is not the identity as the examples in [BGP16, BGHP17] are not homotopic to identity and so cannot be discretized Anosov flows.

fixes a leaf of $\widetilde{\mathcal{W}}^{cs}$, and f -minimality follows. We show that (ii) implies leaf conjugacy of (possibly an iterate of) f to a time one map of an Anosov flow on a Seifert fibered manifold, and Theorem A follows. We also completely classify the partially hyperbolic diffeomorphisms for which it is necessary to take an iterate, as opposed to f itself, to get a discretized Anosov flow (see Remark 7.3).

1.3. Remarks and references. The definition of a partially hyperbolic diffeomorphism traces back to [HPS77] and [BP73].

The classification problem for 3-dimensional partially hyperbolic diffeomorphisms has attracted significant attention since the pioneering work of [BW05, BBI04, BI08], which was partially motivated by Pujals' conjecture (see also [PS04, §20] or [HP06, §2.2.6]). For example, there is interesting recent work done under some restrictions on the center dynamics [Zha17, BZ19] or assuming some conditions on the derivatives [CPR19]. See also the surveys [CRRU15, HP18, Pot18].

Besides its intrinsic interest, the classification problem for partially hyperbolic diffeomorphisms has dynamical implications. For example, several finer dynamical and ergodic properties have been studied under the assumption of having a discretized Anosov flow (while not using that terminology), for instance in [AVW15, BFT19] (see also [Pot18, Wil10]). Our results here and in [BFFPa] make that condition checkable. Several of the techniques presented here also yield information about the dynamics along the center direction, which is so far poorly understood (see, e.g. [FP18]).

In addition, this article contains several new results of independent interest. Indeed, important steps in our study do not use partial hyperbolicity, but instead only use the more general setting of foliation preserving diffeomorphisms. Thus some results (see §3 and §8 for instance) are much more widely applicable. In particular, in §8 we use regulating pseudo-Anosov flows to understand the dynamics of a diffeomorphism that translates the leaves of an \mathbb{R} -covered foliation, showing that any such diffeomorphism has “invariant cores” that shadow the closed orbits of the corresponding flow.

1.4. Acknowledgments. We would like to thank Christian Bonatti, Andrey Gogolev, and Andy Hammerlindl for many helpful comments and discussions.

T. Barthelmé was partially supported by the NSERC (Funding reference number RGPIN-2017-04592).

S. Fenley was partially supported by Simons Foundation grant number 280429.

S. Frankel was partially supported by National Science Foundation grant number DMS-1611768. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

R. Potrie was partially supported by CSIC 618 and ANII-FCE-135352.

2. OUTLINE AND DISCUSSION

In this section we will set some basic terminology, outline our major arguments, and detail the organization of this paper.

Definition 2.1. A C^1 -diffeomorphism $f: M \rightarrow M$ on a 3-manifold M is *partially hyperbolic* if there is a Df -invariant splitting of the tangent bundle TM into three 1-dimensional bundles

$$TM = E^s \oplus E^c \oplus E^u$$

such that for some $n > 0$, one has

$$\begin{aligned} \|Df^n|_{E^s(x)}\| &< 1, \\ \|Df^n|_{E^u(x)}\| &> 1, \text{ and} \\ \|Df^n|_{E^s(x)}\| &< \|Df^n|_{E^c(x)}\| < \|Df^n|_{E^u(x)}\|, \end{aligned}$$

for all $x \in M$.

See Appendix F for more details. Our major goal is to show that large classes of partially hyperbolic diffeomorphisms are discretized Anosov flows:

Definition 2.2. A *discretized Anosov flow* is a partially hyperbolic diffeomorphism $g: M \rightarrow M$ on a 3-manifold M that is of the form $g(p) = \Phi_{t(p)}(p)$ for a topological Anosov flow Φ and a map $t: M \rightarrow (0, \infty)$.

The precise definition of a topological Anosov flow is given in Appendix G, where we also explain the relationship between discretized Anosov flows and the more common notion of partially hyperbolic diffeomorphisms that are leaf conjugate to time-1 maps of Anosov flows.

Consider a discretized Anosov flow $g: M \rightarrow M$ on a closed 3-manifold M . We will see (Proposition G.2) that g is dynamically coherent, and that the center leaves of g are exactly the orbits of the underlying flow. This means that g fixes each leaf of the center foliation. Moreover, it has a natural lift $\tilde{g}: \tilde{M} \rightarrow \tilde{M}$ to the universal cover that fixes the lift of each center leaf, but fixes no point in \tilde{M} . Indeed, such a lift may be obtained by flowing points along lifted orbits. That is, $\tilde{g}(p) = \tilde{\Phi}_{t(\pi(p))}(p)$, where $\tilde{\Phi}$ is the lifted flow and $\pi: \tilde{M} \rightarrow M$ is the covering map.

In fact, to show that a partially hyperbolic diffeomorphism $f: M \rightarrow M$ is a discretized Anosov flow, it will suffice to find a lift $\tilde{f}: \tilde{M} \rightarrow \tilde{M}$ with this property, i.e., that fixes the leaves of the lifted center foliation, but fixes no point in \tilde{M} . This argument is essentially given in [BW05, Section 3.5] — see Section 6.2.

2.1. Setup. We will now set some basic definitions and outline our major arguments. We will assume some familiarity with 3-manifold topology, taut foliations, and leaf spaces; see Appendices A and B for an outline of the necessary background.

In this paper, M will be a closed 3-manifold, and $f: M \rightarrow M$ will be a dynamically coherent partially hyperbolic diffeomorphism that is homotopic to the identity.

The center stable, center unstable, stable, unstable, and center foliations on M are denoted by \mathcal{W}^{cs} , \mathcal{W}^{cu} , \mathcal{W}^s , \mathcal{W}^u , and \mathcal{W}^c . These lift by the universal covering map $\pi: \tilde{M} \rightarrow M$ to foliations on \tilde{M} which we denote by $\tilde{\mathcal{W}}^{cs}$, $\tilde{\mathcal{W}}^{cu}$, $\tilde{\mathcal{W}}^s$, $\tilde{\mathcal{W}}^u$, and $\tilde{\mathcal{W}}^c$.

Convention: Throughout this paper we will assume that $\pi_1(M)$ is not virtually solvable.

This assumption implies that there is no closed surface tangent to either E^{cs} or E^{cu} (Theorem F.1), a fact that we will use often.

The classification of partially hyperbolic diffeomorphisms on manifolds with virtually solvable fundamental group is complete [HP14, HP15], and our assumption does not affect our main results (see Theorem F.5).

2.1.1. Good lifts. Since f is homotopic to the identity, we can lift such a homotopy to \tilde{M} , and obtain a lift $\tilde{f}: \tilde{M} \rightarrow \tilde{M}$ that is *good*:

Definition 2.3. A lift $\tilde{f}: \tilde{M} \rightarrow \tilde{M}$ of a homeomorphism $f: M \rightarrow M$ is called a *good lift* if

- (i) it moves each point a uniformly bounded distance (i.e., there exists $K > 0$ such that $d_{\tilde{M}}(x, \tilde{f}(x)) < K$ for all $x \in \tilde{M}$), and
- (ii) it commutes with every deck transformation.

Throughout this paper we will always take \tilde{f} to be a good lift of f .

Remark 2.4. In fact, it is easy to show that (i) follows from (ii) on a closed manifold.

A homeomorphism may have more than one good lift. Indeed, composing a good lift with a deck transformation in the center of the fundamental group yields another good lift. Conversely, the existence of more than one good lift implies that the fundamental group has non-trivial center. By the Seifert fibered space conjecture [CJ94, Gab92], this implies that the manifold is Seifert-fibered with orientable Seifert fibration.

2.2. Outline of the paper. Recall that there are no closed surfaces tangent to the center stable or center unstable bundles. In particular, \mathcal{W}^{cs} and \mathcal{W}^{cu} have no closed leaves, which implies that they are taut. Furthermore, \tilde{M} is homeomorphic to \mathbb{R}^3 , and each leaf of $\tilde{\mathcal{W}}^{cs}$ or $\tilde{\mathcal{W}}^{cu}$ is a properly embedded plane that separates \tilde{M} into two open balls (cf. Theorem B.1).

2.2.1. Dichotomies for foliations. In §3 we study the basic structure of the center stable and center unstable foliations, and the way that \tilde{f} permutes their lifted leaves. Much of this section applies more generally to a homeomorphism that is homotopic to the identity and preserves a foliation.

The basic tool is Lemma 3.3, which says that the complementary components of a lifted center stable or center unstable leaf are “large” in the sense that they contains balls of arbitrary radius. Since \tilde{f} moves points a uniformly bounded distance, this has immediate consequences for the way that it acts on the leaf spaces of $\tilde{\mathcal{W}}^{cs}$ and $\tilde{\mathcal{W}}^{cu}$.

In particular, we deduce that the set of center stable leaves that are fixed by \tilde{f} is closed in the leaf space \mathcal{L}^{cs} of $\tilde{\mathcal{W}}^{cs}$, each complementary component of this set is an open interval that is acted on by \tilde{f} as a translation, and any two leaves in one of these “translation regions” are a finite Hausdorff distance apart (Proposition 3.7). The same holds for the center unstable foliation.

When \mathcal{W}^{cs} is f -minimal, or M is hyperbolic or Seifert-fibered, we use this to show that either:

- (★)
 - \tilde{f} fixes every leaf of $\tilde{\mathcal{W}}^{cs}$, or
 - \mathcal{W}^{cs} is \mathbb{R} -covered and uniform, and \tilde{f} acts as a translation on the leaf space of $\tilde{\mathcal{W}}^{cs}$.

Recall that \mathbb{R} -covered means that the leaf space in the universal cover is homeomorphic to \mathbb{R} , and *uniform* means that any two leaves in the universal cover are a finite Hausdorff distance apart.

This dichotomy is easy to show under the assumption of f -minimality, where it does not use partial hyperbolicity (Corollary 3.10). It takes significantly more work assuming instead that M is hyperbolic or Seifert-fibered (Proposition 3.15).³

If \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal, or M is hyperbolic or Seifert-fibered, we are left with three possibilities:

³This dichotomy holds even without the assumption of dynamical coherence, but the proof is substantially more difficult (see [BFFPa]).

- (1) **double invariance:** \tilde{f} fixes every leaf of both $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$;
- (2) **mixed behavior:** \tilde{f} fixes every leaf of either $\widetilde{\mathcal{W}}^{cs}$ or $\widetilde{\mathcal{W}}^{cu}$, and acts as a translation on the leaf space of the other, which is \mathbb{R} -covered and uniform;
or
- (3) **double translation:** \tilde{f} acts as a translation on both $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$, which are \mathbb{R} -covered and uniform.

The remainder of the argument is arranged around these three possibilities. We will see in §5 that mixed behavior cannot happen. In §6 we show that double invariance implies that f is a discretized Anosov flow. The double translation case is ruled out for Seifert-fibered manifolds in §7, and for hyperbolic manifolds in §8–§9.

2.2.2. *Center dynamics in fixed leaves.* In §4, we work under the assumption that \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^{cs}$, and study the dynamics *within* each center stable leaf. In particular, we show (Proposition 4.4):

- (**) If \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^{cs}$, then any leaf of $\widetilde{\mathcal{W}}^{cs}$ that is fixed by a nontrivial deck transformation contains a center leaf that is fixed by \tilde{f} .

This immediately eliminates the possibility of mixed behavior (see §5). It will also be used in §6 to show that double invariance implies that f is a discretized Anosov flow.

Consider a center stable leaf L that is fixed by a deck transformation γ . The proof of (**) comes down to understanding the topology of the stable foliation within L “in the direction of” γ . The formal meaning of this is the *axis* for the action of \tilde{f} on the stable leaf space in L (see Appendix E), but it can be understood intuitively as the set of all stable leaves that cross the core of the cylinder $\widetilde{M}/\langle\gamma\rangle$ essentially.

Suppose that there is an line’s worth of stable leaves in this direction, which corresponds to circle’s worth in $\widetilde{M}/\langle\gamma\rangle$ as depicted (roughly) in the left half of Figure 1. Then one can find a curve representing γ that is transverse to the stable foliation, and a “graph transform argument” finds a corresponding center leaf preserved by both γ and \tilde{f} (Lemma 4.5).




FIGURE 1. Axes

The other possibility is that one finds gaps, which look roughly like Reeb components as in the right half of Figure 1. We eliminate the possibility of such

gaps by combining the dynamics coming from partial hyperbolicity with two conflicting forces:

- (i) On one hand, the topology of the stable and center foliations within L forces the existence of a center ray within this gap that is expanded by f (Lemma 4.7).
- (ii) On the other hand, we find from the geometry of L that the entire gap, and any center leaf within it, must be coarsely contracted (Lemma 4.9).

These conclusions are contradictory, so there can be no gaps, completing the proof of $(\star\star)$.

The existence of the expanded center ray (i) is delicate, and depends crucially on dynamical coherence (see Remark 4.8 and Figure 6). The coarse contraction of gaps (ii) is more robust, and will be used again in the dynamically incoherent case in [BFFPa].

2.2.3. *Double invariance.* In the doubly invariant case (1), one would like to show that \tilde{f} fixes each center leaf. Since by assumption it fixes each center stable and center unstable leaf, it fixes the intersection between any two such leaves. Each component of this intersection is a collection of center leaves, but there is no a priori reason for it to have a single component. In §6.1, we show that \tilde{f} fixes either every center leaf or no center leaf, and so by $(\star\star)$ it fixes every center leaf.

Once we know that \tilde{f} fixes every center leaf, we can use the arguments of Bonatti–Wilkinson [BW05] to show that the center foliation is the orbit foliation of a topological Anosov flow, and hence that f is a discretized Anosov flow. This is done in §6.2, completing the proof of Theorem C.

2.2.4. *Double translation in Seifert-fibered manifolds.* The double translation case (3) turns out to be the trickiest. The preceding results work with either topological conditions (M being Seifert fibered or hyperbolic) or dynamical conditions (minimality or f -minimality). To handle double translations we will need topological restrictions.

Part of the difficulty is that double translation *do* in fact exist (Remark 7.3)! However, these examples live in Seifert-fibered manifolds, and have iterates that are discretized Anosov flows. Using a similar idea (see section 6.2 of [BFFPb]), one can build homeomorphisms that act as a double translation on any manifold which admits an \mathbb{R} -covered Anosov flow which is not a suspension, but our techniques do show that they cannot be dynamically coherent partially hyperbolic (even in a topological sense) when the ambient manifold is hyperbolic.

Eliminating double translations when M is a Seifert manifold relies on a trick: Since there are many good lifts, we show in §7 that some good lifts (of a power of f) must fix the leaf of at least one foliation. This completes the proof of Theorem A.

2.2.5. *Double translation in hyperbolic manifolds.* We are left to treat the case of double translations in hyperbolic manifolds, which we do in §8 and §9.

In §8, we prove a result about \mathbb{R} -covered foliations that is of general interest. In a hyperbolic 3-manifold, an \mathbb{R} -covered foliation admits a transverse regulating pseudo-Anosov flow (see Appendix D). We will use this flow to understand the dynamics of any homeomorphism that acts as a translation on its leaf space (Proposition 8.1):

Let $f: M \rightarrow M$ be a homeomorphism on a closed hyperbolic 3-manifold that is homotopic to the identity and preserves a taut, \mathbb{R} -covered foliation \mathcal{T} . Suppose that a good lift acts as a translation on $\widetilde{\mathcal{T}}$.

(***). Then for each periodic orbit γ of the regulating pseudo-Anosov flow Φ , there is a corresponding invariant “core” T_γ for f . Moreover, the dynamics of f at T_γ is coarsely identical to the dynamics of Φ at γ (in the sense that they have the same Lefschetz index).

There is a little lie in this description, as the core T_γ is in fact in the cover $\widetilde{M}/\langle\gamma\rangle$ and is invariant under the appropriate lift of f to that cover.

In fact, having a hyperbolic manifold is not essential — we use similar techniques on Seifert-fibered manifolds in [BFFP20].

The result (***) is the main ingredient in §9, where we show that double translations cannot occur in hyperbolic manifolds and complete the proof of Theorem B.

The rough idea is that (***) gives a circle invariant by (a lift of) f and with at least one fixed point, but partial hyperbolicity implies that any fixed point must be, say, repulsive. But the devil is in the details, and while one can make this rough idea precise in the case of a foliation, it does not lead to a contradiction for branching foliation. This is why Theorem B requires dynamical coherence. In the non dynamically coherent case, treated in [BFFPa], while we do not prove or disprove its existence, we obtain a detailed description of the permitted structure in the translation case that in particular produces some topological obstructions for the existence of partially hyperbolic diffeomorphisms in hyperbolic 3-manifolds.

3. FOLIATIONS AND GOOD LIFTS

In this section we will study the way that a good lift \widetilde{f} of a dynamically coherent partially hyperbolic diffeomorphism $f: M \rightarrow M$ that is homotopic to the identity permutes the leaves of the lifted center stable and center unstable foliations.

Most of the arguments in this section apply to any homeomorphism of a 3-manifold that preserves an appropriate foliation and is homotopic to the identity, so we will work for a while in this more general setting. At the end, we obtain the following results for our setting.

Proposition 3.1. *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a closed 3-manifold that is homotopic to the identity, and let \widetilde{f} be a good lift of f . If \mathcal{W}^{cs} is f -minimal, or M is hyperbolic or Seifert fibered, then either*

- (1) \mathcal{W}^{cs} is \mathbb{R} -covered and uniform, and \widetilde{f} acts on the leaf space of the lifted foliations as a translation, or
- (2) \widetilde{f} fixes each leaf of the lifted foliation.

The same holds for the center unstable foliation \mathcal{W}^{cu} .

3.1. General homeomorphisms. Let \mathcal{T} be a taut foliation on a closed 3-manifold M that is not finitely covered by $S^2 \times S^1$. Then the universal cover \widetilde{M} is homeomorphic to \mathbb{R}^3 , and each leaf of the lifted foliation $\widetilde{\mathcal{T}}$ is a properly embedded plane (see Theorem B.1).

Fix a homeomorphism $f: M \rightarrow M$ that preserves \mathcal{T} and is homotopic to the identity, and a good lift $\widetilde{f}: \widetilde{M} \rightarrow \widetilde{M}$ (Definition 2.3).

3.1.1. *Complementary regions.* Being a properly embedded plane, each leaf $K \in \tilde{\mathcal{T}}$ separates \tilde{M} into two open balls. We will call these two components of $\tilde{M} \setminus K$ the *complementary regions* of K . The closure of such a complementary region U is called a *side* of K and is simply $\bar{U} = U \cup K$.

If $K, L \in \tilde{\mathcal{T}}$ are distinct leaves, then $K \cup L$ separates \tilde{M} into three open complementary regions, which can be built from the complementary regions of K and L : Let U, U' be the complementary regions of K , labeled so that $L \subset U'$, and let W, W' be the complementary regions of L , labeled so that $K \subset W'$. Then the complementary regions of $K \cup L$ are $U, V = U' \cap W'$, and W . See Figure 2. We call V the *(open) region between K and L* . Its closure, which is simply $\bar{V} = K \cup V \cup L$, is called the *closed region between K and L* .




FIGURE 2. The region between two leaves

Since \tilde{M} is simply connected, the lifted foliation $\tilde{\mathcal{T}}$ is coorientable. A coorientation determines a labeling of the complementary regions of each leaf $L \in \tilde{\mathcal{T}}$ as a *positive* complementary region denoted L^\oplus and a *negative* complementary region denoted L^\ominus . The corresponding positive and negative sides are denoted by $L^+ = L^\oplus \cup L$ and $L^- = L^\ominus \cup L$.

Remark 3.2. We stress that a priori, some deck transformations or lifts of \mathcal{T} -preserving homeomorphisms may exchange the coorientations of $\tilde{\mathcal{T}}$.

3.1.2. *The big half-space lemma.* The following lemmas will be used to understand the way that \tilde{f} can act on the leaf space of $\tilde{\mathcal{T}}$.

Lemma 3.3. *For every leaf $L \in \tilde{\mathcal{T}}$, and every $R > 0$, there is a ball of radius R contained in each of the complementary regions of L .*

Proof. If necessary, we pass to a double cover of M for which \mathcal{T} is coorientable, and choose such a coorientation. Then every deck transformation preserves the corresponding coorientation on $\tilde{\mathcal{T}}$ and orientation on the leaf space $\mathcal{L} = \mathcal{L}_{\tilde{\mathcal{T}}}$.

Fix a ball $B \subset \tilde{M}$ of arbitrary radius, and a leaf $L \in \tilde{\mathcal{T}}$. We will find a deck transformation g that takes B into L^\oplus ; a similar argument would find a deck transformation that takes B into L^\ominus , completing the proof.

Since B is compact, we can find a leaf $F \in \tilde{M}$ such that $B \subset F^\oplus$. Indeed, the quotient map $\nu : \tilde{M} \rightarrow \mathcal{L}$ takes B to a compact subset $\nu(B)$ of the leaf space, which can be covered by a finite collection of open intervals I_1, I_2, \dots, I_n ; We may assume that $\nu(B)$ intersects every one of these intervals. At least one of these intervals has an initial point (with respect to the orientation on \mathcal{L}) that is not contained in any other interval, and is therefore disjoint from $\nu(B)$. Then B is contained in the positive complementary region of the leaf F corresponding to this initial point.

Let us now find a deck transformation g that takes F^\oplus , and hence $B \subset F^\oplus$, into L^\oplus . Since \mathcal{T} is taut, we can find a positively oriented closed transversal $\gamma: [0, 1] \rightarrow M$, based at a point in $\pi(F)$, that passes through $\pi(L)$. Let $\tilde{\gamma}$ be a lift of γ based at a point in F , which passes through some lift L' of $\pi(L)$. Then we can take $g = h' \circ h$, where h is the deck transformation that takes $\tilde{\gamma}(0)$ to $\tilde{\gamma}(1)$, and h' takes L' to L . The oriented transversal $\tilde{\gamma}$ certifies that $h(F^\oplus) \subset L'^\oplus$, and $h'(L'^\oplus) \subset L^\oplus$ because our deck transformations preserve coorientation. \square

It follows that \tilde{f} can never take a complementary region of a leaf off of itself: This would mean that it takes an arbitrarily large ball off of itself, which contradicts the fact that \tilde{f} moves points a uniformly bounded distance. This has important consequences for the way that \tilde{f} behaves with respect to each leaf.

In particular, if \tilde{f} fixes a leaf, then it cannot interchange its complementary components, and we have:

Corollary 3.4. *If $L \in \tilde{\mathcal{T}}$ is fixed by \tilde{f} , then \tilde{f} preserves coorientations at L .*




FIGURE 3. Translation-like behavior

3.1.3. *Translated leaves.* Similarly, if \tilde{f} moves some leaf, then it does so in a “translation-like” manner as is illustrated in Figure 3. In fact, something a bit stronger is true:

Proposition 3.5. *Let $L \in \tilde{\mathcal{T}}$ be a leaf that is not fixed by \tilde{f} , then*

- (1) *the closed region between L and $\tilde{f}(L)$ is foliated as a product,*
- (2) *\tilde{f} takes each coorientation at L to the corresponding coorientation at $\tilde{f}(L)$,*
and
- (3) *the closed region between L and $\tilde{f}(L)$ is contained in the closed $2R$ -neighborhood of L , where $R = \max_{y \in \tilde{M}} d(y, \tilde{f}(y))$.*

Proof. As in Figure 3, let U, U' be the complementary components of L , labeled so that $\tilde{f}(L) \subset U'$, and let W, W' be the complementary components of $\tilde{f}(L)$, labeled so that $L \subset W'$. Then $V = U' \cap W'$ is the open region between L and $\tilde{f}(L)$.

Note that \tilde{f} must take U to either W or W' . But W is disjoint from U , so we cannot have $\tilde{f}(U) = W$ by Lemma 3.3. Thus \tilde{f} takes U to W' , and U' to W . This is what is meant formally by the aforementioned “translation-like” behavior.

- (1) It follows, in particular, that \tilde{f} takes V off of itself and into W . To see that $\bar{V} = K \cup V \cup L$ is foliated as a product, it suffices to show that every leaf that lies in V separates K from L . Suppose that some leaf $F \subset V$ does not separate K from L . Then K and L are contained in the same complementary region of F , so the other complementary region is contained entirely in the open region V between K and L . But this means that V contains balls of arbitrary radius, which contradicts the fact that \tilde{f} takes V off of itself. Thus every leaf that lies in V separates K from L , and \bar{V} is foliated as a product.
- (2) Since $\bar{V} = K \cup V \cup L$ is foliated as a product, it follows that a coorientation taking $L^\oplus = U'$ will take $f(L)^\oplus = W$. We have already seen that $\tilde{f}(U') = W$, so (2) follows.
- (3) Suppose for a contradiction that there is a point $p \in \bar{V}$ with $d(p, L) = 2R + \epsilon$ for some $\epsilon > 0$. Then since $d(L, \tilde{f}(L)) \leq R$, it follows from the triangle inequality that $d(p, \tilde{f}(L)) \geq R + \epsilon$. This means that the open ball $B_{R+\epsilon}(p)$ at p of radius $R + \epsilon$ is contained in V . But we have already seen that \tilde{f} takes V off of itself, so this implies that $d(p, \tilde{f}(p)) \geq R + \epsilon > R$, a contradiction.

□

It follows that if $L \in \tilde{\mathcal{T}}$ is not fixed by \tilde{f} , then one can string together the \tilde{f} -translates of the closed region \bar{V} between L and $\tilde{f}(L)$ to see that their union

$$U = \dots \cup \tilde{f}^{-2}(\bar{V}) \cup \tilde{f}^{-1}(\bar{V}) \cup \bar{V} \cup \tilde{f}^1(\bar{V}) \cup \tilde{f}^2(\bar{V}) \cup \dots$$

is an open, product-foliated set that is preserved by \tilde{f} . This corresponds to an open interval in the leaf space on which \tilde{f} acts as a translation.

Let $X \subset \tilde{M}$ be the union of all leaves of $\tilde{\mathcal{T}}$ that are fixed by \tilde{f} . Then U is contained in a connected component of $\tilde{M} \setminus X$. In fact, the following lemma says that U is exactly a connected component of $\tilde{M} \setminus X$.

Lemma 3.6. *Let L be a leaf of $\tilde{\mathcal{T}}$ that is not fixed by \tilde{f} , and let $U = \bigcup_{i=-\infty}^{\infty} f^i(\bar{V})$, where \bar{V} is the closed region between L and $\tilde{f}(L)$. Then each leaf in $\partial U = \bar{U} \setminus U$ is fixed by \tilde{f} .*

Proof. The frontier ∂U can be broken into “forwards” and “backwards” frontiers

$$\partial_\omega U = \limsup_{i \rightarrow \infty} f^i(L) \quad \text{and} \quad \partial_\alpha U = \limsup_{i \rightarrow -\infty} f^i(L),$$

each of which is preserved by f .

Let K be a leaf in $\partial_\omega U$, and suppose that $\tilde{f}(K) \neq K$. Then the closed region between K and $\tilde{f}(K)$ would be product foliated, and it follows that either K separates U from $\tilde{f}(K)$ or $\tilde{f}(K)$ separates U from K . This contradicts the fact that $K, \tilde{f}(K) \subset \partial_\omega U$, so we must have $\tilde{f}(K) = K$. A similar argument shows that every leaf in $\partial_\alpha U$ is fixed by \tilde{f} . □

3.1.4. *The dichotomy.* We summarize the preceding discussion in terms of the leaf space:

Proposition 3.7. *Let M be a closed 3-manifold that is not finitely covered by $S^2 \times S^1$, $f: M \rightarrow M$ a homeomorphism homotopic to the identity that preserves a taut foliation \mathcal{T} , and \tilde{f} a good lift.*

The set $\Lambda \subset \mathcal{L}_{\tilde{f}}$ of leaves that are fixed by \tilde{f} is closed and $\pi_1(M)$ -invariant. Moreover, each connected component I of $\mathcal{L}_{\tilde{f}} \setminus \Lambda$ is an open interval that \tilde{f} preserves and acts on as a translation, and every pair of leaves in I are a finite Hausdorff distance apart.

Proof. The only detail that needs to be pointed out is that Λ is $\pi_1(M)$ -invariant, which follows from the fact that \tilde{f} commutes with every deck transformation. \square

In particular, one may have $\Lambda = \emptyset$:

Corollary 3.8. *Let M be a closed 3-manifold that is not finitely covered by $S^2 \times S^1$, $f: M \rightarrow M$ a homeomorphism homotopic to the identity that preserves a taut foliation \mathcal{T} , and \tilde{f} a good lift.*

If \tilde{f} fixes no leaf of $\tilde{\mathcal{T}}$, then \mathcal{T} is \mathbb{R} -covered and uniform, and \tilde{f} acts on $\mathcal{L}_{\tilde{f}} \simeq \mathbb{R}$ as a translation.

This leads to a simple dichotomy when the foliation is f -minimal. Recall:

Definition 3.9. A foliation \mathcal{T} that is preserved by a map $f: M \rightarrow M$ is said to be f -minimal if the only closed sets that are both f -invariant and saturated are M and \emptyset .

Corollary 3.10. *Let M be a closed 3-manifold that is not finitely covered by $S^2 \times S^1$, $f: M \rightarrow M$ a homeomorphism homotopic to the identity that preserves a taut foliation \mathcal{T} , and \tilde{f} a good lift.*

If \mathcal{T} is f -minimal, then either

- (1) *\tilde{f} fixes every leaf of $\tilde{\mathcal{T}}$, or*
- (2) *\mathcal{T} is \mathbb{R} -covered and uniform, and \tilde{f} acts as a translation on the leaf space of $\tilde{\mathcal{T}}$.*

Proof. Since \tilde{f} commutes with each deck transformation, each deck transformation preserves the set $\Lambda \subset \mathcal{L}$ of fixed leaves. In particular, if I is a component of $\mathcal{L} \setminus \Lambda$ and $g \in \pi_1(M)$ then one has either $g(I) = I$ or $g(I) \cap I = \emptyset$.

If $\Lambda = \emptyset$ then we are in case (2) by the preceding corollary.

Suppose that $\Lambda \neq \emptyset$. If $\Lambda \neq \mathcal{L}$, then it corresponds to a closed, \mathcal{T} -saturated subset of M that is preserved by f . Furthermore, this subset is not all of M since it cannot accumulate on a leaf lying in the interior of a complementary interval to Λ . This contradicts f -minimality, so we have $\Lambda = \mathcal{L}$ and are in case (1). \square

3.1.5. *Bounded movement inside leaves.* We end this section by showing that a good lift that fixes every leaf will be within a bounded distance of the identity not only in \tilde{M} but also in each leaf.

Lemma 3.11. *Let M be a closed 3-manifold that is not finitely covered by $S^2 \times S^1$, $f: M \rightarrow M$ a homeomorphism homotopic to the identity that preserves a taut foliation \mathcal{T} , and \tilde{f} a good lift.*

If \tilde{f} fixes every leaf of $\tilde{\mathcal{T}}$, then there is a uniform bound $K > 0$ such that for any leaf $L \in \tilde{\mathcal{T}}$ one has

$$d_L(x, \tilde{f}(x)) < K \text{ for all } x \in L,$$

where d_L is the path metric on L .

Proof. Suppose for a contradiction that there is a sequence of points $x_i \in \widetilde{M}$ for which $d_{L_i}(x_i, \widetilde{f}(x_i))$ tends to infinity, where d_{L_i} is the path metric on the leaf $L_i \in \widetilde{\mathcal{T}}$ containing x_i .

Since M is compact, we can pass to a subsequence and find a sequence of deck transformations γ_i such that $\gamma_i(x_i)$ converges to a point x_∞ . Since \widetilde{f} commutes with γ_i , we have that $\gamma_i(\widetilde{f}(x_i)) = \widetilde{f}(\gamma_i(x_i))$ converges to $\widetilde{f}(x_\infty)$.

Now, since $d_{L_i}(\gamma_i(\widetilde{f}(x_i)), \gamma_i(x_i)) = d_{L_i}(\widetilde{f}(x_i), x_i)$ goes to infinity, the points $\widetilde{f}(x_\infty)$ and x_∞ must be in different leaves of $\widetilde{\mathcal{T}}$. This contradicts the fact that \widetilde{f} fixes each leaf of $\widetilde{\mathcal{T}}$. \square

Remark 3.12. This lemma applies as well to a leaf of a closed sublamination of \mathcal{T} whose lift is leafwise fixed by \widetilde{f} .

3.2. Consequences for partially hyperbolic systems. Let us now specialize, and fix a closed 3-manifold M whose fundamental group is not virtually solvable, a dynamically coherent partially hyperbolic diffeomorphism $f: M \rightarrow M$ that is homotopic to the identity, and a good lift $\widetilde{f}: \widetilde{M} \rightarrow \widetilde{M}$.

We denote by \mathcal{W}^{cs} , \mathcal{W}^{cu} , \mathcal{W}^s , \mathcal{W}^u , and \mathcal{W}^c the center stable, center unstable, stable, unstable, and center foliations.

3.2.1. Fixed points and the topology of leaves.

Lemma 3.13. *Let $L \in \widetilde{\mathcal{W}}^{cs}$ be a leaf that is fixed by \widetilde{f} . If there is a sequence of leaves $L_i \in \widetilde{\mathcal{W}}^{cs}$ that are fixed by \widetilde{f} and accumulate on L , then there are no points in L fixed by non-trivial power of \widetilde{f} .*

Proof. Suppose that \widetilde{f}^n , $n > 0$, fixes some point $x \in L$. Then it fixes the unstable leaf $\widetilde{\mathcal{W}}^u(x)$ through that point. When i is sufficiently large, $\widetilde{\mathcal{W}}^u(x)$ intersects L_i at a single point x_i , which must therefore also be fixed by \widetilde{f}^n . This contradicts the fact that \widetilde{f}^n contracts unstable leaves. \square

Proposition 3.14. *Let $L \in \widetilde{\mathcal{W}}^{cs}$ be a leaf that is fixed by \widetilde{f} . If \widetilde{f} fixes no point in L , then $A = \pi(L)$ has cyclic fundamental group (and is therefore a plane, cylinder, or Möbius band).*

Proof. Let \mathcal{L} be the leaf space of the stable foliation within L .

Since \widetilde{f} fixes no point in L , it cannot fix any stable leaf in L , since a stable leaf that is fixed by \widetilde{f} would contain a fixed point. In other words, \widetilde{f} acts freely on \mathcal{L} , and hence has an axis A_f by Proposition E.2.

Consider two elements $\gamma_1, \gamma_2 \in \pi_1(M)$ that fix L . Since the stable foliation can have no circular leaves, neither of these elements may fix a stable leaf. Hence each γ_i acts freely on \mathcal{L} with an axis A_i .

As \widetilde{f} commutes with both γ_1 and γ_2 it follows from Proposition E.2 that in fact these axes are the same, i.e., $A_1 = A_f = A_2$. Proposition E.2 further implies that the subgroup generated by γ_1 and γ_2 is abelian. Since there are no compact leaves in \mathcal{W}^{cs} , it follows that this subgroup is cyclic, and hence $\gamma_1^n = \gamma_2^m$ for some n, m . \square

3.2.2. Minimality in hyperbolic or Seifert manifolds. The following proposition implies that the dichotomy in Corollary 3.10 holds, without the assumption of f -minimality, when M is hyperbolic or Seifert-fibered.

Proposition 3.15. *Let M be a closed 3-manifold that is hyperbolic or Seifert-fibered, $f: M \rightarrow M$ a dynamically coherent partially hyperbolic diffeomorphism that is homotopic to the identity, and \widetilde{f} a good lift.*

If \tilde{f} fixes one leaf of $\widetilde{\mathcal{W}}^{cs}$, then \mathcal{W}^{cs} is a minimal foliation, and \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^{cs}$.

The same statement holds for \mathcal{W}^{cu} .

Proof. Without loss of generality, we may assume, by passing to a finite cover of M and power of f , that \mathcal{W}^{cs} is orientable and coorientable, and f preserves all orientations and coorientations.

Let $X \subset \widetilde{M}$ be the union of all leaves of $\widetilde{\mathcal{W}}^{cs}$ that are fixed by \tilde{f} . This set is non-empty by hypothesis, $\pi_1(M)$ -invariant as \tilde{f} is a good lift and closed by Proposition 3.7. It follows that $\pi(X) \subset M$ is compact and non-empty. By Zorn's lemma, we can find a minimal compact, non-empty, \mathcal{W}^{cs} -saturated subset $\Lambda \subset \pi(X)$. We will show that $\Lambda = M$, which implies both that \mathcal{W}^{cs} is minimal and that \tilde{f} fixes every leaf.

Note that Λ cannot contain any isolated leaves. Indeed, it cannot consist solely of isolated leaves since then these leaves would be compact, and \mathcal{W}^{cs} has no compact leaves. Deleting an isolated leaf from Λ still leaves a closed, saturated subset, so the existence of an isolated leaf would contradict our minimality assumption.

Let $\widetilde{\Lambda}$ be the preimage of Λ in \widetilde{M} . Since no leaf in Λ is isolated, every leaf in $\widetilde{\Lambda}$ is accumulated on by a sequence of leaves in $\widetilde{\Lambda}$. Since these leaves are all fixed by \tilde{f} , Lemma 3.13 implies that \tilde{f} has no fixed points in $\widetilde{\Lambda}$. It follows from Proposition 3.14 that each leaf of Λ is either a cylinder or a plane.

Assume for a contradiction that $\Lambda \neq M$, and hence $\widetilde{\Lambda} \neq \widetilde{M}$. Then we can choose a nontrivial connected component V of $\widetilde{M} \setminus \widetilde{\Lambda}$.

Claim 3.16. *The projection $\pi(\partial V)$ to M consists of finitely many leaves.*

This is a standard fact in the theory of foliations [CC00, Lemma 5.2.5].

For each $x \in \partial V$, let u_x be the maximal connected unstable segment that starts at x and is contained in \overline{V} , which is either a closed interval or a ray. That is, u_x is the component of $\widetilde{\mathcal{W}}^u(x) \cap \overline{V}$ that contains x . Given $r > 0$, and a leaf $L \subset \partial V$ define

$$A_L^r = \{x \in L \mid \ell(J_x) \geq r\}.$$

Claim 3.17. *For any leaf $L \subset \partial V$, and any $r > 0$, the set $\pi(A_L^r)$ is compact as a subset of $\pi(L)$.*

Proof. It is straightforward to see that $\pi(A_L^r)$ is closed as a subset of $\pi(L)$. If it is not compact, then one can find sequence of points $x_i \in \pi(A_L^r)$ that escapes every compact subset of $\pi(L)$. After taking a subsequence we can assume that the x_i converges in M to some point x . Take a chart around x of the form $D^2 \times (0, 1)$ where each $D^2 \times \{y\}$ is a plaque of \mathcal{W}^{cs} , and each $\{p\} \times (0, 1)$ is an oriented plaque of \mathcal{W}^u . Since the x_i escape every compact subset of $\pi(L)$, we can pass to a subsequence such that each x_i is contained in a different plaque. Then it is easy to see that the lengths of the unstable segments at x_i that stay in $\pi(\overline{V})$ must go to 0, a contradiction. \square

Claim 3.18. *Each leaf $L \subset \partial V$ corresponds to an annulus $\pi(L)$ in M .*

Proof. Fix a leaf $L \subset \partial V$ and an $r > 0$ for which $A_L := A_L^r$ is non-empty. As $\pi(L)$ is either a plane or an annulus, we assume for a contradiction that it is a plane. Then the covering map π restricts to a homeomorphism on L , so the fact that $\pi(A)$ is compact means that A is compact. Let D be a disk in L containing A_L in its interior.

Since the leaves of ∂V are fixed by \tilde{f} , and a positive iterate of \tilde{f} expands the lengths of unstable arcs, we can find an $n \geq 1$ for which $\tilde{f}^n(D) \subset A_L \subset D$.

Then Brouwer's fixed point theorem implies that \tilde{f}^n has a fixed point in L , which contradicts Lemma 3.13. So L must be an annulus. \square

Now we can complete the proof of Proposition 3.15. Let L_1, \dots, L_k be a finite collection of leaves in ∂V that cover $\pi(\partial V)$, and fix $r > 0$ such that each $A_i := A_{L_i}^r$ is nonempty. Choose a compact annulus C_i in each $\pi(L_i)$ that contains $\pi(A_i)$. Since f preserves orientations and coorientations, we can join each C_i to an adjoining C_j with an annulus built out of unstable segments u_x for points $x \in \partial C_i$. Iterating this procedure, we obtain a torus T that consists of alternating annuli contained in leaves of \mathcal{W}^{cs} and annuli transverse to \mathcal{W}^{cs} inside $W = \pi(V)$.

We will now (for the first time) use the assumption that M is hyperbolic or Seifert-fibered to see that T bounds a solid torus.

If M is hyperbolic, then T either bounds a solid torus or is contained in a 3-ball (Lemma A.2). If T is contained in a 3-ball, then the annuli C_i are contained in that ball, so the \mathcal{W}^{cs} leaf containing C_i is compressible. This contradicts the fact that \mathcal{W}^{cs} is a taut foliation (see Theorem B.1), so T bounds a closed solid torus U .

If M is Seifert-fibered, then \mathcal{W}^{cs} is a horizontal foliation. That is, one can isotope \mathcal{W}^{cs} so that all leaves are transverse to the Seifert fibers of M (Theorem F.3). It follows that the complementary regions of the lamination Λ are horizontal. In particular, the region $\pi(V \cup \partial V)$ is a product, which means that the torus T is made up of two horizontal C_i and two transverse annuli, and hence bounds a closed solid torus U .

We will now use a ‘‘volume vs. length’’ argument to get a contradiction. We refer to [HPS18, Proposition 5.2] for a detailed proof and give only a sketch: Consider an unstable arc inside U from a point in some $\pi(A_i)$ to some C_j . Fix some $\epsilon > 0$, and call u the non-empty part of that unstable segment that is at distance $\geq \epsilon$ from both C_i and C_j . Up to taking $\epsilon > 0$ smaller if necessary, we can then assume that u is at distance at least $\epsilon > 0$ from T . Consider a lift $\tilde{u} \subset V$ of u , and note that for any positive n , $\tilde{f}^n(\tilde{u})$ stays a bounded distance away from the corresponding lift \tilde{T} of T . The length of $\tilde{f}^n(\tilde{u})$ will grow exponentially in n , while the volume of its maximal tubular neighborhood can only grow linearly, as \tilde{f} is at bounded distance from the identity and the fundamental group of T is \mathbb{Z} . This means that $\tilde{f}^n(\tilde{u})$ must auto-accumulate in \tilde{M} , contradicting the fact that it is transverse to $\tilde{\mathcal{W}}^{cs}$. Thus $\Lambda = M$ as desired. \square

Remark 3.19. We point out here that the hypothesis of M being hyperbolic or Seifert fibered is used in a single place, but it is crucial. To see this, it is enough to consider the time-one map of Franks-Williams intransitive Anosov flow [FW80] (or any other non-transitive Anosov flow), for which neither the center stable nor the center unstable foliations are minimal.

3.3. Gromov hyperbolicity of leaves. In this section we show that Candel's Theorem (Theorem C.1) applies under the assumptions that f is partially hyperbolic and that \tilde{f} fixes the leaves of the center stable foliation. It is known that the assumption for Candel's Theorem is always satisfied for hyperbolic 3-manifolds (see e.g., [Cal07]), as well as for horizontal foliations in Seifert fibered manifolds with exponential growth of fundamental group (which is automatic in our case thanks to Theorem F.3). However, in order to deal with other 3-manifolds, we need a more general version.

Lemma 3.20. *Let M be a closed 3-manifold, $f: M \rightarrow M$ a dynamically coherent partially hyperbolic diffeomorphism homotopic to the identity, and \tilde{f} a good lift.*

Suppose \mathcal{W}^{cs} has no compact leaves, and \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}^{cs}}$. Then every leaf of \mathcal{W}^{cs} is Gromov-hyperbolic. Moreover, there is a metric on M which restricts to a metric of constant negative curvature on each leaf of \mathcal{W}^{cs} .

Proof. Thanks to Candel's Theorem (Theorem C.1), all we have to show is that \mathcal{W}^{cs} does not admit a holonomy invariant transverse measure.

So we suppose that there is an invariant transverse measure μ to \mathcal{W}^{cs} . Let S be its support. First notice that, as there are no compact leaves in \mathcal{W}^{cs} , μ has no atoms, so there are no isolated leaves in S .

Let $\tilde{\mu}$ be the lift of μ to \widetilde{M} .

The fact that \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}^{cs}}$ implies that the measure μ is f -invariant. To see this, consider τ a small transversal to \mathcal{W}^{cs} , and $\tilde{\tau}$ its lift to \widetilde{M} . Then, since \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}^{cs}}$, the transversals $\tilde{\tau}$ and $\tilde{f}(\tilde{\tau})$ intersect the same set of leaves of \mathcal{W}^{cs} . Hence, $\tilde{\tau}$ and $\tilde{f}(\tilde{\tau})$ have the same $\tilde{\mu}$ -measure (because μ is an invariant transverse measure), thus $\mu(\tau) = \mu(f(\tau))$ as desired.

Now let τ be a closed segment on an unstable leaf and call x one of its endpoints. Note that τ is a transversal to \mathcal{W}^{cs} , and, up to taking a different unstable segment, we assume that $\mu(\tau) > 0$.

We can choose a sequence (n_i) of negative integers converging to $-\infty$ such that $(f^{n_i}(x))$ converges to some $y \in M$.

Then, as the n_i are negative integers, f^{n_i} contracts the unstable length, so the sequence of segments $(f^{n_i}(\tau))$ also converges to y . Now, since μ is f -invariant, it implies that $\mu(f^{n_i}(\tau)) = \mu(\tau) > 0$, for all n_i . By taking the limit, we get that the \mathcal{W}^{cs} leaf containing y must be an atom of μ , in contradiction with the fact recalled earlier that μ has no atoms.

Thus \mathcal{W}^{cs} does not admit an invariant transverse measure and Candel's Theorem yields the conclusion of our lemma. \square

We will use the metric given by this lemma on M in the specific situations where a hyperbolic metric makes the proof less technical. But all such results only need a Gromov-hyperbolic metric in the center stable or center unstable leaves.

3.4. Summary. For convenience, we summarize the results obtained in Section 3.

Corollary 3.21. *Let $f: M \rightarrow M$ be a partially hyperbolic, dynamically coherent, diffeomorphism of a 3-manifold M that is homotopic to the identity. Suppose that \mathcal{W}^{cs} is f -minimal, or that M is hyperbolic or Seifert fibered. Let $\tilde{f}: \widetilde{M} \rightarrow \widetilde{M}$ be any good lift of f .*

Then, \tilde{f} has no fixed points and either

- (1) *the foliation \mathcal{W}^{cs} is \mathbb{R} -covered and uniform, and \tilde{f} acts as a translation on the leaf space of $\widetilde{\mathcal{W}^{cs}}$; or,*
- (2) *the map \tilde{f} leaves every leaf of $\widetilde{\mathcal{W}^{cs}}$ fixed and every leaf of \mathcal{W}^{cs} is a plane, an annulus or a Möbius band. Moreover, there is a metric on M that restricted to each leaf has constant negative curvature -1 .*

Proof. By Proposition 3.7, either the foliation \mathcal{W}^{cs} is \mathbb{R} -covered and uniform, and \tilde{f} acts as a translation on the leaf space of $\widetilde{\mathcal{W}^{cs}}$, or, if \tilde{f} does not act as a translation, then it must fix at least one leaf.

Thus, if M is hyperbolic or Seifert fibered, we can apply Proposition 3.15 and deduce that \mathcal{W}^{cs} is f -minimal.

Hence, if \tilde{f} does not act as a translation, then we can apply Corollary 3.10 to get that \tilde{f} must fix every leaf of $\widetilde{\mathcal{W}}^{cs}$.

Now, if no leaf of $\widetilde{\mathcal{W}}^{cs}$ is fixed by \tilde{f} then \tilde{f} cannot have fixed points. On the other hand, if all leaves of $\widetilde{\mathcal{W}}^{cs}$ are fixed, then we can apply Lemma 3.13 to deduce that \tilde{f} still does not fix points.

Finally, Proposition 3.14 implies that when all leaves of $\widetilde{\mathcal{W}}^{cs}$ are fixed then every leaf is a plane, an annulus, or a Möbius band.

The existence of the claimed metric follows from Lemma 3.20. \square

The same statement holds for the foliation \mathcal{W}^{cu} . Notice that, in principle, the behavior of each foliation is independent. The goal of the next few sections is to show that the behavior of one of the foliations forces the same behavior in the other foliation.

4. CENTER DYNAMICS IN FIXED LEAVES

In this section we will study the dynamics *within* center stable leaves. The main result is Proposition 4.4, which will be used to understand the doubly invariant and mixed cases (see §2.2.1).

4.1. Perfect fits. Much of this section will be concerned with transverse pairs of foliations of a plane — in particular, the stable and center foliations within a center stable leaf. We begin by introducing some basic tools, in particular the idea of “perfect fits” first used by Barbot and the second author [Fen94, Bar95].

Let L be a complete plane equipped with a transverse pair of one-dimensional foliations \mathcal{S} and \mathcal{C} . We denote by $\mathcal{L}^s := L/\mathcal{S}$ and $\mathcal{L}^c := L/\mathcal{C}$ their respective leaf spaces. These are simply-connected, separable 1-manifolds which may not be Hausdorff (see e.g., [Bar98, Cal07, CLN85]).

Definition 4.1. A leaf $c \in \mathcal{C}$ and leaf $s \in \mathcal{S}$ are said to make a *CS-perfect fit*, if they do not intersect, but there is a local transversal τ to \mathcal{C} through c , such that every leaf $c' \in \mathcal{C}$ that intersects τ on one side of c must intersect s .

On the other hand, if there exists τ' a local transversal to \mathcal{S} through s , such that every leaf $s' \in \mathcal{S}$ that intersects τ' on one side of s has to intersect c , we will say that s and c make a *SC-perfect fit*.

If c and s make both a CS-perfect fit and a SC-perfect fit, we say that they make a *perfect fit*.

Lemma 4.2. *If two leaves $c \in \mathcal{C}$ and $s \in \mathcal{S}$ make a CS-perfect fit, then there exists $s' \in \mathcal{S}$, possibly distinct from s such that c and s' make a perfect fit. The symmetric statement holds for SC-perfect fits.*

Proof. Fix a small transversal τ to c . Let c' near enough c which also intersects s . Let $p = c' \cap \tau$ and $q = c' \cap s$. For any x in c' between p and q and near enough p , the stable leaf of x intersects c . Let y in c' between p and q be the first point such that the stable leaf of y does not intersect c . Let s' be this stable leaf. Then c, s' form a perfect fit. \square

A straightforward argument shows the following — see, e.g., [Fen98, Claim in Theorem 3.5].

Lemma 4.3. *If two leaves $s, s' \in \mathcal{S}$ are nonseparated in the leaf space \mathcal{L}^s , then there is a unique leaf $c \in \mathcal{C}$ that separates s from s' and makes a perfect fit with s .*




FIGURE 4. The leaves c and s make a CS -perfect fit, but not a SC -perfect fit. The leaves c and s' make a perfect fit.

4.2. Finding fixed center leaves. The following proposition is the main result of this section.

Proposition 4.4. *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism homotopic to the identity, and \tilde{f} a good lift.*

Suppose that \tilde{f} fixes every leaf of \tilde{W}^{cs} . Then any leaf of \tilde{W}^{cs} that is fixed by a nontrivial element of $\pi_1(M)$ contains a center leaf that is fixed by \tilde{f} .

The proof of this will span the rest of this section. Let us fix M , f , and \tilde{f} as above, along with a leaf $L \in \tilde{W}^{cs}$ and a nontrivial element $\gamma \in \pi_1(M)$ that fixes L . Our goal is to find a center leaf $c \subset L$ that is fixed by \tilde{f} .

Let \mathcal{L}^s and \mathcal{L}^c be the leaf spaces of the foliations \tilde{W}^s and \tilde{W}^c restricted to L . These are simply connected, separable, 1-manifolds that may not be Hausdorff.

Since \tilde{f} fixes every center stable leaf, Lemma 3.13 implies that \tilde{f} has no fixed points. This means that \tilde{f} cannot fix any stable leaf, since such a leaf would be contracted and hence contain a fixed point, so \tilde{f} acts freely on \mathcal{L}^s .

Since there are no circular stable leaves downstairs, γ must also act freely on \mathcal{L}^s . By Proposition 3.14, the stabilizer of L is cyclic, so we may take γ to be a generator.

Since \tilde{f} and γ commute and act freely on \mathcal{L}^s , they preserve an axis $A^s \subset \mathcal{L}^s$, which is either a line or a \mathbb{Z} -union of intervals (see Proposition E.2 and Remark E.3).

The following lemma completes the proof of Proposition 4.4 when A^s is a line.

Lemma 4.5. *If $A^s \simeq \mathbb{R}$, then there exists a center leaf $c \subset L$ that is fixed by both \tilde{f} and γ .*

Proof. We will use the graph transform argument (Lemma H.1).

Since A^s is homeomorphic to \mathbb{R} , one can find a bi-infinite curve η in L that is transverse to the stable foliation and invariant under γ . For instance, pick a point x in L and an arc a from x to γx transverse to the stable foliation. Concatenating the positive and negative iterates of a by γ gives such a curve η (that can be smoothed if required).

In particular, η represents the axis A^s of γ , in the sense that a stable leaf is in A^s if and only if it intersects η .

Since A^s is also the axis for the action of \tilde{f} on \mathcal{L}^s , every \tilde{f} -iterate of η also represents A^s . In particular, $\tilde{f}(\eta)$ and η intersect the exact same set of stable leaves.

So the curve η satisfies the two hypothesis of Lemma H.1, and we obtain a curve β in L that is tangent to E^c and invariant under both \tilde{f} and γ . It remains to show that β is in fact a center leaf.

Choose a point $x \in \beta$, and let β' be the compact subsegment of β running from x to γx . This is a fundamental domain for the action of γ on β .

At each point $y \in \beta$, one can find a compact center segment c_y through y that intersects the same set of stable leaves as some compact subsegment β_y of β , where the interior of β_y contains y . By compactness of β' , one can find a finite collection c_1, c_2, \dots, c_k of these center segments such that the corresponding subsegments of β covers β' .

Projecting to M , we have a finite union of center segments $\bigcup \pi(c_i)$ that intersects the same set of stable leaves as the closed curve $\pi(\beta)$. Since f contracts stable segments, $f^n(\bigcup \pi(c_i))$ converges to $\pi(\beta)$ as $n \rightarrow \infty$. Since a sequence of center segments can only converge to a center segment, it follows that $\pi(\beta)$ is a center leaf, and so is β . \square

4.2.1. *Gaps.* To complete the proof of Proposition 4.4 we will show that A^s is indeed a line.

Suppose that A^s is not a line. Then it is a \mathbb{Z} -union of closed intervals

$$A^s = \bigcup_{i \in \mathbb{Z}} [s_i^-, s_i^+].$$

We will call each of the pairs s_i^+, s_{i+1}^- a *gap* in this axis.

The following lemma says that some positive power of f fixes the image of each gap in M .

Lemma 4.6. *There are $m \neq 0$ and $n > 0$ such that $h = \gamma^m \circ \tilde{f}^n$ fixes every s_i^\pm .*

Proof. Since \tilde{f} and γ act freely on A^s , they act freely on the index set of the collection of intervals, which is \mathbb{Z} . It follows that some nontrivial element of the group generated by \tilde{f} and γ acts trivially on \mathbb{Z} . This element is of the form $h := \gamma^m \circ \tilde{f}^n$. Since both γ , and \tilde{f} act freely on A^s , neither n nor m can be equal to zero, and we can take $n > 0$. Since h fixes each interval, it fixes the endpoints of each interval as desired. \square

For the remainder of this section, we fix h as in this lemma, and look at a single gap, setting $s^+ = s_i^+$ and $s^- = s_{i+1}^-$ for some fixed i .

Let us name some features of this gap — refer to Figure 5.

Proposition E.2 says that s^+ is non-separated from s^- in the leaf space \mathcal{L}^s , so Lemma 4.3 provides a center leaf c that makes a perfect fit with s^+ and separates s^+ from s^- . Since there is a unique such leaf, it follows that h fixes c .

Note that h eventually contracts stable leaves; this is because γ is an isometry, \tilde{f} eventually contracts stable leaves, and $n > 0$. Up to an iterate we can assume that this contraction is immediate. It follows that h fixes a single point x within s^+ . Let c' be the center ray that starts at x on the side of c .

We will show that h expands c' in Lemma 4.7, and that it contracts c' in Corollary 4.13. This is contradictory, so there are no gaps in A^s , i.e., it is a line, and the proof of Proposition 4.4 is complete.




FIGURE 5. A perfect fit forces expansion on a center ray.


4.2.2. *Perfect fits and expanded center rays.* In the following lemma we find that the topology of the stable and center foliations in L forces a stable ray in our gap to expand.

Lemma 4.7. *h acts as an expansion on c' with unique fixed point x .*

Proof. Note that the stable leaf $s_y = \widetilde{W}^s(y)$ through any point $y \in c'$ that is sufficiently close to x will intersect c . This is because s^+ and c make a perfect fit, and c' is a transversal to \mathcal{S} on the side of c . Given such a point, let s'_y be the compact segment of s_y that runs from c' to c . Since the lengths of h -iterates of this segment go to zero, i.e., $\lim_{n \rightarrow \infty} \ell(h^n(s'_y)) = 0$, it follows that the h -iterates of y eventually escape every compact set. Indeed, otherwise one would find that c and c' intersect at some accumulation point of $h^n(y)$. See Figure 5.

The lemma follows since we can take $y \in c'$ arbitrarily close to x . \square

Remark 4.8. The proof of Lemma 4.7 uses the structure of the transverse pair of foliations in an essential way. It does not hold when the center leaves are allowed to merge — see Figure 6. This is exactly the type of behavior that arises in the examples of [RHRHU16].

FIGURE 6. The center ray may land by merging in c in the non-dynamically coherent case.

4.2.3. *Coarse contraction in stable gaps.* In the following lemma we find that the geometry of the gap forces it to contract laterally. This contradicts the expansion found in Lemma 4.7 — see Corollary 4.13.

Lemma 4.9. *There is a rectangle R bounded by segments of s^+ and s^- that contain the fixed points, together with two arcs τ_1, τ_2 , such that $h(R)$ is contained in the interior of R .*

See Figure 7.




FIGURE 7. The domain R is mapped onto itself by h .

We will need two lemmas. The first is that the gap is “uniformly thin”:

Lemma 4.10. *The leaves s^+ and s^- are a bounded Hausdorff distance apart with respect to the path metric on L .*

Proof. Since this gap is part of the axis $A^s = \bigcup_{k \in \mathbb{Z}} [s_k^-, s_k^+]$, it follows that s^- separates s^+ from either $\tilde{f}(s^+)$ or $\tilde{f}^{-1}(s^+)$. Then Lemma 3.11 implies that the Hausdorff distance between s^+ and $\tilde{f}^{\pm 1}(s^+)$ is uniformly bounded above, and the same holds for the Hausdorff distance between s^+ and s^- . \square

Recall that, since \tilde{f} fixes all leaves of \tilde{W}^{cs} , Candel’s theorem (Theorem C.1) implies that there is a metric g on M such that \mathcal{W}^{cs} is leafwise hyperbolic. Let d be the associated path metric on the leaf L .

Lemma 4.11. *For any $K_0 > 0$, and any ray $r \subset s^+$, there exists $y \in r$ such that $d(y, h(y)) > K_0$.*

Proof. Let r be a ray of s^+ . Suppose for a contradiction that there exists K_0 such that for all y in r one has $d(y, h(y)) < K_0$.

Recall that $h = \gamma^m \tilde{f}^n$, where m and n are fixed. By Lemma 3.11, there exists a constant K_1 such that, for any z in L , $d(z, \tilde{f}^n(z)) < K_1$.

Thus, by assumption, for any $y \in r$,

$$d(y, \gamma^m y) \leq d(y, \tilde{f}^n(\gamma^m y)) + d(\gamma^m y, \tilde{f}^n(\gamma^m y)) < K_0 + K_1.$$

Now γ is an hyperbolic isometry for d (since γ acts without fixed points on L , and γ is not parabolic because M is compact). Hence, since $d(y, \gamma^m y)$ stays bounded for all y in r , it implies that r has to stay a bounded distance from the (or any when d is only supposed Gromov-hyperbolic) geodesic in L that is the axis for the action of γ on L .

So $\pi(r)$ stays a bounded distance away from the (or all) geodesic in $A = \pi(L)$ that lifts to the axis of γ . Thus, Poincaré–Bendixon Theorem implies that $\pi(r)$ must accumulate onto a closed stable leaf in M , which is impossible (see Figure 8). \square

Remark 4.12. Notice that this is the only place in the proof of Proposition 4.4 that Theorem C.1 is used.

In fact, the proof does not actually need d to come from a Riemannian hyperbolic metric — only that it is Gromov-hyperbolic — so Lemma 4.11 will hold as long as we know Gromov-hyperbolicity of the leafwise metric. We will need this in [BFFPa].




FIGURE 8. If a stable ray in L stays close to the axis of the deck transformation γ which is a hyperbolic isometry, then its projection in M has to accumulate on a circle stable leaf.

Proof of Lemma 4.9. Let y_1 and y_2 be points in s^+ that lie on either side of, and far away from, the fixed point x^+ , and let τ_1 and τ_2 be geodesic arcs from y_1 and y_2 to s^- . See Figure 7. Note that the lengths of τ_i , $i = 1, 2$, are uniformly bounded by Lemma 4.10, and since f has bounded derivatives, the length of $h(\tau_i)$ is bounded as well. By Lemma 4.11, we can ensure that h moves y_i far enough to ensure that $h(\tau_i)$ is disjoint from τ_i , and the lemma follows. \square

Corollary 4.13. *Some subsegment of c' is contracted by h .*

Proof. This follows from Lemma 4.9, noting that c' must intersect either τ_1 or τ_2 . \square

This completes the proof of Proposition 4.4.

Remark 4.14. Note that Lemma 4.9 also implies that the center leaf c that separates s^+ from s^- is “coarsely contracted” by h , in the sense that certain sufficiently large subsegments of c are taken properly into themselves. To show this one needs to note that the set of fixed points of h in c is compact and this follows from the same argument as in Lemma 4.11.

This then generalizes as follows:

Lemma 4.15. *Let c be a center leaf in a center stable leaf $L \subset \widetilde{M}$. Suppose that L is Gromov-hyperbolic, and fixed by \tilde{f} and some nontrivial $\gamma \in \pi_1(M)$. Moreover, assume that there exist two stable leaves s_1, s_2 on L such that:*

- (1) *The center leaf c is in the region between s_1 and s_2 ;*

- (2) The leaves s_1 and s_2 are a bounded Hausdorff distance apart;
 (3) The leaves c , s_1 and s_2 are all fixed by $h = \gamma^n \circ \tilde{f}^m$, $m \neq 0$.

Then, there exists a compact segment $I \subset c$, such that h (if $m > 0$) or h^{-1} (if $m < 0$) acts as a contraction on $c \setminus \dot{I}$.

This remains true without assuming dynamical coherence — we will use it in [BFFPa]. The proof of this lemma is very similar to that of Lemma 4.9. Note that we do not need c to make a perfect fit with s_1 or s_2 , nor do we need that c necessarily goes to both ends of the band determined by s_1 and s_2 as in Figure 7. All we need is that c is between s_1 and s_2 , and that both rays of c escape every compact set in L . That last fact is true of any center leaf in \widetilde{M} .

5. MIXED BEHAVIOR

We can now eliminate mixed behavior in our cases of interest.

Theorem 5.1. *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism homotopic to the identity. Assume that \mathcal{W}^{cu} is f -minimal or that M is hyperbolic or Seifert.*

If a good lift \tilde{f} fixes all the leaves of $\widetilde{\mathcal{W}}^{cs}$, then it also fixes all the leaves of $\widetilde{\mathcal{W}}^{cu}$.

Proof. Since M is not \mathbb{T}^3 (recall that $\pi_1(M)$ is not virtually solvable), Proposition B.2 says that we can find a leaf in \mathcal{W}^{cs} with nontrivial fundamental group. Let $L \in \widetilde{\mathcal{W}}^{cs}$ be a lift of such a leaf, which is invariant by some nontrivial $\gamma \in \pi_1(M)$.

By Proposition 4.4, \tilde{f} fixes some center leaf c in L , so it must fix the center unstable leaf $K \in \widetilde{\mathcal{W}}^{cu}$ that contains L . From the dichotomy in Corollary 3.21, it follows that \tilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^{cu}$ as desired. \square

In particular, under the assumptions of this theorem, one rules out the mixed case (see item (2) of section 2.2.1).

6. DOUBLE INVARIANCE

In this section we show that, under the appropriate conditions, the *doubly invariant case* (see item (1) of section 2.2.1) leads to a discretized Anosov flow.

Theorem 6.1. *Let f be a dynamically coherent partially hyperbolic diffeomorphism. Assume that M is hyperbolic or Seifert or that \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal. If there exists a good lift \tilde{f} which fixes a leaf of $\widetilde{\mathcal{W}}^{cs}$, then f is a discretized Anosov flow.*

Thanks to Proposition 3.15, under any of the hypothesis of the Theorem, both \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal. Notice also that Theorem 5.1 implies that \tilde{f} must fix every leaf of both $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$.

Notice that Theorem 6.1 together with the dichotomy of Corollary 3.10 proves Theorem C from the introduction.

We will first show that connected components of the intersections of center stable and center unstable leaves are fixed by \tilde{f} (i.e., that \tilde{f} fixes leaves of the center foliation). Proving that f is a discretized Anosov flow will then follow rather easily.

We will prove that the set of connected components of intersections fixed by \tilde{f} is both open and closed and then that it is non-empty, thus proving that all center leaves are fixed.

6.1. Fixing center leaves. The main step in the proof of Theorem 6.1 is the following proposition. Recall that the lift $\widetilde{\mathcal{W}}^c$ of the center foliation \mathcal{W}^c consists of the connected components of the intersections between leaves of $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$.

Proposition 6.2. *Let f be a dynamically coherent partially hyperbolic diffeomorphism homotopic to the identity. Let \widetilde{f} be a good lift of f which fixes every leaf of $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$. Suppose that \mathcal{W}^{cs} and \mathcal{W}^{cu} are f -minimal in M . Then \widetilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^c$.*

The key point in the proof of this proposition is to show that either all leaves of $\widetilde{\mathcal{W}}^c$ are fixed by \widetilde{f} , or no leaf of $\widetilde{\mathcal{W}}^c$ is fixed by \widetilde{f} . In the latter case we will use an argument similar to that of the analysis of the mixed behavior case, reaching a contradiction from the results of section 4.

Lemma 6.3. *The set*

$$\text{Fix}_{\widetilde{f}}^c := \left\{ x \in \widetilde{M} \mid \text{the center leaf through } x \text{ is fixed by } \widetilde{f} \right\}$$

is open in \widetilde{M} . In addition $\text{Fix}_{\widetilde{f}}^c$ is invariant under deck transformations.

Proof. Let $c \in \widetilde{\mathcal{W}}^c$ be such that $\widetilde{f}(c) = c$. Let $L = \widetilde{\mathcal{W}}^{cs}(c)$ be the center-stable leaf containing c .

Let $\epsilon > 0$ be small enough so that the center and stable foliations restricted to any ball of radius ϵ in L is product (i.e., every stable and central leaf in the ball intersect each other).

Let $x \in c$. By continuity of f , pick $\delta > 0$ such that if $d(x, y) < \delta$ then $d(\widetilde{f}(x), \widetilde{f}(y)) < \epsilon$. Up to taking δ smaller, and since $\widetilde{f}(x) \in c$, we can assume that for any $y \in L$ such that $d(x, y) < \delta$, we have that $c(y)$, the central leaf through y , intersects $s(\widetilde{f}(x))$, the stable leaf through $\widetilde{f}(x)$. This δ a priori depends on x .

Let $y \in L$ such that $d(x, y) < \delta$, then $c(y) \cap s(\widetilde{f}(x)) \neq \emptyset$. Moreover, since $d(\widetilde{f}(x), \widetilde{f}(y)) < \epsilon$, we have that $c(\widetilde{f}(y)) \cap s(\widetilde{f}(x)) \neq \emptyset$. So the stable leaf $s(\widetilde{f}(x))$ intersects both $c(y)$ and $c(\widetilde{f}(y))$.

Now, as \widetilde{f} fixes the leaves of the central unstable foliations, we have that $\widetilde{\mathcal{W}}^{cu}(c(y)) = \widetilde{\mathcal{W}}^{cu}(c(\widetilde{f}(y)))$. But $s(\widetilde{f}(x))$ is transverse to $\widetilde{\mathcal{W}}^{cu}$, so it cannot intersect the same leaf of $\widetilde{\mathcal{W}}^{cu}$ more than once (see Theorem B.1). Hence $c(\widetilde{f}(y)) = c(y)$.

Thus, the set of center leaves fixed by \widetilde{f} in a center stable leaf is open in that center stable leaf. As the same argument applies to center unstable leaves, we obtain that the union of points in center leaves in $\text{Fix}_{\widetilde{f}}^c$ is open in \widetilde{M} .

Finally, since \widetilde{f} commutes with every deck transformation, $\text{Fix}_{\widetilde{f}}^c$ is $\pi_1(M)$ -invariant. \square

We will think of $\text{Fix}_{\widetilde{f}}^c$ as both a subset of \widetilde{M} and a collection of center leaves in $\widetilde{\mathcal{W}}^c$.

Let $D := \pi(\text{Fix}_{\widetilde{f}}^c)$. By Lemma 6.3, D is open in M , and, obviously, f -invariant.

Lemma 6.4. *Either $\text{Fix}_{\widetilde{f}}^c = \widetilde{M}$ or $\text{Fix}_{\widetilde{f}}^c = \emptyset$.*

Proof. Assume that $\text{Fix}_{\widetilde{f}}^c \neq \emptyset$, and thus, $D \neq \emptyset$.

We start by showing that every leaf of $\widetilde{\mathcal{W}}^{cs}$ has at least some fixed center leaves: Let E be the $\widetilde{\mathcal{W}}^{cs}$ -saturation of Fix_f^c , and suppose for a contradiction that there exists L a leaf of $\widetilde{\mathcal{W}}^{cs}$ such that $L \cap E = \emptyset$.

Since Fix_f^c is $\pi_1(M)$ -invariant, so is E . Hence, for any $\gamma \in \pi_1(M)$, we have $\gamma L \cap \text{Fix}_f^c = \emptyset$. Therefore, in M , we have

$$\pi(L) \cap \pi(E) = \emptyset.$$

So $\pi(L)$ is contained in the set $M \setminus \pi(E)$, which is thus non empty. But $\pi(E)$ is the \mathcal{W}^{cs} -saturation of D , hence open since D is open. The set $\pi(E)$ is also f -invariant since D is. Therefore, $M \setminus \pi(E)$ is a non empty, closed, f -invariant subset of M saturated by \mathcal{W}^{cs} . The f -minimality of \mathcal{W}^{cs} implies that $M \setminus \pi(E) = M$, which is in contradiction with the fact that $\pi(E)$ is non empty.

It follows that, for any center-stable leaf L , we have $L \cap \text{Fix}_f^c \neq \emptyset$.

Our next step is to prove that any center stable leaf that has a non-trivial stabilizer in $\pi_1(M)$ is contained in Fix_f^c . Let L be a leaf of $\widetilde{\mathcal{W}}^{cs}$ such that its projection $A = \pi(L)$ is not simply connected (in which case it must be an annulus or a Möbius band according to Corollary 3.21). As we proved above, we know that $L \cap \text{Fix}_f^c \neq \emptyset$. We now want to show that $L \subset \text{Fix}_f^c$. Let us assume for a contradiction that $\text{Fix}_f^c \cap L \neq L$.

Recall that Fix_f^c is open (by Lemma 6.3), thus so is $B = \text{Fix}_f^c \cap L$ (for the relative topology on L). Notice that, since both Fix_f^c and L are invariant by \widetilde{f} , so is B , and in turn, so is its boundary ∂B .

Let c_1 be a center leaf in ∂B . Then $\widetilde{f}(c_1) \neq c_1$, but arbitrarily near c_1 there are some fixed center leaves.

Since c_1 and $\widetilde{f}(c_1)$ are both in ∂B , they are non separated from each other in the leaf space of the center foliation in L . Indeed, if one takes a sequence (c_n) of central leaves in B that accumulates on c_1 , then, since $\widetilde{f}(c_n) = c_n$, the sequence also accumulates on $\widetilde{f}(c_1)$.

As c_1 and $\widetilde{f}(c_1)$ are not separated in the center leaf space of L , it follows that there exists a stable leaf s_1 making a perfect fit with c_1 , such that s_1 separates c_1 from $\widetilde{f}(c_1)$.

If some power of \widetilde{f} fixes s_1 , then that power of \widetilde{f} has a fixed point in s_1 , contradicting Lemma 3.13 (since \widetilde{f} fixes every leaves of $\widetilde{\mathcal{W}}^{cs}$).

It follows that the sequence $(\widetilde{f}^i(s_1))$ is infinite. Moreover, there exists $c \in \text{Fix}_f^c$ that intersects all the leaves $(\widetilde{f}^i(s_1))$. Indeed, taking $c \in \text{Fix}_f^c$ to be a central leaf close enough to c_1 so that $c \cap s_1 \neq \emptyset$, then c intersects every $\widetilde{f}^i(s_1)$, because $\widetilde{f}(c) = c$.

Furthermore, for all i , $\widetilde{f}^i(s_1)$ separates $\widetilde{f}^{i-1}(s_1)$ from $\widetilde{f}^{i+1}(s_1)$, because \widetilde{f} acts as a translation on c (because \widetilde{f} cannot have a fixed point in L by Lemma 3.13).

As \widetilde{f} acts freely on the stable leaf space in L (again thanks to Lemma 3.13), then \widetilde{f} has an axis $A^s(\widetilde{f})$ for this action. By definition, all the leaves $\widetilde{f}^i(s_1)$ are in this axis. Since all the leaves $\widetilde{f}^i(s_1)$ also intersect a common transversal c , we deduce that $A^s(\widetilde{f})$ is a line (see Figure 9).

Now recall that $C = \pi(L)$ is an annulus or a Möbius band, (thanks to Corollary 3.21). Let γ be the deck transformation associated with the generator of $\pi_1(C)$, so that γ fixes L .




FIGURE 9. The combination of fixed and non-fixed center leaves allows to construct a center leaf intersecting s_1 and $\tilde{f}(s_1)$ in the axis $A^s(\tilde{f}) = A^s(\gamma)$.

Recall that, since there does not exist closed stable leaves in M , γ must act freely on the stable leaf space in L . Thus γ admits an axis $A^s(\gamma)$. Since \tilde{f} and γ commute, then $A^s(\tilde{f}) = A^s(\gamma)$ (see Proposition E.2). In particular, $A^s(\gamma)$ is a line.

Therefore there exists a γ -invariant curve in L , that we call α , such that α is transverse to the stable foliation, and intersects each stable leaf in $A^s(\gamma) = A^s(\tilde{f})$ exactly once. It follows that $\tilde{f}(\alpha)$ and α intersect exactly the same set of stable leaves in L . So we can use the Graph Transform argument (Lemma H.1) on α and obtain that there exists a curve c_0 in L , tangent to the central direction⁴ E^c , and invariant by both \tilde{f} and γ .

Since c_0 intersects s_1 , and the leaves s_1 and c_1 make a perfect fit, we deduce that there exists s , close to s_1 , that intersects both c_0 and c_1 . Let $x = c_0 \cap s$, $y = c_1 \cap s$ and $z = c_0 \cap s_1$. Up to choosing s closer to s_1 , we may assume that the distance between x and z is less than some fixed $K > 0$, the length of the closed curve $\pi(c_0)$. Now, since c_0 is invariant by γ , we have that, for all n ,

$$d(\tilde{f}^n(x), \tilde{f}^n(z)) \leq K.$$

Moreover, since \tilde{f} contracts stable length, we have that $d(\tilde{f}^n(x), \tilde{f}^n(y))$ converges to 0 as n goes to $+\infty$.

Using the above, together with the invariance of c_0 by \tilde{f} and the fact that c_0 is tangent to the central direction, we deduce that for some large enough n , the leaf $\tilde{f}^n(c_1)$ intersects $\tilde{f}^n(s_1)$, contradicting the fact that s_1 and c_1 do not intersect.

Hence, we proved thus far that for any L a center-stable leaf with non-trivial stabilizer, we have $L \subset \text{Fix}_{\tilde{f}}^c$. We can now finish the proof of Lemma 6.4.

Let c be any center leaf in \tilde{M} . Let x be a point in c , let U be the center unstable leaf containing c , and let τ be a small unstable segment in $\tilde{W}^u(x)$ that contains x in its interior.

⁴In fact, since there exists a central leaf that is transverse to the axis, an argument used in the proof of Lemma 4.5 shows that c_0 is not just tangent to the central direction, but an actual central leaf. However, just having tangency to the central direction is enough to finish the proof.




FIGURE 10. The fixed center circle and the circles in the boundary of $\pi(B)$ are joined by a stable leaf.

Recall (see Proposition B.2) that there exists leaves in $\widetilde{\mathcal{W}}^{cs}$ with non-trivial stabilizer. Then f -minimality implies that such leaves are dense. Thus, we may assume that both endpoints of τ are on center stable leaves with non-trivial stabilizer. Call c_1 and c_2 the center leaves through the two endpoints of τ . We proved above that both c_1 and c_2 are fixed by \widetilde{f} (since they are on center stable leaves with non-trivial stabilizer).

Since c intersects τ , an unstable segment from c_1 to c_2 , it follows that c separates c_1 from c_2 in U . As \widetilde{f} fixes both c_1 and c_2 then $\widetilde{f}(c)$ also separates c_1 from c_2 in U . This implies that $\widetilde{f}(c)$ also intersects τ . As argued before, since \widetilde{f} fixes every center stable leaves, c and $\widetilde{f}(c)$ must be in the same center stable leaf, and, since they both intersect τ , which is a transversal to the center stable foliation, we deduce that $c = \widetilde{f}(c)$.

Therefore, we proved that \widetilde{f} fixes every center leaf, i.e., $\text{Fix}_{\widetilde{f}}^c = \widetilde{M}$, as desired. \square

We can now prove Proposition 6.2.

Proof of Proposition 6.2. By assumption, \widetilde{f} fixes every leaf of $\widetilde{\mathcal{W}}^{cs}$, and, by Proposition B.2, there exists some center stable leaf with non-trivial stabilizer. Thus, Proposition 4.4 implies that there exists at least one fixed center leaf, i.e., $\text{Fix}_{\widetilde{f}}^c \neq \emptyset$. Lemma 6.4, then yields that $\text{Fix}_{\widetilde{f}}^c = \widetilde{M}$, which is what we wanted to prove. \square

6.2. Showing that the map is a discretized Anosov flow.

Proposition 6.5 (Leaf conjugacy to a topological Anosov flow). *Let f be a partially hyperbolic diffeomorphism on a 3-manifold M . Suppose that there exists a lift \widetilde{f} to the universal cover \widetilde{M} such that \widetilde{f} moves points a bounded distance and \widetilde{f} fixes every center leaf. Then the center foliation is the orbit foliation of a topological Anosov flow.*

The proof is very similar to that given in [BW05, Section 3.5]. We sketch the main points of the proof. We also refer to Appendix G for the precise definition of a topological Anosov flow, and more discussion about discretized Anosov flows.

Proof. Fix a metric on M and consider X^c a unit vector field in E^c which we first assume orientable. In the universal cover, using that \widetilde{f} fixes every center leaf, one can show that \widetilde{f} does not fix any point in \widetilde{M} , that there is a uniform estimate

for $d_c(x, \tilde{f}(x))$, and it is indeed continuous (see [BW05, Lemma 3.4] for a proof with less hypothesis). In particular, we can assume that $[x, \tilde{f}(x)]^c$ is positively oriented with respect to X^c .

Now, let c_1, c_2 be two center leaves in the same center stable leaf such that for some $x \in c_1$ one has that $\tilde{\mathcal{W}}^s(x) \cap c_2 \neq \emptyset$. Then, letting y be the point of intersection, we have that $d(\tilde{f}^n(x), \tilde{f}^n(y)) \rightarrow 0$ as $n \rightarrow \infty$. As the points are moving forward by \tilde{f} along the orbits of X^c at bounded speed, this shows that the flow is locally contracted on center stable manifolds. The symmetric arguments gives local contraction for the past in center-unstable manifolds. Notice that the fact that \tilde{f} acts as a translation in all center leaves and that center leaves are fixed by \tilde{f} implies that no deck transformation can reverse orientation of the center, this implies that our initial assumption is verified.

This shows that the flow generated by X^c is expansive. Moreover, it preserves the transverse foliations \mathcal{W}^{cs} and \mathcal{W}^{cu} , which do not have singularities. Thus, the work of Paternain [Pat93] implies that the flow generated by X^c is a topological Anosov flow (see also Appendix G). \square

Putting together Theorem 5.1, Proposition 6.2, Proposition 6.5 and Proposition G.2 one finishes the proof of Theorem 6.1 and of Theorem C.

7. PROOF OF THEOREM A WITH DYNAMICAL COHERENCE

We are now ready to finish the proof of Theorem A when the diffeomorphism is assumed to be dynamically coherent.

We start by proving that, in a Seifert manifold, one can always choose a good lift in such a way that it fixes one center-stable leaf.

Proposition 7.1. *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a Seifert manifold. Suppose that f is homotopic to the identity and that the Seifert fibration in M is orientable. Then there exists a good lift of an iterate of f which fixes a leaf (and therefore every leaf) of $\tilde{\mathcal{W}}^{cs}$.*

Proof. To prove the result, partial hyperbolicity will only be used to get that M has non-zero Euler class ([HPS18, Theorem B]).

First up to taking a finite lift we assume that M is an orientable circle bundle over a higher genus (orientable) surface Σ .

Consider the leaf space \mathcal{L}^{cs} of the center-stable foliation and let δ be the deck transformation associated with the center of M . As the foliation is horizontal (see Theorem F.3), it follows that the leaf space \mathcal{L}^{cs} is homeomorphic to the reals. In addition, $\mathcal{L}^{cs}/\langle \delta \rangle$ is a circle that we will call S_δ^1 .

Consider a good lift \tilde{f} of f . The map \tilde{f} induces a homeomorphism $\hat{f}: S_\delta^1 \rightarrow S_\delta^1$. Moreover, \hat{f} commutes with the image of the homeomorphisms $\hat{\rho}(\gamma): S_\delta^1 \rightarrow S_\delta^1$ which are defined for all $\gamma \in \pi_1(M)$. These homeomorphisms are well defined up to composition with δ . So $\hat{\rho}$ naturally induces a quotient representation $\rho: \pi_1(\Sigma) \rightarrow \text{Homeo}_+(S_\delta^1)$ when using the identification $\pi_1(M)/\langle \delta \rangle \cong \pi_1(\Sigma)$. The Euler class of M coincides with the one of the representation ρ (see [CC03, Chapter 4]).

We first show that \hat{f} has rational rotation number. We proceed by contradiction: Assume that \hat{f} has irrational rotation number.

Suppose first that \hat{f} is minimal. It directly implies that \hat{f} is conjugate to an irrational rotation by a homeomorphism $h: S_\delta^1 \rightarrow S_\delta^1$. Conjugating the homeomorphisms $\rho(\gamma)$ (that is, $h^{-1}\rho(\gamma)h$), since they commute with an irrational rotation

they must commute with every rotation. Therefore the homeomorphisms $\rho(\gamma)$ are all conjugate by h to rigid rotations. This implies that $\rho: \pi_1(\Sigma) \rightarrow \text{Homeo}_+(S^1)$ is conjugate to a representation into $SO(2, \mathbb{R})$. This allows to construct a path to the trivial representation, because one can move freely along $SO(2, \mathbb{R})$ until one gets to the identity without altering the relations. Therefore the representation has zero Euler class. See [Man18, Sections 5.2 and 5.3].

If \hat{f} is not minimal, it is a Denjoy counterexample, one can see that the representation of $\pi_1(\Sigma)$ into $\text{Homeo}(S^1)$ is semi-conjugate to a representation which commutes with a minimal homeomorphism, and so it also has to have zero-Euler class (see [Man18, Section 5.2]). This proves that \hat{f} has rational rotation number.

Now we go back to the original manifold. Since in the finite cover, the corresponding map \hat{f} had rational rotation number, the same is true for \hat{f} associated with the original manifold. In particular, \hat{f} has a periodic point, which means that for some $i \neq 0$, $\delta^n \tilde{f}^i$ has a fixed point. So $\delta^n \tilde{f}^i$ is the sought good lift (note that it is a good lift because the Seifert fibration is orientable, and thus δ is in the center of $\pi_1(M)$) This finishes the proof. \square

Notice that the symmetric statement holds for $\widetilde{\mathcal{W}}^{cu}$ but a priori not for both simultaneously.

Remark 7.2. In this proof, we did not really use dynamical coherence (see [BFFPa]). One could give a slightly simpler proof that uses dynamical coherence. However, since we will need this result in the non-dynamically coherent case in [BFFPa], it is more efficient to give the general proof.

So we can now prove Theorem A.

Proof of Theorem A. If the result holds in a finite cover of M , then, by projection, it also holds in M . So, by lifting to a double cover, we may assume that M has orientable Seifert fibration. Let \tilde{f} be a good lift of some power f^k given by Proposition 7.1. Then \tilde{f} does not act as a translation on both center-stable and center-unstable leaf spaces, so is not in case (ii) of Theorem C. Thus it is in case (i) of Theorem C, i.e., f^k is a discretized Anosov flow. \square

Remark 7.3. Note that, in Theorem A, we need to take a power of f to get a discretized Anosov flow, whereas Theorem C holds for the original f . This condition is necessary, i.e., there are some dynamically coherent partially hyperbolic diffeomorphisms homotopic to the identity on a Seifert manifold that are not discretized Anosov but such that a (non-trivial) iterate is. We will give such an example below and also classify all such examples.

Consider Σ a hyperbolic surface (or orbifold) and g^t the geodesic flow on $T^1\Sigma$. Let M be a k -fold cover of T^1M obtained by unwrapping the fiber and $g_M^t: M \rightarrow M$ be a lift of g^t to M . Call $s: M \rightarrow M$ the map obtained by lifting the “rotation by 2π ” along the fiber in $T^1\Sigma$. Then for any $i = 1, \dots, k-1$, the diffeomorphism $f_{k,i} := g_M^1 \circ s^i$ is a partially hyperbolic diffeomorphism, dynamically coherent, homotopic to the identity, and it is *not* a discretized Anosov flow (but $f_{k,i}^k$ is a discretized Anosov). Notice that the action of any good lift of $f_{k,i}$ on the center stable and center unstable leaf spaces is by translations.

Now, suppose that M is a Seifert manifold and f is a dynamically coherent partially hyperbolic diffeomorphisms homotopic to the identity. Then, by Theorem A, there exists k such that f^k is a discretized Anosov flow. Thus, by the classification of Anosov flows on Seifert manifolds (see [Ghy84, Bar96]), M is a finite lift of the unit tangent bundle of an orbifold Σ and f^k is leaf conjugate to

the time-one map of the lift of the geodesic flow. Then the action of (a good lift of) f on both the central stable and central unstable leaf spaces is conjugated to the action of (a good lift of) a diffeomorphism $f_{k,i}$ as above. So f and $f_{k,i}$ are leaf conjugate.

8. COARSE DYNAMICS OF TRANSLATIONS

In this section, we consider a homeomorphism $f: M \rightarrow M$ of a hyperbolic 3-manifold that preserves a uniform, \mathbb{R} -covered foliation \mathcal{F} and acts as a translation on its leaf space. We show that the dynamics of f is comparable to the dynamics of the pseudo-Anosov flow Φ (given by Theorem D.3) that regulates \mathcal{F} . More precisely, for every periodic orbit of Φ , we show that there exists a compact core (in a lift of M) invariant by f that plays the role of the periodic orbit of Φ .

So in particular, the result of this section does not require f to be partially hyperbolic and is of independent interest. The description of the dynamics of f in periodic leaves of \mathcal{F} (if any) can be compared to the global shadowing for pseudo-Anosov homeomorphisms done in [Han85]. We will use the results obtained here to complete the proof of Theorem B in the next section.

To make this comparison precise we will introduce some more objects. Let $f: M \rightarrow M$ be a hyperbolic 3-manifold. We assume that f is homotopic to the identity, and preserves a foliation \mathcal{F} . Furthermore, we suppose that \mathcal{F} is \mathbb{R} -covered and uniform and such that a good lift \tilde{f} of f acts as a translation on the leaf space of $\tilde{\mathcal{F}}$.

Since \tilde{f} commutes with any deck transformation and acts as a translation on the leaf space of $\tilde{\mathcal{F}}$, it implies that the foliation \mathcal{F} is actually transversely orientable. Hence Theorem D.3 applies and we call Φ a transverse regulating pseudo-Anosov flow. We denote by $\tilde{\Phi}$ its lift to the universal cover \tilde{M} .

Let $\gamma \in \pi_1(M)$ be an element associated with a periodic orbit of Φ (i.e., such that there is a flow line of $\tilde{\Phi}$ invariant under γ). Let $M_\gamma := \tilde{M}/\langle \gamma \rangle$ be the cover of M associated with that deck transformation.

The foliation $\hat{\mathcal{F}}_\gamma$ lifted from \mathcal{F} to M_γ is a foliation by planes. Indeed, since Φ is regulating, each orbit of $\tilde{\Phi}$ can represent the leaf space $\mathcal{L}_{\tilde{\mathcal{F}}}$ of $\tilde{\mathcal{F}}$. Thus γ , and all of its powers, act as a translation on $\mathcal{L}_{\tilde{\mathcal{F}}}$, so no leaf of $\tilde{\mathcal{F}}$ can be fixed by a power of γ . Therefore, $\hat{\mathcal{F}}_\gamma$ is a foliation by planes (see, e.g., [Fen02] for more details).

Since \tilde{f} is a good lift of f it induces a lift \hat{f}_γ of f in M_γ .

We can now state precisely the main result of this section.

Proposition 8.1. *Let $M, f: M \rightarrow M, \mathcal{F}$ and Φ be as above.*

Then, for every $\gamma \in \pi_1(M)$ associated with a periodic orbit of Φ , there is a compact \hat{f}_γ -invariant set T_γ in M_γ which intersects every leaf of $\hat{\mathcal{F}}_\gamma$.

Moreover, if an iterate \hat{f}_γ^k of \hat{f}_γ fixes a leaf L of $\hat{\mathcal{F}}_\gamma$, and γ fixes all the prongs of a periodic orbit associated with γ , then the fixed set of \hat{f}_γ^k in L is contained in $T_\gamma \cap L$ and has negative Lefschetz index.

Remark 8.2. In fact, the proof will show that the total Lefschetz index $I_{T_\gamma \cap L}(\hat{f}_\gamma^k|_L)$ equals -1 if the periodic orbit of Φ is a regular periodic orbit, and equals $1 - p$ if the periodic orbit is a p -prong, $p \geq 3$, assuming that γ fixes the prongs of the orbit.

We also remark that, by construction, the set T_γ is *essential* in the sense that any neighborhood of it contains a curve homotopic to (a power of) γ .

To prove this proposition, we first need to explore some properties of the pseudo-Anosov flow Φ and its interaction with the foliation \mathcal{F} .

Let Λ_s and Λ_u be the weak stable and weak unstable (singular) foliations of the pseudo-Anosov flow Φ . We denote by $\tilde{\Lambda}_s$ and $\tilde{\Lambda}_u$ their lift to the universal cover.

For any leaf L of $\tilde{\mathcal{F}}$, we denote by \mathcal{G}_L^s and \mathcal{G}_L^u the one dimensional (singular) foliations obtained by intersecting the foliations $\tilde{\Lambda}_s$ and $\tilde{\Lambda}_u$ with L .

Fact 8.3. *The length along foliations \mathcal{G}_L^s and \mathcal{G}_L^u is uniformly efficient up to a multiplicative distortion at measuring distances in the leaves of $\tilde{\mathcal{F}}$. That is, the rays of \mathcal{G}_L^s and \mathcal{G}_L^u are uniform quasi-geodesics for the path metric on L .*

Proof. This fact is a consequence of the construction of the foliations $\tilde{\Lambda}_s$ and $\tilde{\Lambda}_u$. They are obtained by blowing down certain laminations that intersect the leaves of $\tilde{\mathcal{F}}_\epsilon$ along geodesics (with respect to the uniformization metric obtained via Candel's Theorem C.1). We refer to [Fen02] or [Cal07] for the construction of these laminations.

In particular, there exists a uniform $K_1 > 1$ such that for every $L \in \tilde{\mathcal{F}}$ and $y \in \mathcal{G}^s(x)$ one has

$$\ell([x, y]^{\mathcal{G}_L^s}) \leq K_1 d_L(x, y) + K_1$$

where $\ell([x, y]^{\mathcal{G}_L^s})$ denotes the length of the arc in \mathcal{G}^s joining x and y . And similarly for $\mathcal{G}^u(x)$. \square

The flow $\tilde{\Phi}$ does not preserve the foliation $\tilde{\mathcal{F}}$, but since it is transverse and regulating to the foliation, it makes sense to consider, given $L_1, L_2 \in \tilde{\mathcal{F}}$ two leaves, the map $\tau_{12}: L_1 \rightarrow L_2$ consisting in flowing along $\tilde{\Phi}$ from one leaf to the other. By construction, the map τ_{12} is a homeomorphism. Notice that since \mathcal{F} is \mathbb{R} -covered and uniform, the Hausdorff distance between L_1 and L_2 is bounded multiplicatively with the flow distance between the leaves – at least for leaves which are sufficiently apart from each other.

By convention, we will always assume that L_2 is taken to be *above* L_1 , in the sense that one has to follow the orbits of $\tilde{\Phi}$ in the positive direction to go from L_1 to L_2 .

Notice that invariance of $\tilde{\Lambda}_s$ and $\tilde{\Lambda}_u$ by $\tilde{\Phi}$ imply that the homeomorphism τ_{12} maps the foliations $\mathcal{G}_{L_1}^s$ and $\mathcal{G}_{L_1}^u$ into the the foliations $\mathcal{G}_{L_2}^s$ and $\mathcal{G}_{L_2}^u$ respectively.

When the leaves L_1, L_2 are understood, we will omit them from the notation. It is a standard fact from the dynamics of pseudo-Anosov flows and the bounded comparison between flow distance and leaves⁵ that the following holds:

Fact 8.4. *For any leaves L_1 and L_2 sufficiently far apart, the map τ_{12} expands lengths (and, equivalently distances) in \mathcal{G}^u exponentially in terms of the Hausdorff distance between L_1 and L_2 . That is, there exists a $\lambda > 0$, independent of L_1, L_2 , such that, for any $x \in L_1$ and $y \in \mathcal{G}_{L_1}^u(x)$, we have*

$$d_{L_2}(\tau_{12}(x), \tau_{12}(y)) \geq e^{\lambda d_{\text{Haus}}(L_1, L_2)},$$

as long as $d_{\text{Haus}}(L_1, L_2)$ is sufficiently big. Similarly, τ_{12}^{-1} expands the lengths in \mathcal{G}^s exponentially in terms of the Hausdorff distance between L_1 and L_2 .

The following simple result will be extremely useful for us.

⁵It is worth noting that the pseudo-Anosov property is invariant under reparametrizations of the flows.

Lemma 8.5. *Suppose that β is a deck transformation that acts freely and decreasingly on the leaf space of $\tilde{\mathcal{F}}$. Let L_1 be a leaf of $\tilde{\mathcal{F}}$. Let τ_{12} be the flow along map from L_1 to $L_2 := \beta^{-1}(L_1)$. Define $g_{\beta, L_1} := \beta \circ \tau_{12}: L_1 \rightarrow L_1$. Then, for every $K > 0$, there exists $R > 0$ such that if $d_{L_1}(x, x_1) > R$ then $d_{L_1}(x, g_{\beta, L_1}(x)) > K$.*

Remark 8.6. Notice that in this Lemma, we do not ask for β to be associated with a periodic orbit of the pseudo-Anosov regulating flow.

Proof. Suppose for a contradiction that there exists $K > 0$ and a sequence y_n escaping to infinity in L_1 and such that $d_{L_1}(y_n, g_{\beta, L_1}(y_n)) \leq K$ for all n .

Up to taking a subsequence, there exists $\gamma_n \in \pi_1(M)$ such that $\gamma_n(y_n)$ converges to y_0 in \tilde{M} .

We define a map $\tau_\beta: \tilde{M} \rightarrow \tilde{M}$ as follows: given x in \tilde{M} , it is in L a leaf of $\tilde{\mathcal{F}}$, then we let $\tau_\beta(x)$ be the intersection of the flow line of $\tilde{\Phi}$ through x with $\beta^{-1}(L)$. Notice that if $x \in L_1$, then $\tau_\beta(x) = \tau_{12}(x)$. In particular, for every n , $\tau_\beta(y_n) = \tau_{12}(y_n)$.

Since $\gamma_n(y_n)$ converges to y_0 , and τ_β consists of flowing along $\tilde{\Phi}$ a uniformly bounded amount, for n big enough, we have that $d(\tau_{12}(y_n), \tau_\beta(\gamma_n^{-1}(y_0)))$ is as small as we want. Hence, for n big enough, we have

$$d(\beta \circ \tau_{12}(y_n), \beta \circ \tau_\beta(\gamma_n^{-1}(y_0))) < 1.$$

Now, $\beta \tau_{12}(y_n) = g_{\beta, L_1}(y_n)$ is at distance less than K from y_n . Thus, after applying γ_n , we get

$$d(\gamma_n(y_n), \gamma_n \circ \beta \circ \tau_\beta(\gamma_n^{-1}(y_0))) < 1 + K.$$

Note that the map τ_β moves every point a bounded distance, the transformations γ_n, β are isometries, and, for n big enough, $d(\gamma_n(y_n), y_0)$ is very small. Therefore, $d(y_0, \gamma_n \circ \beta \circ \tau_\beta(\gamma_n^{-1}(y_0))) < K'$ for all n big enough and a fixed constant K' .

So we can extract a converging subsequence once more, and get that for any n, m big enough, the distance between $\gamma_n \beta \gamma_n^{-1}(y_0)$ and $\gamma_m \beta \gamma_m^{-1}(y_0)$ is smaller than the injectivity radius of M . It follows that

$$\gamma_m \beta \gamma_m^{-1} = \gamma_n \beta \gamma_n^{-1},$$

for all n, m large enough.

Now we use that M is hyperbolic. So β is a hyperbolic isometry of $\mathbb{H}^3 \cong \tilde{M}$. It has an axis with ideal points a, b .

Let n_0 be fixed large enough. Then, since $\gamma_m \beta \gamma_m^{-1} = \gamma_{n_0} \beta \gamma_{n_0}^{-1}$, we have that $\gamma_m(a) = \gamma_{n_0}(a)$ and $\gamma_m(b) = \gamma_{n_0}(b)$.

Let $c := \gamma_{n_0}(a)$ and $d := \gamma_{n_0}(b)$. Notice that (for all n big enough) the axis of the isometry $\gamma_n \beta \gamma_n^{-1}$ has endpoints c and d .

Let α be the generator of the group of deck transformations fixing c, d . Then, for all n large enough, $\gamma_n = \alpha^{i_n} \gamma_{n_0}$. In addition since y_n escapes compact sets but $\gamma_n(y_n)$ converges to y_0 , it follows that $|i_n|$ converges to infinity.

Notice that, since γ_{n_0} sends the axis of β to the axis of α , a power of α is conjugated to a power of β by γ_{n_0} . Now, since β acts freely on the leaf space of $\tilde{\mathcal{F}}$, then so does α . But $\gamma_n(y_n)$ converges to y_0 . So $\alpha^{i_n}(L_1) = \gamma_n \gamma_{n_0}^{-1}(L_1)$ does not escape in the leaf space, which contradicts the fact that α acts freely. \square

We remark that this proof only uses geometry of M and foliations. That is, this proof works for any regulating flow transverse to a transversely oriented, \mathbb{R} -covered, uniform foliation in a hyperbolic 3-manifold.

We will also need the following consequence of Lemma 8.5.

Lemma 8.7. *Let $\gamma \in \pi_1(M)$ be associated with a periodic orbit δ_0 of Φ . Let δ be the unique lift of δ_0 to \widetilde{M} . For any leaf L of $\widetilde{\mathcal{F}}$, let x_L be the unique intersection point of L and δ .*

For any $K > 0$, there exists $R > 0$ such that, for any leaves L_1, L_2 in $\widetilde{\mathcal{F}}$ with L_2 above $\gamma(L_1)$ and for any $x \in L_1$, we have that, if $d_{L_1}(x, x_{L_1}) > R$ then $d_{L_2}(\tau_{12}(x), x_{L_2}) > K$.

Proof. Let $K > 0$. First since the Hausdorff distance from L to γL is bounded, then the amount of flowing between them is bounded by a constant C , for any L leaf of $\widetilde{\mathcal{F}}$. Now, using the fact that the leaves are uniformly properly embedded (see [Cal07, Lemma 4.48]), there is a constant K_1 such that if $d_L(u, v) > K_1$ then $d_{\widetilde{M}}(u, v) > K + 2C$.

Let L_1 be a leaf in $\widetilde{\mathcal{F}}$ and let L_2 be such that γL_1 separates L_1 from L_2 . Then there exists $n > 0$ such that L_2 is in the interval (in the leaf space of $\widetilde{\mathcal{F}}$) between $\gamma^n L_1$ and $\gamma^{n+1} L_1$.

By Lemma 8.5 and the fact that deck transformations are isometries in the leafwise metrics, there exists R such that for any $x \in L_1$, if $d_{L_1}(x, x_{L_1}) > R$ then $d_{\gamma^n L_1}(\tau_n(x), x_{\gamma^n L_1}) > K_1$, where $\tau_n: L_1 \rightarrow \gamma^n L_1$ is the flow along Φ map. By the choice of K_1 it follows that $d_{\widetilde{M}}(\tau_n(x), x_{\gamma^n L_1}) > K + 2C$.

Now, since L_2 is between $\gamma^n L_1$ and $\gamma^{n+1} L_1$, the flow distance from $\gamma^n L_1$ to L_2 is bounded above by C . It follows that $d_{\widetilde{M}}(\tau_{12}(x), x_{L_2}) > K$. In particular $d_{L_2}(\tau_{12}(x), x_{L_2}) > K$, which proves the lemma. \square

The reason we will be able to compare the dynamics of \widetilde{f} and $\widetilde{\Phi}$ is thanks to the fact that they are a uniform bounded distance apart. That is, we have the following.

Lemma 8.8. *Let $f: M \rightarrow M$ be a homeomorphism of a hyperbolic 3-manifold M preserving an \mathbb{R} -covered uniform foliation \mathcal{F} and \widetilde{f} a good lift to \widetilde{M} . There exists $R_1 > 0$ so that for every $L_1 \in \widetilde{\mathcal{F}}$, if $L_2 = \widetilde{f}(L_1)$ and $x \in L_1$ then $d_{L_2}(\widetilde{f}(x), \tau_{12}(x)) < R_1$.*

Proof. Since \widetilde{f} is a good lift it follows that one can join x with $\widetilde{f}(x)$ by an arc of bounded length. In particular, since the foliation \mathcal{F} is \mathbb{R} -covered and uniform, it follows that the Hausdorff distance between L_1 and $L_2 = \widetilde{f}(L_1)$ is uniformly bounded above and below independently of $L_1 \in \widetilde{\mathcal{F}}$. Therefore, as explained before the statement of Fact 8.4 the amount of flowing needed to go from L_1 to L_2 is also uniformly bounded below and above.

It follows that $d_{\widetilde{M}}(\widetilde{f}(x), \tau_{12}(x))$ is uniformly bounded. Again we use the fact that leaves of an \mathbb{R} -covered taut foliation are uniformly properly embedded in the universal cover (see [Cal07, Lemma 4.48]). The result follows. \square

Now we are ready to prove Proposition 8.1.

Proof of Proposition 8.1. Let $\gamma \in \pi_1(M)$ be represented by a periodic orbit δ_0 of Φ and take δ the unique lift of δ_0 to \widetilde{M} fixed by γ .

We will build the core T_γ that we seek by considering a very large tubular neighborhood of δ and taking the intersection of this tubular neighborhood with all of its forward and backwards image under \widetilde{f} (see figure 11). We will prove that this infinite intersection is non-empty, thus its projection to M_γ will have the desired properties.

Note that, if we build the core for a power γ^{k_1} and \widetilde{f}^{k_2} instead, then taking its intersection with its images by $\gamma, \dots, \gamma^{k_1-1}$ and $\widetilde{f}, \dots, \widetilde{f}^{k_2-1}$ will give a core

with the properties we want. So, in this proof we may take any finite power of γ or \tilde{f} .

Thus, if δ_0 is a p -prong, we replace γ by a power if necessary, so that $\tilde{\gamma}$ fixes every prongs of δ . Furthermore, we take a power of f so that for any L , $\tilde{f}(L)$ is above $\gamma(L)$. For notations sake, we assume this is the original f .




FIGURE 11. The image of a large tubular neighborhood of the lift of the prong by \tilde{f} in a given center stable leaf.

As in Lemma 8.7 for any leaf L in $\tilde{\mathcal{F}}$, we write x_L to be the (unique) intersection of δ with L .

Let a_L^i , with $i = 1, \dots, p$, be all the ideal points on the boundary at infinity of L of the weak unstable leaf (of $\tilde{\Phi}$) through δ , where $p = 2$ if δ is a regular orbit and otherwise δ is a p -prong orbit. Equivalently, a_L^i is the ideal point determined by each ray of $\mathcal{G}_L^u(x_L)$.

Similarly, we define r_L^i , $i = 1, \dots, p$, to be the ideal ends of the rays of $\mathcal{G}_L^s(x_L)$.

For every $L \in \tilde{\mathcal{F}}$ and for every i , we choose P_L^i and N_L^i neighborhoods (in $L \cup \partial_\infty L$) of, respectively a_L^i and r_L^i . We also choose these neighborhoods such that their boundary (in L) are geodesics for the path metric on L . Furthermore, we choose these neighborhoods in such a way that they depend continuously on $L \in \tilde{\mathcal{F}}$ and they are γ -invariant, i.e., $\gamma(P_L^i) = P_{\gamma(L)}^i$ and $\gamma(N_L^i) = N_{\gamma(L)}^i$.

Up to taking the neighborhoods smaller, we assume that for any L and any i, j , $P_L^i \cap N_L^j = \emptyset$; and for any $i \neq j$, $P_L^i \cap P_L^j = \emptyset$, $N_L^i \cap N_L^j = \emptyset$.

We define a map $\tau_f: \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$ in the following way: For any L in $\tilde{\mathcal{F}}$ and any $x \in L$, $\tau_f(x)$ is the intersection of the orbit of $\tilde{\Phi}$ through x with $\tilde{f}(L)$.

Let R_1 be the constant given by Lemma 8.8 (i.e., such that \tilde{f} and τ_f are R_1 -close). Using that leaves of $\mathcal{G}_L^u(x_L)$ and $\mathcal{G}_L^s(x_L)$ are quasigeodesics and that the flow in a given time expands and contracts their length by a factor different from one, one can construct neighborhoods P_L^i and N_L^i sufficiently small so that:

- (i) For any L and any i ,

$$\tau_f(P_L^i) \subset P_{\tilde{f}(L)}^i \text{ and } d_{\tilde{f}(L)}\left(\tau_f(P_L^i), \partial P_{\tilde{f}(L)}^i\right) > 10R_1.$$

(ii) For any L and any i ,

$$\tau_f^{-1}(N_L^i) \subset N_{\tilde{f}^{-1}(L)}^i \text{ and } d_{\tilde{f}^{-1}(L)}\left(\tau_f^{-1}(N_L^i), \partial N_{\tilde{f}^{-1}(L)}^i\right) > 10R_1.$$

To make sense of the distance between sets in $L \cup \partial_\infty L$ above, we decide that ideal points are at infinite distance from any other point. A direct consequence of the conditions above is that

- (1) For any L and any i , $\tilde{f}(P_L^i) \subset P_{\tilde{f}(L)}^i$
- (2) For any L and any i , $\tilde{f}^{-1}(N_L^i) \subset N_{\tilde{f}^{-1}(L)}^i$

Lemma 8.8 shows that, for any L , the maps $\tau_f|_L$ and $\tilde{f}|_L$ are a finite distance from each other. Thus their extension to the circles at infinity $\partial_\infty L$ is the same. Now, recall that τ_f corresponds to flowing along the pseudo-Anosov flow $\tilde{\Phi}$. Hence, up to replacing \tilde{f} by a very high power of \tilde{f} , we can moreover assume that:

- (3) For any L and i , if ω is an ideal endpoint of ∂N_L^i , then $\tilde{f}(\omega) \in P_{\tilde{f}(L)}^j$, for some j (where j is the unique index such that a_L^j is the first attractor on the side of ω from r_L^i);
- (4) For any L and i , if ω is an ideal endpoint of ∂P_L^i , then $\tilde{f}^{-1}(\omega) \in N_{\tilde{f}^{-1}(L)}^j$, for some j (where j is uniquely determined as above).

Note that conditions (1) and (2) still are satisfied by our high power of \tilde{f} .

Now we choose a constant R large enough so that for every L and every i , the ball $D_L := B(x_L, R)$, of radius R around x_L , intersects every P_L^i . Moreover, we choose it to satisfy:

$$\begin{aligned} D_L \supset \tilde{f}\left(\partial N_{\tilde{f}^{-1}(L)}^i\right) \setminus \bigcup_j P_L^j \\ D_L \supset \tilde{f}^{-1}\left(\partial P_{\tilde{f}(L)}^i\right) \setminus \bigcup_j N_L^j \end{aligned}$$

This is possible because the ideal points of $A = \tilde{f}(\partial N_{\tilde{f}^{-1}(L)}^i)$ are contained in the interior of the ideal boundary of the union of the P_L^j . It follows that for each L only a compact part of $A \cap L$ is outside the union, and this varies continuously with L . By choosing R big enough one satisfies the equations above.

Let

$$V := \bigcup_{L \in \tilde{\mathcal{F}}} D_L.$$

We will show that the set $\bigcap_{n \in \mathbb{Z}} \tilde{f}^n(V)$ is non-empty and thus its projection to M_γ is the core T_γ that we seek.

The proof will be done by induction. In order to make that induction work, we need the following

Claim 8.9. *Let L be a leaf in $\tilde{\mathcal{F}}$. Let $C \subset D_L$ be any compact and path-connected set that does not intersect any N_L^i .*

If there exists i_1, i_2 distinct such that C intersect both $P_L^{i_1}$ and $P_L^{i_2}$, then there exists a path-connected component of $\tilde{f}(C) \cap D_{\tilde{f}(L)}$ that intersects $P_{\tilde{f}(L)}^{i_1}$ and $P_{\tilde{f}(L)}^{i_2}$, for some $j_2 \neq i_1$ (j_2 is not necessarily i_2) and that does not intersect any $N_{\tilde{f}(L)}^i$.

Proof. Since C intersects $P_L^{i_1}$ and $P_L^{i_2}$, $\tilde{f}(C)$ also intersects both $P_{\tilde{f}(L)}^{i_1}$ and $P_{\tilde{f}(L)}^{i_2}$ (thanks to the condition (1)).

Now, since C does not intersect any N_L^i , because of condition (3) and the choice of D_L , the intersections of $\tilde{f}(C)$ with $\partial D_{\tilde{f}(L)}$ are contained in the union of the $P_{\tilde{f}(L)}^i$.

Thus, as claimed, $\tilde{f}(C) \cap D_{\tilde{f}(L)}$ contains a connected component that intersects $P_{\tilde{f}(L)}^{i_1}$ and $P_{\tilde{f}(L)}^{j_2}$, for some $j_2 \neq i_1$ (j_2 is not necessarily i_2) and that does not intersect any $N_{\tilde{f}(L)}^i$. \square

Figure 12 shows a case where j_2 is not equal to i_2 : It may be that $\tilde{f}(C)$ stretches well into $P_{\tilde{f}(L)}^{j_2}$ and out of $D_{\tilde{f}(L)}$. Thus, as in the figure, the intersection $\tilde{f}(C) \cap D_{\tilde{f}(L)}$ can have two components C_1 and C_2 , neither of which intersects both $P_{\tilde{f}(L)}^{i_1}$ and $P_{\tilde{f}(L)}^{i_2}$.




FIGURE 12. The intersection $\tilde{f}(C) \cap D_{\tilde{f}(L)}$ may not have a connected set joining P^{i_1} to P^{i_2} .

The same proof as above, using \tilde{f}^{-1} instead (and the conditions (2) and (4)), gives

Claim 8.10. *Let L be a leaf in $\tilde{\mathcal{F}}$. Let $C \subset D_L$ be any compact and path-connected set that does not intersect any $P_{\tilde{f}(L)}^i$.*

If there exists i_1, i_2 distinct such that C intersect both $N_L^{i_1}$ and $N_L^{i_2}$, then there exists a path-connected component of $\tilde{f}^{-1}(C) \cap D_{\tilde{f}^{-1}(L)}$ that intersects $N_{\tilde{f}^{-1}(L)}^{i_1}$ and $N_{\tilde{f}^{-1}(L)}^{j_2}$, for some $j_2 \neq i_1$ and that does not intersect any $P_{\tilde{f}^{-1}(L)}^i$.

For any leaf L and any integer $n \geq 0$, define

$$R_L^n = \bigcap_{k=0}^n \tilde{f}^k(D_{\tilde{f}^{-k}(L)}) \quad \text{and} \quad Q_L^n = \bigcap_{k=0}^n \tilde{f}^{-k}(D_{\tilde{f}^k(L)}).$$

We will show:

Claim 8.11. *For every i and every n , R_L^n contains a subset C , compact and path-connected that does not intersect any N_L^j but does intersect P_L^i and some $P_L^{i_2}$ (for some $i_2 \neq i$).*

Similarly, for every i and every n , Q_L^n contains a subset C , compact and path-connected that does not intersect any P_L^j but does intersect N_L^i and some $N_L^{i_2}$ (for some $i_2 \neq i$).

Proof. We only do the proof for R_L^n , as the claim for Q_L^n follows similarly.

First, since $R_L^0 = D_L$, the claim is true for $n = 0$ and any leaf L (because D_L clearly contains such a subset). Let us assume that the claim holds for R_L^{n-1} and for any L .

Then, Claim 8.9 implies that (for any L) $\tilde{f}(R_L^{n-1}) \cap D_{\tilde{f}(L)}$ has a compact and path-connected subset that does not intersect any N_L^j but does intersect P_L^i and some $P_L^{i_2}$ (for some $i_2 \neq i$).

But, by definition, we have

$$R_L^n = \bigcap_{k=0}^n \tilde{f}^k(D_{\tilde{f}^{-k}(L)}) = \tilde{f}\left(R_{\tilde{f}^{-1}(L)}^{n-1}\right) \cap D_L.$$

Thus the claim is proved. \square

Now, since for any L , the ideal points a_L^i and r_L^i alternate, the properties of R_L^n and Q_L^n given by Claim 8.11 imply that, for all n , $R_L^n \cap Q_L^n$ is a non-empty compact set.

Since R_L^n and Q_L^n are decreasing sets, the set

$$T_L := \bigcap_{n \geq 0} (R_L^n \cap Q_L^n)$$

is (for any L) non-empty and compact. Thus

$$T := \bigcup_{L \in \tilde{\mathcal{F}}} T_L$$

is non-empty, and, by construction, \tilde{f} -invariant (note also that $T = \bigcap_{n \in \mathbb{Z}} \tilde{f}^n(V)$ as we claimed above).

Hence, the projection T_γ of T to M_γ is non-empty, compact and \hat{f}_γ -invariant.

Once T_γ is built, the second half of Proposition 8.1 follows directly from the homotopy invariance of Lefschetz index together with Lemma 8.8. So we finished the proof of Proposition 8.1. \square

In the proof of Proposition 8.1, we obtained the following result which we state independently for future reference.

Lemma 8.12. *Let $f: M \rightarrow M$ be a homeomorphism of a hyperbolic 3-manifold, f homotopic to the identity. Suppose that f preserves an \mathbb{R} -covered uniform foliation \mathcal{F} and that a good lift \tilde{f} of f acts as a translation on the leaf space of $\tilde{\mathcal{F}}$. Let $\gamma \in \pi_1(M)$ be a deck transformation.*

If $h = \gamma \circ \tilde{f}^n$ fixes some leaf $L \in \tilde{\mathcal{F}}$ (with $n \neq 0$) then the set of fixed points of h in L is contained in a compact subset of L .

Moreover, given $n > 0$ big enough, then for every $R > 0$ there is a compact set $D \subset L$ such that if $y \notin D$ then $d_L(y, h(y)) > R$.

Finally, let P be the set of ideal points in the boundary at infinity $S^1(L)$ that are attracting and fixed under the map $\gamma \circ \tau_{12}$, where $\tau_{12}: L \rightarrow \tilde{f}^n(L)$ is the

flow along Φ map. Then, for any $y \in P$, there exists a neighborhood U of y in $L \cup S^1(L)$ such that

- (1) $h(\bar{U})$ is strictly contained in \bar{U} , and
- (2) $\bigcap_{i \geq 0} h^i(U) = \{y\}$.

Remark 8.13. Notice that the results of this section should be adaptable to the case of a homeomorphism acting as a translation on the leaf space of a manifold with one atoroidal piece. What would be required is some sort of analogue of Theorem D.3. That is, we would need to know that there exists a transverse regulating flow such that any orbit that stays in the atoroidal piece is a hyperbolic p -prong ($p \geq 2$). Although that result seems likely to be true, it has not been proven. A similar case is dealt with in a companion paper [BFFP20] where we study integrability for partially hyperbolic diffeomorphisms not homotopic to identity in Seifert manifolds.

9. DOUBLE TRANSLATIONS IN HYPERBOLIC MANIFOLDS

In this section we prove Theorem B.

Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism of a hyperbolic 3-manifold M . Recall that we denote by \mathcal{W}^{cs} and \mathcal{W}^{cu} a pair of f -invariant foliations tangent respectively to E^{cs} and E^{cu} . Up to taking an iterate, one has that f is homotopic to identity and therefore has a good lift \tilde{f} to M . We fix that good lift.

We want to show that \tilde{f} fixes the leaves of both foliations $\widetilde{\mathcal{W}}^{cs}$ and $\widetilde{\mathcal{W}}^{cu}$. By Theorem 6.1 this is enough to prove Theorem B. Notice that by Corollary 3.21 and Theorem 5.1 we can assume by contradiction that both foliations are \mathbb{R} -covered and uniform and that \tilde{f} acts as translation on both leaf spaces.

We in fact will get a contradiction using just one of the translations thanks to the Proposition 8.1, together with the following result. Notice that we thus obtain an alternative proof, albeit much more complicated, of the fact that there cannot be a mixed behavior in a hyperbolic manifold.

Proposition 9.1. *Assume that a good lift \tilde{f} of f acts as a translation on the foliation $\widetilde{\mathcal{W}}^{cs}$ and let Φ be a transverse regulating pseudo-Anosov flow for \mathcal{W}^{cs} . Then, for every $\gamma \in \pi_1(M)$ associated to the inverse of a periodic orbit γ of Φ there is $n > 0, m > 0$ such that $h = \gamma^n \circ \tilde{f}^m$ fixes a leaf L of $\widetilde{\mathcal{W}}^{cs}$.*

By symmetry, the same result holds if applied to \mathcal{W}^{cu} . Notice that once one knows that h fixes a leaf L of $\widetilde{\mathcal{W}}^{cs}$, the second part of Proposition 8.1 applies to f .

Proof. Thanks to Proposition 8.1, we can consider the cover $M_\gamma = \widetilde{M} / \langle \gamma \rangle$ and let V be a compact solid torus in M_γ such that $\bigcap_{n \in \mathbb{Z}} \hat{f}_\gamma^n(V) = T_\gamma$ is compact and far from ∂V .

Let $z \in T_\gamma$. Let $y \in T_\gamma$ be an accumulation point of $(\hat{f}_\gamma^n(z))$.

Take i, j big enough, with j much bigger than i , such that $\hat{f}_\gamma^i(z)$ and $\hat{f}_\gamma^j(z)$ are both very close to y .

Consider t a small closed unstable segment containing $\hat{f}_\gamma^i(z)$ in its interior.

Since \hat{f}_γ^{j-i} increases the unstable length, every leaf of $\hat{\mathcal{W}}^{cs}$ through t intersects the interior of $\hat{f}_\gamma^{j-i}(t)$. This set of $\hat{\mathcal{W}}^{cs}$ leaves is an interval. This produces a fixed $\hat{\mathcal{W}}^{cs}$ leaf under \hat{f}_γ^{j-i} . Lifting to \widetilde{M} proves the proposition. \square

We can now finish the proof of Theorem B.

Proof of Theorem B. Let \tilde{f} be a good lift of f and let L_0 be a leaf fixed by $h := \gamma \circ \tilde{f}^k$ for some $k > 0$ and $\gamma \in \pi_1(M) \setminus \{\text{id}\}$ given by Proposition 9.1.

For any leaf L fixed by h , the map h has negative Lefschetz index (according to Proposition 8.1). Thus there exists a point $x_L \in L$ fixed by h . Now, h is partially hyperbolic, so any fixed leaf L is repelling along the unstable manifold through x_L .

But this is impossible, as in the leaf space of $\widetilde{\mathcal{W}}^{cs}$, the closed interval between L_0 and $\gamma(L_0)$ is fixed so cannot contain only repelling fixed points.

This contradiction implies that \tilde{f} cannot act as a translation on either leaf spaces. It follows that \tilde{f} has to fix every center stable and center unstable leaf. Therefore by Theorem 6.1, it is conjugate to a discretized Anosov flow. This proves Theorem B. \square

APPENDIX A. SOME 3-MANIFOLD TOPOLOGY

We collect here some concepts from 3-manifold topology that were used in this article. We refer the reader to [Hem76, Hat, Jac80] for more background.

A 3-manifold (which we always mean to be a smooth manifold) is *irreducible*, if every smoothly embedded sphere bounds a ball. It is well known that closed 3-manifolds admitting taut foliations are irreducible (see [Ros68]).

An irreducible compact, 3-manifold M is said to be *homotopically atoroidal* if every π_1 -injective map of a torus in M is homotopic to a map into the boundary of M . The manifold is *geometrically atoroidal* if every π_1 -injective, embedded smooth torus is homotopic to the boundary of M .

If a manifold with exponential growth of fundamental group is homotopically atoroidal, then by the Geometrization Theorem of Perelman [Per02, Per03b, Per03a] it is *hyperbolic*, i.e., the interior of M admits a complete, Riemannian metric of constant negative curvature.

Notice that when M is homotopically atoroidal, $\pi_1(M)$ does not contain any subgroup isomorphic to \mathbb{Z}^2 .

A 3-manifold is called a *Seifert manifold* if it admits a partition by distinct circles such that a tubular neighborhood of each fiber is homeomorphic by a fiber-preserving homeomorphism to either:

- A fibered solid torus of type (p, q) . This is a torus obtained from $\mathbb{D}^2 \times [0, 1]$ by identifying $\mathbb{D}^2 \times \{0\}$ to $\mathbb{D}^2 \times \{1\}$ via the map $(z, 0) \mapsto (z \exp(2\pi ip/q), 1)$. The fiber $\{0\} \times S^1$ is called *regular* if $p = 0$ and *exceptional* otherwise. Or,
- A fibered solid Klein bottle, obtained from $\mathbb{D}^2 \times [0, 1]$ by identifying $\mathbb{D}^2 \times \{0\}$ to $\mathbb{D}^2 \times \{1\}$ via the map $(z, 0) \mapsto (\bar{z}, 1)$. The fibers $\{z\} \times S^1$, $z \in \mathbb{R}$, are also called *exceptional*.

The quotient of a Seifert manifold by the Seifert fibration, called the *base*, B , has a structure of a 2-orbifold (without corner reflectors). The exceptional fibers separate into two sets: The axis of the fibered solid torus projected to isolated points in the interior of B (called *conical points*), while the exceptional fibers coming from fibered solid Klein bottles projects to a closed 1-submanifold of the boundary of B (and each connected component is called a *reflector curve*).

Putting together work of Epstein [Eps72] and Tollefson [Tol78], one can notice that a 3-manifold is Seifert if and only if it admits a foliation by circles.

Remark A.1. The definition above is not the one originally taken by Seifert. Indeed, the fibered solid Klein bottles neighborhood were not allowed in the original definition. However, it is now more common to use this definition (see,

e.g., [Sco83]). In particular with this definition then all 3-manifolds foliated by circles are Seifert.

Note that both the original definition and the one chosen here agree when the manifold is assumed orientable.

If a Seifert manifold has fundamental group with exponential growth, then it is finitely covered by a circle bundle over a surface of genus ≥ 2 . In particular, thanks to the classification of Seifert manifolds (see [Sco83, Theorem 3.8]), the Seifert fibration is unique in this case.

If a 3-manifold M is geometrically atoroidal but not homotopically atoroidal then the proof of the Seifert fibered conjecture [CJ94, Gab92] implies that M is closed and Seifert. The base surface has to be a sphere with 3-singular fibers. Unless the difference between geometric and homotopic atoroidal is essential we only refer to it as atoroidal.

The JSJ decomposition theorem implies that compact, irreducible, and orientable 3-manifolds admit a decomposition into finitely many pieces, which are either geometrically atoroidal or Seifert [Hem76, Hat].

The following lemma was used when establishing minimality of foliations in Seifert and hyperbolic 3-manifolds:

Lemma A.2. *If T is an embedded torus inside an orientable closed hyperbolic 3-manifold M , then T either bounds a solid torus or is contained in a 3-dimensional ball.*

Proof. This is standard result in 3-manifold topology, so we only sketch the proof. Since M is orientable and hyperbolic, T is two sided, and not π_1 -injective. Since T is furthermore embedded, Dehn's lemma [Hem76] implies that there is a compressing disk D . That is, D is embedded and $D \cap T = \partial D$. Cutting T along D and capping with two copies of D produces a sphere. Since M is hyperbolic, it is irreducible [Ros68], so it follows that the sphere bounds a ball. This implies that either T bounds a solid torus or T is contained in a 3-ball [Hem76, Jac80]. \square

We also use the following consequence of Mostow rigidity [Mos68]

Proposition A.3. *If M is a hyperbolic 3-manifold and $f: M \rightarrow M$ a homeomorphism, then it has an iterate which is homotopic to identity.*

Proof. Mostow rigidity [Mos68] implies that every homeomorphism is homotopic to an isometry. Isometries in a compact manifold have iterates which are close to the identity, so homotopic to them. \square

APPENDIX B. TAUT FOLIATIONS IN 3-MANIFOLDS

All the foliations considered in this article are continuous foliations, with C^1 leaves, tangent to a continuous distribution of a 3-manifold (so they are foliations of regularity $C^{0,1+}$ in the terminology of [CC00]). In this appendix, all foliations are 2-dimensional.

A foliation on M is called *taut* if it admits a closed transversal that intersects every leaf of \mathcal{T} .

An important consequence of Novikov's theorem [Nov65] is that if a 2-dimensional foliation of a 3-manifold does not have compact leaves then it is taut (see, e.g., [CC00, CC03, CLN85, Cal07]).

Let $\tilde{\mathcal{T}}$ denote the lift of the foliation \mathcal{T} to \tilde{M} . The *leaf space* of \mathcal{T} is defined as the set $\mathcal{L}_{\tilde{\mathcal{T}}} := \tilde{M}/\tilde{\mathcal{T}}$ equipped with the quotient topology.

The following theorem gathers some known properties of taut foliations (see, for instance, [Cal07, Chapter 4] for the proofs) and relies particularly on the celebrated theorems by Novikov [Nov65] and Palmeira [Pal78].

Theorem B.1. *A foliation without compact leaves in a 3-manifold M is taut.*

If M is a 3-manifold that is not finitely covered by $S^2 \times S^1$ and admitting a taut foliation \mathcal{T} ⁶ then \widetilde{M} is homeomorphic to \mathbb{R}^3 . Moreover, every leaf of \mathcal{T} lifts to a plane $L \in \widetilde{\mathcal{T}}$ which is properly tamely embedded in \widetilde{M} and separates \widetilde{M} in two half spaces.

The leaf space $\mathcal{L}_{\widetilde{\mathcal{T}}}$ is a one dimensional (non necessarily Hausdorff), simply connected (separable) manifold. Furthermore, every point in $\mathcal{L}_{\widetilde{\mathcal{T}}}$ is contained in the interior of an interval in $\mathcal{L}_{\widetilde{\mathcal{T}}}$.

In particular, if β is a transversal to $\widetilde{\mathcal{T}}$, then β intersects a leaf of $\widetilde{\mathcal{T}}$ at most once.

When $\mathcal{L}_{\widetilde{\mathcal{T}}}$ is Hausdorff, then it is homeomorphic to the real numbers \mathbb{R} . In this case, the foliation \mathcal{T} is called \mathbb{R} -covered.

Since \widetilde{M} is simply connected, $\widetilde{\mathcal{T}}$ is transversely orientable (but deck transformations of \widetilde{M} may flip this transverse orientation).

For reference, we cite the following result that we used in this article (for the C^0 version of the arguments used, see [CC00, §9]).

Proposition B.2 (Rosenberg [Ros68]). *Let M be a closed 3-manifold which is not \mathbb{T}^3 , and let \mathcal{T} be a foliation on M . Then some leaf of \mathcal{T} is not a plane.*

APPENDIX C. UNIFORMIZATION OF LEAVES

The following result is very helpful to understand the action of deck transformations inside leaves of the foliations of a partially hyperbolic diffeomorphism.

Theorem C.1 (Candel [Can93]). *Let \mathcal{F} be a taut foliation of a 3-manifold M and assume that it has no transverse invariant measure. Then, there is a metric in M which restricts to a (2-dimensional) hyperbolic metric in each leaf of \mathcal{F} .*

A transverse invariant measure is an assignment of a non negative number to each arc transverse to \mathcal{F} , such that it satisfies the properties of measures under countable union and restriction. In addition the measure is unchanged if we homotope the transverse arcs keeping each point in its respective leaf. The statement of [Can93] gives further properties on the transverse invariant measure (also called holonomy invariant transverse measure), stating that it has zero Euler characteristic, but we will avoid defining this (see [Cal07] for a detailed treatment).

It is well known that every taut foliation in a hyperbolic 3-manifold (see e.g., [Cal07]) or the horizontal foliations in a Seifert manifold used in this article satisfy the conclusion of Theorem C.1. We remark that it is possible to show that any minimal foliation on a manifold with non-virtually solvable fundamental group satisfies the hypothesis of Theorem C.1 (see [FP18, Section 5.1]).

⁶Note that since M is not finitely covered by $S^2 \times S^1$ no leaves of \mathcal{T} can be a sphere or a projective plane.

APPENDIX D. UNIFORM FOLIATIONS AND TRANSVERSE PSEUDO-ANOSOV FLOWS

Uniform foliations were introduced by Thurston [Thu], and have been intensively studied, particularly when M is a hyperbolic 3-manifold. They are intimately related to the notion of *slitherings* (see [Thu] or [Cal07, Chapter 9]).

Definition D.1. An \mathbb{R} -covered foliation \mathcal{T} is called *uniform* if the Hausdorff distance between any pair of leaves L, L' of $\widetilde{\mathcal{T}}$ is finite. That is, there exists $K > 0$ (depending on L and L') such that $L \subset B_K(L')$ and $L' \subset B_K(L)$ where $B_K(X)$ denotes the set of points at distance less than K from $X \subset \widetilde{M}$.

Thurston build a special pseudo-Anosov flow associated with a \mathbb{R} -covered foliation in a hyperbolic manifold.

Definition D.2. Let \mathcal{F} be a foliation of a 3-manifold. A flow $\Psi: M \rightarrow M$ is called *regulating* for \mathcal{F} if every orbit of the lifted flow $\widetilde{\Psi}$ intersects every leaf of the lifted foliation $\widetilde{\mathcal{F}}$ in the universal cover \widetilde{M} .

Theorem D.3 (Thurston, Calegari, Fenley [Thu, Cal00, Fen02]). *A transversely oriented, \mathbb{R} -covered, uniform foliation in a hyperbolic 3-manifold admits a regulating transverse pseudo-Anosov flow Φ . Moreover, Φ can be chosen so that the singular foliations have C^1 leaves outside the prongs.*

Recall that a pseudo-Anosov flow Φ is a flow generated by a vector field X which preserves two singular foliations Λ^s and Λ^u and such that, outside a finite number of singular orbits, the flow is locally modeled on a (topological) Anosov flow (see Appendix G). The foliations glue along the singularities forming p -prongs (with $p \geq 3$). We refer the reader to [Cal07] for more details. We note also that every *expansive* flow is orbit equivalent to a pseudo-Anosov flow [Pat93, IM90].

Work of Barbot and the second author implies that Thurston's regulating flow is genuinely pseudo-Anosov:

Proposition D.4. *If Φ is a pseudo-Anosov flow regulating and transverse to a uniform foliation in a non virtually solvable 3-manifold, then, Φ is not a topological Anosov flow. In particular, there are singular periodic orbits which are p -prongs with $p \geq 3$.*

Proof. This fact can be found in [Fen13], but we recall the argument: every element of the fundamental group that represents a periodic orbit of Φ acts as a translation on the leaf space of $\widetilde{\mathcal{F}}$. However, this is inconsistent with the fact that every Anosov flow on a 3-manifold, except for suspensions of Anosov diffeomorphisms (which do not exist on non virtually solvable manifolds), admits pairs of periodic orbit that are freely homotopic to the inverse of each other (this fact follows from work of Barbot and the second author, see [BBGR17, Theorem 2.15]). \square

APPENDIX E. AXES

Here, we recall some needed results from the theory of *axes* for free actions on one-dimensional, non-Hausdorff, simply connected, manifolds. These results extend similar results for trees. We refer the reader to [Fen03, Bar98] for a more detailed account. All of the results we state are true for homeomorphisms of one-dimensional, non-Hausdorff, simply connected, separable manifolds. However, to

keep the terminology close to the core of this article, we phrase our results in the setting of homeomorphisms preserving a foliation.

Let L be a complete plane. Let \mathcal{C} be a foliation such that its leaf space $\mathcal{L}_{\mathcal{C}}$ is a one dimensional (not necessarily Hausdorff) simply connected manifold.

Definition E.1 (Axis of a foliation-preserving homeomorphism). Let $g: L \rightarrow L$ be a homeomorphism which preserves \mathcal{C} . The *axis* of g (or, if one needs to be more precise, the *\mathcal{C} -axis* of g) is the set of leaves $c \in \mathcal{C}$ such that $g(c)$ separates c from $g^2(c)$.

For the statement below, we recall that a \mathbb{Z} -union of intervals means an ordered set consisting of countably many closed (possibly degenerate) intervals which are ordered according to \mathbb{Z} .

Proposition E.2. *Let $g: L \rightarrow L$ be a homeomorphism that preserves \mathcal{C} without leaving any leaf of \mathcal{C} fixed. Then, the \mathcal{C} -axis for the action of g in $\mathcal{L}_{\mathcal{C}}$ is non empty. In addition, the axis is either a line or an ordered \mathbb{Z} -union of intervals. In the second case, the axis is $\cup I_i$, where $I_i = [x_i, y_i]$ is a closed interval and y_i is not separated from x_{i+1} in the leaf space of \mathcal{C} .*

Moreover, suppose that $g, h: L \rightarrow L$ are two \mathcal{C} -preserving homeomorphisms that do not fix any \mathcal{C} -leaves, and that share the same axis. If the group generated by g and h acts freely⁷ on this axis, then it is abelian.

Proof. This is proven in section 3 of [Fen03]. See in particular Lemma 3.5, Theorem 3.8 and Proposition 3.10 there. The last statement uses Hölder's Theorem (see e.g., [Nav11, Section 2.2.4]) to deduce that the group generated by g and h must be abelian. \square

Remark E.3. Two commuting homeomorphisms that act freely have the same axis (see [Bar98, Section 2] or [Fen03, Section 3]). We remark a couple of subtle points:

- (1) The fact that f acts freely does not imply that any power of f acts freely, and in fact there are easy counterexamples,
- (2) Unlike in the case of trees, f acting freely does not necessarily imply that the axis is properly embedded in the leaf space. If the axis is a bi-infinite union of intervals, then it is properly embedded (see Lemma E.5 below). If the axis is the reals, it may fail to be properly embedded, even if all powers of f act freely.

Notice that as a consequence we obtain:

Corollary E.4. *Let $f, g, h: L \rightarrow L$ be three \mathcal{C} -preserving homeomorphisms such that the action of the group is free in the \mathcal{C} -leaf space. Assume moreover that f commutes with g and with h . Then, the group generated by f, g, h is abelian.*

Proof. Notice that as f commutes with g and h then all three homeomorphisms of the leaf spaces share the same axis ([Bar98, Section 2]). Now the result follows from the previous proposition. \square

Another useful fact about axes is the following:

Lemma E.5. *If A is the axis of a \mathcal{C} -preserving homeomorphism f and A is a \mathbb{Z} -union of intervals, then A is properly embedded in the leaf space of \mathcal{C} .*

⁷Recall that we say that a group acts *freely* if no element different from the identity has a fixed point.

Proof. Let $A = \cup_{\mathbb{Z}} I_i$, where $I_i = [x_i, y_i]$, with y_i and x_{i+1} not separated in the leaf space of \mathcal{C} .

If A is not properly embedded, then there exists a leaf $c \in \mathcal{C}$ such that (x_i) and (y_i) converges to c as, say, n goes to $+\infty$. Now, for any i , the interval I_i separates I_{i-1} from I_{i+1} . Thus, if τ is a transversal to \mathcal{C} through c , then τ intersects every I_i , for i big enough. So in particular, for some i big enough, τ intersects both y_i and x_{i+1} . But this is impossible since these two leaves are not separated. \square

APPENDIX F. ON PARTIAL HYPERBOLICITY

Here we state some facts about partial hyperbolicity that are used in the article but are well-known to the experts.

Recall that a C^1 -diffeomorphism $f: M \rightarrow M$ is partially hyperbolic if there exists a Df -invariant splitting $TM = E^s \oplus E^c \oplus E^u$ into 1-dimensional bundles and an $n > 0$ such that for every $x \in M$ we have

$$\|Df^n|_{E^s(x)}\| < \min\{1, \|Df^n|_{E^c(x)}\|\} \leq \max\{1, \|Df^n|_{E^c(x)}\|\} < \|Df^n|_{E^u(x)}\|.$$

By changing the Riemannian metric, one can always assume that $n = 1$ (see [Gou07, CP]).

A partially hyperbolic diffeomorphism is called *dynamically coherent* if there exists f -invariant foliations \mathcal{W}^{cs} and \mathcal{W}^{cu} tangent to $E^{cs} = E^s \oplus E^c$ and $E^{cu} = E^c \oplus E^u$. Taking the intersection of \mathcal{W}^{cs} and \mathcal{W}^{cu} gives a one-dimensional foliation \mathcal{W}^c tangent to E^c and f -invariant. Note that these foliations are not assumed to be unique in any sense (see [BW08] for a discussion).

Partially hyperbolic diffeomorphisms need not be dynamically coherent, but when they are, the standard notion of equivalence (which goes back to [HPS77]) is that of *leaf conjugacy*: Two dynamically coherent partially hyperbolic diffeomorphisms $f: M \rightarrow M$ and $g: N \rightarrow N$ are said to be *leaf conjugate* if there exists a homeomorphism $h: M \rightarrow N$ that maps the center foliation \mathcal{W}_f^c of f to the center foliation \mathcal{W}_g^c of g . More precisely, h is such that $h(\mathcal{W}_f^c(f(x))) = \mathcal{W}_g^c(g(h(x)))$ for all $x \in M$. We refer the reader to [Pot18] and references therein for more discussions.

We state the following result of Hertz, Hertz and Ures in a way that fits our particular needs.

Theorem F.1 ([RHRHU11]). *Let $f: M \rightarrow M$ be a partially hyperbolic diffeomorphism admitting a compact manifold⁸ tangent to E^{cs} (or E^{cu}). Then, M has solvable fundamental group (indeed, it is a torus bundle over the circle).*

In particular, if the fundamental group of M is not virtually solvable, and f is dynamically coherent, then the center-stable and center-unstable foliations are taut.

Hence, using the fundamental results of Burago and Ivanov [BI08] one gets:

Corollary F.2 ([BI08, Par10]). *Let M be a 3-manifold with non-solvable fundamental group. Suppose that M admits a partially hyperbolic diffeomorphism f such that the bundles E^s, E^c, E^u are orientable and Df preserves these orientations. Then M admits a taut foliation. In particular, M is irreducible, and aspherical.*

We recall that when $\pi_1(M)$ is (virtually) solvable a complete classification of partially hyperbolic diffeomorphisms is known [HP14, HP15, HP19].

⁸Notice that a compact manifold tangent to E^{cs} is necessarily a torus, see e.g., [RHRHU11].

In the setting of Seifert manifolds we used the following result from [HPS18] which is partially based on the study of horizontal and vertical laminations in Seifert manifolds [Bri93].

Theorem F.3 ([HPS18]). *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism on a Seifert manifold M whose fundamental group is not (virtually) solvable. Then, M is a finite cover of T^1S where S is a 2-dimensional hyperbolic orbifold, and the center-stable and center-unstable foliations of f are horizontal. That is, there exists a Seifert fibration $p: M \rightarrow \Sigma$ for which every leaf of the center-stable and center-unstable (2-dimensional) foliations is transverse to the (1-dimensional) fibers.*

An invariant foliation is called f -minimal if the only non-empty, saturated, closed set invariant by f is the whole manifold. The following result motivates asking for f -minimality of the foliations as a hypothesis as it covers the most important (from a dynamical standpoint) cases.

Proposition F.4 (see Lemma 1.1 of [BW05]). *Let $f: M \rightarrow M$ be a dynamically coherent partially hyperbolic diffeomorphism. If f is either volume preserving or transitive,⁹ then the center-stable and center-unstable foliations are f -minimal.*

Proof. Assume that there is a compact, non-empty f -invariant set Λ saturated by center-stable leaves. If $\Lambda \neq M, \emptyset$ then it must be a repeller, so f cannot be transitive nor volume preserving. \square

We remark that the property of f -minimality of \mathcal{W}^{cs} and \mathcal{W}^{cu} is a strictly weaker hypothesis than (chain-)transitivity (as seen, for instance, in the examples of [BG10]).

Finally, we recall the classification of partially hyperbolic diffeomorphisms in manifolds with virtually solvable fundamental group under the assumption that f is homotopic to the identity (see [HP14, HP15, HP19] for the general case):

Theorem F.5. *Let $f: M \rightarrow M$ be a partially hyperbolic diffeomorphism homotopic to identity in a 3-manifold with virtually solvable fundamental group. Then M is not Seifert fibered and if f is dynamically coherent then it is a discretized Anosov flow. Moreover, if there are no tori tangent to E^{cs} or E^{cu} then f is dynamically coherent.*

APPENDIX G. DISCRETIZED ANOSOV FLOWS

Let $\varphi_t: M \rightarrow M$ be a continuous flow generated by a continuous vector field $X = \frac{\partial \varphi_t}{\partial t}|_{t=0}$. It is called a *topological Anosov flow* if it preserves two topologically transverse codimension one continuous¹⁰ foliations \mathcal{F}^{ws} and \mathcal{F}^{wu} (called *weak stable* and *weak unstable*) such that:

- (i) For every pair of points $x, y \in \mathcal{F}^{ws}$ (resp. $x, y \in \mathcal{F}^{wu}$), there exists an increasing continuous reparametrization $h: \mathbb{R} \rightarrow \mathbb{R}$ so that $d(\varphi_t(x), \varphi_{h(t)}(y)) \rightarrow 0$ as $t \rightarrow +\infty$ (resp. as $t \rightarrow -\infty$);
- (ii) There exists $\epsilon > 0$ such that for every $x, y \in \mathcal{F}^{ws}$ (resp. $x, y \in \mathcal{F}^{wu}$) not on the same orbit, there exists $t \leq 0$ (resp. $t \geq 0$) such that $d(\varphi_t(x), \varphi_t(y)) > \epsilon$.

⁹It in fact suffices that f be *chain-recurrent*, that is, if a non-empty open set U verifies that $f(\overline{U}) \subset U$ then $U = M$, see [CP] for equivalences.

¹⁰We emphasize here that we do not require a priori the foliations to have C^1 -leaves.

As mentioned earlier in Appendix D, thanks to the work of Paternain [Pat93] and Inaba and Matsumoto [IM90], the definition of topological Anosov flow can be replaced by asking for the flow to be expansive and to preserve two (non singular, i.e., without prongs) foliations. Note also that just condition (i) is not enough for a flow to be topological Anosov as condition (i) does not imply condition (ii).

Conditions (i) and (ii) allow one to obtain the same classical results as for Anosov flows (e.g., there are no closed \mathcal{F}^s or \mathcal{F}^u leaves; the foliations \mathcal{F}^{ws} and \mathcal{F}^{wu} are taut; the leaves are planes, annuli or Möbius bands — these last two possibilities arising only when the leaves contain a periodic orbit; periodic points are dense in the non-wandering set, etc., see [Bar05] and references therein).

We say that a diffeomorphism $f: M \rightarrow M$ is a *discretized Anosov flow* if there exists a topological Anosov flow $\varphi_t: M \rightarrow M$ and a continuous function $\tau: M \rightarrow \mathbb{R}_{>0}$ such that $f(x) = \varphi_{\tau(x)}(x)$.

Remark G.1. Note that one should not expect a discretized Anosov flow to be the time-1 map of a reparametrization of the Anosov flow. To see this, one can for instance restrict to a periodic orbit of the flow: Then the discretized Anosov flow f restricts to a diffeomorphism of the circle. But diffeomorphisms of the circle do not all come from the time-1 map of a non-singular flow on the circle.

The following result relates the notion of discretized Anosov flows with the usual form of equivalence between partially hyperbolic systems.

Proposition G.2. *Let $f: M \rightarrow M$ be a partially hyperbolic diffeomorphism. The following are equivalent:*

- (1) f is a discretized Anosov flow;
- (2) f is dynamically coherent, the center leaves are fixed by f and the center foliation is the flow line foliation of a topological Anosov flow.

Proof. The fact that the second condition implies the first follows from arguments in [BW05], as was done in section 6.2.

The newer result is the other implication, which we now prove. Let $\varphi_t: M \rightarrow M$ be a topological Anosov flow and $\tau: M \rightarrow \mathbb{R}_{>0}$ be the positive continuous function such that $f(x) = \varphi_{\tau(x)}(x)$. Let F be the distribution generated by the vector field X generating φ_t . First, we claim that $F = E^c$. To prove this we will first show that F cannot be equal to E^s or E^u at any point and then deduce that F has to be E^c .

Suppose that there is $x \in M$ such that $F(x) = E^s(x)$. Then, the invariance of the orbits of the flow φ_t by f together with the uniqueness of the stable manifold implies that there is an arc $I = \varphi_{[-\epsilon, \epsilon]}(x)$ of the orbit of x by φ_t which is tangent to E^s . That fact is proven in [CP], we give here a brief explanation and the precise references. Consider a small cone field around E^s . Let α be the orbit of φ^t through x . Since $F(x) = E^s(x)$, it follows that, near $f^n(x)$, the curves $f^n(\alpha)$ are uniformly Lipschitz and their tangent are inside the cone field around E^s . Notice also that the family of curves $\{f^n(\alpha)\}$ is also invariant under f . The uniqueness of the stable manifold implies that Lipschitz curves that are invariant and inside the cone have to be the stable manifold near the point (this is done in [CP, Sections 4.2 and 4.3]). Hence $f^n(\alpha)$ must contain an open interval inside the stable manifold near x .

Iterating I backwards, we get that the length of $f^{-n}(I)$ grows exponentially, contradicting the continuity of $\tau: M \rightarrow \mathbb{R}_{>0}$.

Thus F is never tangent to E^s . The same argument shows that it is never tangent to E^u .

Now, suppose that there is a point y such that F is not inside E^{cs} at y . Then applying Df^n to $F(y)$ will get $F(f^n(y))$ closer and closer to $E^u(f^n(y))$. Hence, for any point z in the ω -limit set of y , one has that $F(z) = E^u(z)$, contradicting the above. So F is everywhere inside E^{cs} and, by the same argument, also inside E^{cu} . Thus $F = E^c$ everywhere.

The last step is to show that f is dynamically coherent, for this, we use the fact that the strong stable saturation of a center curve is tangent to $E^s \oplus E^c$ (see [BI08, Proposition 3.1]). We stress that a center curve here means a curve whose tangent everywhere is in the center bundle E^c . Since f is a discretized Anosov flow, the saturation of a flow line by strong stable leaves of f is contained in the weak stable foliation of the Anosov flow φ_t (cf. the proof of Proposition 6.5). In particular this implies that the weak stable foliation \mathcal{F}^{ws} of φ_t (which a priori could be only continuous) has C^1 leaves and is everywhere tangent to $E^s \oplus E^c$. This establishes dynamical coherence and completes the proof. \square

We remark that a long-standing conjecture (see [BW05]) states that every topological Anosov flow is orbit equivalent to an Anosov flow. If this conjecture is true, then condition (2) above is equivalent to saying that f is dynamically coherent and *leaf conjugate* to the time one map of an Anosov flow. We remark here that it has been recently announced that this conjecture is true in the setting of transitive topological Anosov flows [Sha].

APPENDIX H. THE GRAPH TRANSFORM ARGUMENT

We give here an application of the general graph transform technique to the particular case we needed it in.

We call *center stable plane* any embedded C^1 -plane tangent to $E^s \oplus E^c$ in \widetilde{M} . Notice that by unique integrability of E^s there is always a stable foliation inside a center stable plane.

Lemma H.1 (Graph Transform Lemma). *Let f be a partially hyperbolic diffeomorphism in M . Suppose that $L \subset \widetilde{M}$ is a center stable plane which is fixed by a lift \hat{f} of f to \widetilde{M} , and by some $\gamma \in \pi_1(M) \setminus \{id\}$. Assume that there is a properly embedded C^1 curve η transverse to the stable foliation in L and such that*

$$\gamma\eta = \eta \quad \text{and} \quad \hat{f}(\eta) \subset \bigcup_{z \in \eta} \widetilde{\mathcal{W}}^s(z).$$

Then in L there is a curve $\hat{\eta}$ which is fixed by both \hat{f} and γ and is everywhere tangent to E^c .

Notice the subtlety in the conclusion of this lemma: The curve $\hat{\eta}$ produced is tangent to the center direction, however, it may *not* be a center leaf since the bundle may not be uniquely integrable.

Remark H.2. The second hypothesis of the lemma is equivalent to saying that the union $\bigcup_{z \in \eta} \widetilde{\mathcal{W}}^s(z)$ is invariant by \hat{f} . In particular, all positive and negative images of η by powers of \hat{f} are contained in this union. To see this, notice that the second condition implies that $f_1(\alpha)$ is freely homotopic to α in $L/\langle \gamma \rangle$ because it is a cylinder, and α (or $f_1(\alpha)$) is not null homotopic in this cylinder. Therefore $\bigcup_{z \in \eta} \widetilde{\mathcal{W}}^s(z) = \bigcup_{z \in \hat{f}(\eta)} \widetilde{\mathcal{W}}^s(z)$. Thus $\bigcup_{z \in \eta} \widetilde{\mathcal{W}}^s(z)$ is \hat{f} invariant. The converse is immediate.

Proof. We work in the quotient $L/\langle\gamma\rangle$ which is an annulus on which η projects to a closed C^1 -circle transverse to the stable foliation, denoted by α . Let $\pi_0: L \rightarrow L/\langle\gamma\rangle$ the quotient map. Let f_1 be the induced diffeomorphism on $L/\langle\gamma\rangle$.

Up to a small modification of α if necessary, we can assume that α is simple, that is, it goes around the cylinder $L/\langle\gamma\rangle$ once.

We parametrize α in $L/\langle\gamma\rangle$ by arclength (for the leaf-wise path metric on $L/\langle\gamma\rangle$). Then, we parametrize $\bigcup_{z \in \alpha} \mathcal{W}^s(z)$ as a cylinder $S^1 \times \mathbb{R}$ contained in $L/\langle\gamma\rangle$, where α is the zero section and the stable leaves are parametrized by arclength.

Since all $\hat{f}^n(\eta)$ are in $\bigcup_{z \in \eta} \mathcal{W}^s(z)$, we can express $\eta_n := \hat{f}^n(\eta)$ as graphs of C^1 -functions¹¹ from S^1 to \mathbb{R} .

We want to show that \hat{f} acts as a contraction on curves transverse to the stable foliation in (at least a compact part of) $\bigcup_{z \in \alpha} \mathcal{W}^s(z) \simeq S^1 \times \mathbb{R}$. First we show that all the η_n stay in a compact subset of $\bigcup_{z \in \eta} \mathcal{W}^s(z)$ (and thus also of $L/\langle\gamma\rangle$).

Our assumptions imply that there exists some $a_0 > 0$, such that $f_1(\alpha)$ is contained in an a_0 stable neighborhood of η . That is, in the union of stable segment of length $2a_0$ centered at η .

Let $\lambda < 1$ be the smallest contraction factor for f_1 along stable leaves. It follows that $f_1^2(\eta)$ is contained in the stable neighborhood of size $a_0 + \lambda a_0$ around α , and so on. Thus, we immediately get that, for all n , $f_1^n(\alpha)$ is contained in a compact subset of the annulus.

Now that we know that all the η_n curves are contained in a compact subset, we can use the fact that f_1 contracts stable leaves more than centers to prove the following:

There exists some constant a_1 such that f_1 globally preserves the space of uniformly bounded (for some appropriately large bound) Lipschitz functions from S^1 to \mathbb{R} with Lipschitz constant less than a_1 . By standard computations, one can see that this acts as a contraction on this complete metric space (this is usually called the *graph transform technique* see e.g., [HPS77] or [CP, Section 4.2] for a more detailed study of this technique and the reason for considering Lipschitz functions).

Therefore, one obtains that there is a unique fixed point of this action which corresponds to the graph of a Lipschitz function from S^1 to \mathbb{R} which is the unique invariant Lipschitz graph under f_1 . It is also standard to show that the tangent cones at each point must actually be degenerate (see [CP, Section 4.2]), i.e., the invariant curve is C^1 . Moreover, since E^c is the only invariant bundle transverse to E^s , the curve must be everywhere tangent to E^c . The lift of this curve to L is the curve we sought. \square

Under the assumptions of the Graph Transform Lemma (Lemma H.1), another thing we easily deduce is that there must be a periodic center leaf of f in the projection of the leaf L :

Lemma H.3. *Let f be a partially hyperbolic diffeomorphism in M . Suppose that $L \subset \widetilde{M}$ is a center stable plane which is fixed by a lift \hat{f} of f to \widetilde{M} , and by some $\gamma \in \pi_1(M) \setminus \{id\}$.*

Assume that there exists a curve $\hat{\eta}$ that is fixed by both \hat{f} and γ . Then there exists a center leaf c in L and two integers n, m , with $m \neq 0$, such that $c = \gamma^n \hat{f}^m c$.

¹¹In this specific case with one dimensional center, one can assume that the stable foliation is C^1 inside center stable leaves so that this makes sense, see [CP, Section 4.7]. If the stable foliation is less regular then one can go through with the proof by taking a smooth approximating foliation instead, and the arguments would be essentially the same.

To prove this lemma, we need to use the center leaf space on L . When the foliations are branching (i.e. in the non-dynamically coherent setting), the center leaf space will be defined in [BFFPa].

Proof. Let $\eta = \pi(\hat{\eta})$ be the projection of $\hat{\eta}$ to M . Since $\hat{\eta}$ is invariant by γ and \hat{f} , the curve η is a circle on which f acts.

Suppose first that f has a periodic point on η . Then, there exists a center leaf through that point that is periodic, as claimed.

Otherwise, there exists a point in η that is inside its ω -limit set for f .

Lifting back to the universal cover, this means that there exists $x \in \hat{\eta}$ such that there exists integers m, n , with m arbitrarily large, such that x and $\gamma^n \hat{f}^m(x)$ can be made arbitrarily close.

Let τ be a compact segment of the stable leaf through x . Since \hat{f} contracts the length of stable segments, we can choose $m \in \mathbb{N}$ large enough so that every center leaf through $\gamma^n \hat{f}^m(\tau)$ intersects the interior of τ . (This is possible as $\gamma^n \hat{f}^m(\tau)$ can be chosen arbitrarily small and arbitrarily close to x , which is in the interior of τ .)

Let \mathcal{L}_L^c be the leaf space of the center foliation in L . Let

$$\tau_c = \{c \in \mathcal{L}_L^c \mid c \cap \tau \neq \emptyset\}.$$

Notice that τ_c is a compact interval in the 1-manifold \mathcal{L}_L^c .

Then consider the function $h: \tau_c \rightarrow \mathcal{L}_L^c$ defined by $h(c) = \gamma^n \hat{f}^m(c)$. The map h is continuous, and, thanks to our choice of m , $h(\tau_c)$ is contained in the interior of τ_c . Hence, there exists $c_0 \in \tau_c$ that is fixed by h , as claimed. \square

REFERENCES

- [AVW15] A. Avila, M. Viana, and A. Wilkinson, *Absolute continuity, Lyapunov exponents and rigidity I: geodesic flows*, J. Eur. Math. Soc. (JEMS) **17** (2015), no. 6, 1435–1462. MR 3353805
- [Bar95] T. Barbot, *Caractérisation des flots d’Anosov en dimension 3 par leurs feuilletages faibles*, Ergodic Theory Dynam. Systems **15** (1995), no. 2, 247–270.
- [Bar96] ———, *Flots d’Anosov sur les variétés graphées au sens de Waldhausen*, Ann. Inst. Fourier (Grenoble) **46** (1996), no. 5, 1451–1517.
- [Bar98] ———, *Actions de groupes sur les 1-variétés non séparées et feuilletages de codimension un*, Ann. Fac. Sci. Toulouse Math. (6) **7** (1998), no. 4, 559–597. MR 1693597
- [Bar05] ———, *De l’hyperbolique au globalement hyperbolique*, Habilitation à diriger des recherches, Université Claude Bernard de Lyon, 2005.
- [BBGR17] T. Barthelmé, C. Bonatti, A. Gogolev, and F. Rodriguez Hertz, *Anomalous Anosov flows revisited*, Proc. Lond. Math. Soc. (2017), arXiv:1712.07755.
- [BFFPa] T. Barthelmé, S. Fenley, S. Frankel, and R. Potrie, *Partially hyperbolic diffeomorphisms homotopic to identity in dimension three II: branching foliations*, arXiv.
- [BFFPb] ———, *Research Announcement: Partially hyperbolic diffeomorphisms homotopic to the identity on 3-manifolds*, 2018 MATRIX Annals.
- [BFFP20] ———, *Dynamical incoherence for a large class of partially hyperbolic diffeomorphisms*, ArXiv e-prints (2020).
- [BDV05] C. Bonatti, L. J. Díaz, and M. Viana, *Dynamics beyond uniform hyperbolicity*, Encyclopaedia of Mathematical Sciences, vol. 102, Springer-Verlag, Berlin, 2005, A global geometric and probabilistic perspective, Mathematical Physics, III. MR 2105774
- [BGHP17] C. Bonatti, A. Gogolev, A. Hammerlindl, and R. Potrie, *Anomalous partially hyperbolic diffeomorphisms III: abundance and incoherence*, arXiv e-prints (2017), arXiv:1706.04962.
- [BGP16] C. Bonatti, A. Gogolev, and R. Potrie, *Anomalous partially hyperbolic diffeomorphisms II: stably ergodic examples*, Invent. Math. **206** (2016), no. 3, 801–836. MR 3573973

- [BG10] C. Bonatti and N. Guelman, *Axiom A diffeomorphisms derived from Anosov flows*, J. Mod. Dyn. **4** (2010), no. 1, 1–63. MR 2643887
- [BPP16] C. Bonatti, K. Parwani, and R. Potrie, *Anomalous partially hyperbolic diffeomorphisms I: Dynamically coherent examples*, Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), no. 6, 1387–1402. MR 3592360
- [BW05] C. Bonatti and A. Wilkinson, *Transitive partially hyperbolic diffeomorphisms on 3-manifolds*, Topology **44** (2005), no. 3, 475–508. MR 2122214
- [BZ17] C. Bonatti and J. Zhang, *Transverse foliations on the torus \mathbb{T}^2 and partially hyperbolic diffeomorphisms on 3-manifolds*, Comment. Math. Helv. **92** (2017), no. 3, 513–550. MR 3682779
- [BZ19] C. Bonatti and J. Zhang, *Transitive partially hyperbolic diffeomorphisms with one-dimensional neutral center*, arXiv e-prints (2019), arXiv:1904.05295.
- [BBI04] M. Brin, D. Burago, and S. Ivanov, *On partially hyperbolic diffeomorphisms of 3-manifolds with commutative fundamental group*, Modern dynamical systems and applications, Cambridge Univ. Press, Cambridge, 2004, pp. 307–312. MR 2090777
- [BP73] M. I. Brin and Ja. B. Pesin, *Partially hyperbolic dynamical systems*, Uspehi Mat. Nauk **28** (1973), no. 3(171), 169–170. MR 0391178
- [Bri93] M. Brittenham, *Essential laminations in Seifert-fibered spaces*, Topology **32** (1993), no. 1, 61–85. MR 1204407
- [BI08] D. Burago and S. Ivanov, *Partially hyperbolic diffeomorphisms of 3-manifolds with abelian fundamental groups*, J. Mod. Dyn. **2** (2008), no. 4, 541–580. MR 2449138
- [BW08] K. Burns and A. Wilkinson, *Dynamical coherence and center bunching*, Discrete Contin. Dyn. Syst. **22** (2008), no. 1-2, 89–100. MR 2410949
- [BFT19] J. Buzzi, T. Fisher, and A. Tahzibi, *A dichotomy for measures of maximal entropy near time-one maps of transitive Anosov flows*, arXiv e-prints (2019), arXiv:1904.07821.
- [Cal00] D. Calegari, *The geometry of \mathbf{R} -covered foliations*, Geom. Topol. **4** (2000), 457–515 (electronic).
- [Cal07] ———, *Foliations and the geometry of 3-manifolds*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2007.
- [CLN85] C. Camacho and A. Lins Neto, *Geometric theory of foliations*, Birkhäuser Boston, Inc., Boston, MA, 1985, Translated from the Portuguese by Sue E. Goodman. MR 824240
- [Can93] A. Candel, *Uniformization of surface laminations*, Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 4, 489–516.
- [CC00] A. Candel and L. Conlon, *Foliations. I*, Graduate Studies in Mathematics, vol. 23, American Mathematical Society, Providence, RI, 2000. MR 1732868
- [CC03] ———, *Foliations. II*, Graduate Studies in Mathematics, vol. 60, American Mathematical Society, Providence, RI, 2003. MR 1994394
- [CRRU15] P. Carrasco, F. Rodriguez Hertz, J. Rodriguez Hertz, and R. Ures, *Partially hyperbolic dynamics in dimension 3*, ArXiv e-prints (2015).
- [CPR19] P. D. Carrasco, E. Pujals, and F. Rodriguez-Hertz, *Classification of partially hyperbolic diffeomorphisms under some rigid conditions*, arXiv e-prints (2019), arXiv:1903.09264.
- [CJ94] A. Casson and D. Jungreis, *Convergence groups and Seifert fibered 3-manifolds*, Invent. Math. **118** (1994), no. 3, 441–456. MR 1296353
- [CP] S. Crovisier and R. Potrie, *Introduction to partially hyperbolic dynamics*, available from the authors webpage.
- [DPU99] L. J. Díaz, E. R. Pujals, and R. Ures, *Partial hyperbolicity and robust transitivity*, Acta Math. **183** (1999), no. 1, 1–43. MR 1719547
- [Eps72] D. B. A. Epstein, *Periodic flows on three-manifolds*, Ann. of Math. (2) **95** (1972), 66–82. MR 0288785
- [FP18] S. Fenley and R. Potrie, *Ergodicity of partially hyperbolic diffeomorphisms in hyperbolic 3-manifolds*, arXiv e-prints (2018), arXiv:1809.02284.
- [Fen94] S. R. Fenley, *Anosov flows in 3-manifolds*, Ann. of Math. (2) **139** (1994), no. 1, 79–115.
- [Fen98] ———, *The structure of branching in Anosov flows of 3-manifolds*, Comment. Math. Helv. **73** (1998), no. 2, 259–297.
- [Fen02] ———, *Foliations, topology and geometry of 3-manifolds: \mathbf{R} -covered foliations and transverse pseudo-Anosov flows*, Comment. Math. Helv. **77** (2002), no. 3, 415–490.

- [Fen03] ———, *Pseudo-Anosov flows and incompressible tori*, *Geom. Dedicata* **99** (2003), 61–102. MR 1998929
- [Fen13] ———, *Rigidity of pseudo-Anosov flows transverse to \mathbb{R} -covered foliations*, *Comment. Math. Helv.* **88** (2013), no. 3, 643–676. MR 3093506
- [FW80] J. Franks and B. Williams, *Anomalous Anosov flows*, *Global theory of dynamical systems* (Proc. Internat. Conf., Northwestern Univ., Evanston, Ill., 1979), *Lecture Notes in Math.*, vol. 819, Springer, Berlin, 1980, pp. 158–174.
- [Gab92] D. Gabai, *Convergence groups are Fuchsian groups*, *Ann. of Math. (2)* **136** (1992), no. 3, 447–510. MR 1189862
- [Ghy84] É. Ghys, *Flots d’Anosov sur les 3-variétés fibrées en cercles*, *Ergodic Theory Dynam. Systems* **4** (1984), no. 1, 67–80.
- [Gou07] N. Gourmelon, *Adapted metrics for dominated splittings*, *Ergodic Theory Dynam. Systems* **27** (2007), no. 6, 1839–1849. MR 2371598
- [HP14] A. Hammerlindl and R. Potrie, *Pointwise partial hyperbolicity in three-dimensional nilmanifolds*, *J. Lond. Math. Soc. (2)* **89** (2014), no. 3, 853–875. MR 3217653
- [HP15] ———, *Classification of partially hyperbolic diffeomorphisms in 3-manifolds with solvable fundamental group*, *J. Topol.* **8** (2015), no. 3, 842–870. MR 3394318
- [HP18] ———, *Partial hyperbolicity and classification: a survey*, *Ergodic Theory Dynam. Systems* **38** (2018), no. 2, 401–443. MR 3774827
- [HP19] ———, *Classification of systems with center-stable tori*, *Michigan Math. J.* **68** (2019), no. 1, 147–166. MR 3934607
- [HPS18] A. Hammerlindl, R. Potrie, and M. Shannon, *Seifert manifolds admitting partially hyperbolic diffeomorphisms*, *J. Mod. Dyn.* **12** (2018), 193–222. MR 3824728
- [Han85] M. Handel, *Global shadowing of pseudo-Anosov homeomorphisms*, *Ergodic Theory Dynam. Systems* **5** (1985), no. 3, 373–377. MR 805836
- [HP06] B. Hasselblatt and Y. Pesin, *Partially hyperbolic dynamical systems*, *Handbook of dynamical systems*. Vol. 1B, Elsevier B. V., Amsterdam, 2006, pp. 1–55. MR 2186241
- [Hat] A. Hatcher, *Notes on basic 3-manifold topology*.
- [Hem76] J. Hempel, *3-Manifolds*, Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1976, *Ann. of Math. Studies*, No. 86. MR 0415619
- [HPS77] M. W. Hirsch, C. C. Pugh, and M. Shub, *Invariant manifolds*, *Lecture Notes in Mathematics*, Vol. 583, Springer-Verlag, Berlin-New York, 1977. MR 0501173
- [IM90] T. Inaba and S. Matsumoto, *Nonsingular expansive flows on 3-manifolds and foliations with circle prong singularities*, *Japan. J. Math. (N.S.)* **16** (1990), no. 2, 329–340. MR 1091166
- [Jac80] W. Jaco, *Lectures on three-manifold topology*, *CBMS Regional Conference Series in Mathematics*, vol. 43, American Mathematical Society, Providence, R.I., 1980. MR 565450
- [Man18] K. Mann, *Rigidity and flexibility of group actions on the circle*, *Handbook of group actions*. Vol. IV, *Adv. Lect. Math. (ALM)*, vol. 41, Int. Press, Somerville, MA, 2018, pp. 705–752. MR 3888699
- [Mos68] G. D. Mostow, *Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms*, *Inst. Hautes Études Sci. Publ. Math.* (1968), no. 34, 53–104. MR 0236383
- [Nav11] A. Navas, *Groups of circle diffeomorphisms*, spanish ed., *Chicago Lectures in Mathematics*, University of Chicago Press, Chicago, IL, 2011. MR 2809110
- [Nov65] S. P. Novikov, *The topology of foliations*, *Trudy Moskov. Mat. Obšč.* **14** (1965), 248–278. MR 0200938
- [Pal78] C. F. B. Palmeira, *Open manifolds foliated by planes*, *Ann. Math. (2)* **107** (1978), no. 1, 109–131.
- [Par10] K. Parwani, *On 3-manifolds that support partially hyperbolic diffeomorphisms*, *Nonlinearity* **23** (2010), no. 3, 589–606. MR 2586372
- [Pat93] M. Paternain, *Expansive flows and the fundamental group*, *Bol. Soc. Brasil. Mat. (N.S.)* **24** (1993), no. 2, 179–199.
- [Per02] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, *ArXiv Mathematics e-prints* (2002).
- [Per03a] ———, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, *ArXiv Mathematics e-prints* (2003).

- [Per03b] ———, *Ricci flow with surgery on three-manifolds*, ArXiv Mathematics e-prints (2003).
- [Pot18] R. Potrie, *Robust dynamics, invariant structures and topological classification*, arXiv e-prints (2018), arXiv:1802.05291.
- [PS04] Charles Pugh and Michael Shub, *Stable ergodicity*, Bull. Amer. Math. Soc. (N.S.) **41** (2004), no. 1, 1–41, With an appendix by Alexander Starkov. MR 2015448
- [RHRHU11] F. Rodriguez Hertz, M. A. Rodriguez Hertz, and R. Ures, *Tori with hyperbolic dynamics in 3-manifolds*, J. Mod. Dyn. **5** (2011), no. 1, 185–202. MR 2787601
- [RHRHU16] ———, *A non-dynamically coherent example on \mathbb{T}^3* , Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), no. 4, 1023–1032. MR 3519530
- [Ros68] H. Rosenberg, *Foliations by planes*, Topology **7** (1968), 131–138. MR 0228011 (37 #3595)
- [Sco83] P. Scott, *The geometries of 3-manifolds*, Bull. London Math. Soc. **15** (1983), no. 5, 401–487. MR 705527
- [Sha] M. Shannon, *Personal communication*.
- [Thu] W. P. Thurston, *3-manifolds, foliations and circles I*.
- [Tol78] J. L. Tollefson, *Involutions of Seifert fiber spaces*, Pacific J. Math. **74** (1978), no. 2, 519–529. MR 0645400
- [Wil10] A. Wilkinson, *Conservative partially hyperbolic dynamics*, Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi, 2010, pp. 1816–1836. MR 2827868
- [Zha17] J. Zhang, *Partially hyperbolic diffeomorphisms with one-dimensional neutral center on 3-manifolds*, arXiv e-prints (2017), arXiv:1701.06176.

QUEEN'S UNIVERSITY, KINGSTON, ON

Email address: `thomas.barthelme@queensu.edu`

URL: `sites.google.com/site/thomasbarthelme`

FLORIDA STATE UNIVERSITY, TALLAHASSEE, FL 32306

Email address: `fenley@math.fsu.edu`

WASHINGTON UNIVERSITY IN ST. LOUIS, ST. LOUIS, MO

Email address: `steven.frankel@wustl.edu`

CENTRO DE MATEMÁTICA, UNIVERSIDAD DE LA REPÚBLICA, URUGUAY

Email address: `rpotrie@cmat.edu.uy`

URL: `http://www.cmat.edu.uy/~rpotrie/`