

# Length 3 Complexes of Abelian Sheaves and Picard 2-Stacks

A. Emin Tatar

Department of Mathematics, Florida State University

Tallahassee, FL 32306-4510, USA

atatar@math.fsu.edu

## Abstract

We define a tricategory  $T^{[-2,0]}$  of length 3 complexes of abelian sheaves, whose hom-bigroupoids consist of weak morphisms of such complexes. We also define a 3-category  $2PIC(S)$  of Picard 2-stacks, whose hom-2-groupoids consist of additive 2-functors. We prove that these categories are triequivalent as tricategories. As a consequence we obtain a generalization of Deligne's analogous result about Picard stacks in SGA4, Exposé XVIII.

## Contents

<b>Introduction</b>	<b>2</b>
<b>1 Preliminaries</b>	<b>4</b>
1.1 Butterflies . . . . .	4
1.2 $(A, B)$ -torsors . . . . .	5
1.3 $(\mathcal{A}, \mathcal{B})$ -torsors . . . . .	5
1.4 Abelian Sheaves and Picard Stacks . . . . .	6
1.5 Tricategories . . . . .	7
<b>2 Picard 2-Stacks as Torsors</b>	<b>7</b>
2.1 2-Stacks . . . . .	7
2.2 Picard 2-Stack Associated to a Complex . . . . .	12
2.3 Homotopy Exact Sequence . . . . .	13
2.4 The 3-category of Picard 2-Stacks . . . . .	14
<b>3 The 3-category of Complexes of Abelian Sheaves</b>	<b>14</b>
3.1 Definition of $C^{[-2,0]}(S)$ . . . . .	14
3.2 Abelian Sheaves and Picard 2-Stacks . . . . .	15
<b>4 Weak Morphisms of Complexes of Abelian Sheaves</b>	<b>17</b>
4.1 Definition of $Frac(A^\bullet, B^\bullet)$ . . . . .	17
4.2 $Frac(A^\bullet, B^\bullet)$ is a bigroupoid . . . . .	19

<b>5</b>	<b>Biequivalence of <math>\text{Frac}(A^\bullet, B^\bullet)</math> and <math>\text{Hom}(A^\bullet, B^\bullet)</math></b>	<b>21</b>
5.1	Morphisms of Picard 2-Stacks as Fractions . . . . .	21
5.2	Hom-categories of $\text{Frac}(A^\bullet, B^\bullet)$ and $\text{Hom}(A^\bullet, B^\bullet)$ . . . . .	25
<b>6</b>	<b>The Tricategory of Complexes of Abelian Sheaves</b>	<b>30</b>
6.1	Definition of $\mathcal{T}^{[-2,0]}(\mathcal{S})$ . . . . .	30
6.2	Main Theorem . . . . .	33
<b>7</b>	<b>Stackification</b>	<b>34</b>

## Introduction

Let  $\mathcal{D}^{[-1,0]}(\mathcal{S})$  be the subcategory of the derived category of category of complexes of abelian sheaves  $A^\bullet$  over a site  $\mathcal{S}$  with  $H^{-i}(A^\bullet) \neq 0$  only for  $i = 0, 1$ . Let  $\text{Pic}^b(\mathcal{S})$  denote the category of Picard stacks over  $\mathcal{S}$  with 1-morphisms isomorphism classes of additive functors. In SGA4 Exposé XVIII, Deligne shows the following.

**Proposition.** [9, Proposition 1.4.15] The functor

$$\wp^b : \mathcal{D}^{[-1,0]}(\mathcal{S}) \longrightarrow \text{Pic}^b(\mathcal{S})$$

given by sending a length 2 complex of abelian sheaves,  $A^\bullet : A^{-1} \rightarrow A^0$  over  $\mathcal{S}$  to its associated Picard stack  $[A^{-1} \rightarrow A^0]^\sim$ , an isomorphism class of fractions from  $A^\bullet$  to  $B^\bullet$  to an isomorphism class of morphisms of associated Picard stacks is an equivalence.

The purpose of this paper is to generalize the above result to Picard 2-stacks over  $\mathcal{S}$ . Let  $2\text{Pic}^{bb}(\mathcal{S})$  denote the category of Picard 2-stacks, whose morphisms are equivalence classes of additive 2-functors. Let  $\mathcal{D}^{[-2,0]}(\mathcal{S})$  be the subcategory of the derived category of category of complexes of abelian sheaves  $A^\bullet$  over  $\mathcal{S}$  with  $H^{-i}(A^\bullet) \neq 0$  for  $i = 0, 1, 2$ .

**Theorem I.** The functor

$$2\wp^{bb} : \mathcal{D}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{Pic}^{bb}(\mathcal{S})$$

given by sending a length 3 complex of abelian sheaves,  $A^\bullet : A^{-2} \rightarrow A^{-1} \rightarrow A^0$  over  $\mathcal{S}$  to its associated Picard 2-stack  $[A^{-2} \rightarrow A^{-1} \rightarrow A^0]^\sim$ , an equivalence class of fractions from  $A^\bullet$  to  $B^\bullet$  to an equivalence class of morphisms of associated Picard 2-stacks is an equivalence.

Basically, it gives a geometric description of the derived category of length 3 complexes of abelian sheaves. It states that any Picard 2-stack over a site  $\mathcal{S}$  is biequivalent to a Picard 2-stack associated to a length 3 complex of abelian sheaves and that any morphism of Picard 2-stacks comes from a fraction of such complexes. A complex of abelian sheaves, whose only non-zero cohomology groups are placed at degrees -2, -1, and 0 can be thought as a length 3 complex of abelian sheaves, and therefore a morphism in  $\mathcal{D}^{[-2,0]}(\mathcal{S})$  between any two complexes  $A^\bullet$  and  $B^\bullet$  is given by an equivalence class of fraction

$$(q, M^\bullet, p) : A^\bullet \xleftarrow{q} M^\bullet \xrightarrow{p} B^\bullet$$

with  $q$  being a quasi-isomorphism.

However, we prove a much stronger statement, so that the latter theorem becomes an immediate consequence of it. Let  $2\text{PIC}(\mathcal{S})$  be the 3-category of Picard 2-stacks where 1-morphisms are additive 2-functors, 2-morphisms are natural 2-transformations, and 3-morphisms are modifications. Length 3 complexes of abelian sheaves over  $\mathcal{S}$  placed in degrees  $[-2, 0]$  form a 3-category  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  by adding to the regular morphisms of complexes, the degree -1 and -2 morphisms. Then we easily construct an explicit trihomomorphism

$$2\wp : \mathcal{C}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{PIC}(\mathcal{S}),$$

that is a 3-functor between 3-categories. Under this construction, length 3 complexes of abelian sheaves correspond to Picard 2-stacks. Although morphisms of such complexes induce morphisms between associated Picard 2-stacks, not all of them are obtained in this way. In this sense, the 1-morphisms of  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  are not geometric and the reason is their strictness. We resolve this problem by weakening  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  as follows: We introduce a tricategory  $\mathcal{T}^{[-2,0]}(\mathcal{S})$  (a tricategory is a weak version of a 3-category in the sense of [11]) with same objects as  $\mathcal{C}^{[-2,0]}(\mathcal{S})$ . For any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , morphisms between  $A^\bullet$  and  $B^\bullet$  in  $\mathcal{T}^{[-2,0]}(\mathcal{S})$  is the bigroupoid  $\text{Frac}(A^\bullet, B^\bullet)$ , whose main property is that it satisfies  $\pi_0(\text{Frac}(A^\bullet, B^\bullet)) \simeq \text{Hom}_{\mathcal{D}^{[-2,0]}(\mathcal{S})}(A^\bullet, B^\bullet)$ , where  $\pi_0$  denotes the isomorphism classes of objects. Roughly speaking, objects of  $\text{Frac}(A^\bullet, B^\bullet)$  are fractions from  $A^\bullet$  to  $B^\bullet$  in the ordinary sense and its 2-morphisms are certain commutative diagrams (4.2) called “*diamonds*”. Then we prove:

**Theorem II.** The trihomomorphism

$$2\wp : \mathcal{T}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{PIC}(\mathcal{S})$$

defined by sending  $A^\bullet$  a length 3 complex of abelian sheaves to its associated Picard 2-stack is a triequivalence.

Since in particular a triequivalence is essentially surjective, every Picard 2-stack is biequivalent to a Picard 2-stack associated to a complex of abelian sheaves. Then by ignoring the 3-morphisms and passing to the equivalence class of morphisms in the triequivalence of Theorem II, we deduce Theorem I.

## Organization of the paper

This paper is organized as follows:

In Section 1, we recall important points of butterflies in abelian context,  $(A, B)$ -torsors, where  $A$  and  $B$  are abelian sheaves, and  $(\mathcal{A}, \mathcal{B})$ -torsors, where  $\mathcal{A}$  and  $\mathcal{B}$  are Picard stacks. We also remind the reader some important results from [3] that we will refer continuously.

In Section 2, we explain briefly the basics on 2-stacks with structures, and exact sequences of Picard 2-stacks. We also give an example of Picard 2-stack, namely  $\text{TORS}(\mathcal{A}, A^0)$ , where  $\mathcal{A}$  is a Picard stack and  $A^0$  is an abelian sheaf. This example will be of great importance for the rest since it will be the Picard 2-stack associated to  $A^\bullet$  a length 3 complex of abelian sheaves. We define at the end of the section the 3-category  $2\text{PIC}(\mathcal{S})$  of Picard 2-stacks, as well.

In Section 3, we first define the 3-category  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  of length 3 complexes of abelian sheaves. We construct an explicit trihomomorphism from this 3-category to the 3-category of Picard 2-stacks.

In Section 4, for any two length 3 complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , we define a bigroupoid  $\text{Frac}(A^\bullet, B^\bullet)$ . It is a weakened version of the hom-2-category  $\text{Hom}_{\mathcal{C}^{[-2,0]}(\mathcal{S})}(A^\bullet, B^\bullet)$  in the sense that  $\pi_0(\text{Frac}(A^\bullet, B^\bullet)) \simeq \text{Hom}_{\mathcal{D}^{[-2,0]}(\mathcal{S})}(A^\bullet, B^\bullet)$ .

In Section 5, we show that for any two length 3 complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , there exists a biequivalence as bigroupoids between  $\text{Frac}(A^\bullet, B^\bullet)$  and the 2-category  $\text{Hom}(A^\bullet, B^\bullet)$  of morphisms of Picard 2-stacks associated to  $A^\bullet$  and  $B^\bullet$ .

In Section 6, we define the tricategory  $\mathcal{T}^{[-2,0]}(\mathcal{S})$ . It consists of same objects as  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  and for any two length 3 complexes  $A^\bullet$  and  $B^\bullet$  of abelian sheaves,  $\text{Frac}(A^\bullet, B^\bullet)$  as the hom-bigroupoid. We extend the trihomomorphism constructed in Section 3 to a trihomomorphism on  $\mathcal{T}^{[-2,0]}(\mathcal{S})$ . We prove Theorem II which says that the latter trihomomorphism is a triequivalence and from which Theorem I follows.

In Section 7, we informally discuss the stack versions of what has been done in the previous sections.

## Acknowledgements

I would like to express my profound gratitude to my advisor, Ettore Aldrovandi, for helping me at all stages of the paper, which is part of my Ph.D. thesis. I would like to thank Behrang Noohi for helpful conversations. I also thank Chris Portwood for proofreading.

## 1 Preliminaries

The method that we are going to adopt to prove our results is going to use mostly the language and techniques developed in the papers of Aldrovandi and Noohi such as butterflies, torsors, etc. So it is worthwhile to mention here some of their work. We finish with a few words about bicategories and tricategories. Before, let us fix our conventions and notations.

Throughout the paper, we will work with sheaves, stacks, etc. defined over a site  $\mathcal{S}$ . For simplicity, we will assume that  $\mathcal{S}$  has fibered products. Fibered 2-categories, 2-functors, and natural 2-transformations will be used in the sense of Hakim [13]. A complex of abelian sheaves will mean a length 3 complex of abelian sheaves over the site  $\mathcal{S}$  unless otherwise stated. It will be denoted as

$$A^\bullet : A^{-2} \xrightarrow{\delta_A} A^{-1} \xrightarrow{\lambda_A} A^0.$$

For any complex of abelian sheaves  $A^\bullet$ ,  $A^{\bullet < 0}$  denotes the complex

$$A^{\bullet < 0} : A^{-2} \xrightarrow{\delta_A} A^{-1} \longrightarrow 0$$

and therefore  $f^{\bullet < 0} : A^{\bullet < 0} \rightarrow B^{\bullet < 0}$  a morphism of complexes between  $A^{\bullet < 0}$  and  $B^{\bullet < 0}$ .

### 1.1 Butterflies

The reader can refer to [17] and [18] for details of butterflies over a point or to [3] for a treatment over a site. Here, we will remind the basic definitions following the latter point of view in an abelian context.

**Definition 1.1.** Let  $A^\bullet : A^{-1} \rightarrow A^0$  and  $B^\bullet : B^{-1} \rightarrow B^0$  be two length 2 complexes of abelian sheaves. A butterfly from  $A^\bullet$  to  $B^\bullet$  is a commutative diagram of abelian sheaf morphisms of the form

$$\begin{array}{ccc}
 A^{-1} & & B^{-1} \\
 \downarrow & \searrow \kappa & \swarrow \iota \\
 & E & \\
 \downarrow & \swarrow \rho & \searrow j \\
 A^0 & & B^0
 \end{array}
 \tag{1.1}$$

where  $E$  is an abelian sheaf, the NW-SE sequence is a complex, and the NE-SW sequence is an extension.  $[A^\bullet, E, B^\bullet]$  will denote a butterfly from  $A^\bullet$  to  $B^\bullet$ .

A morphism of butterflies  $\varphi : [A^\bullet, E, B^\bullet] \rightarrow [A^\bullet, E', B^\bullet]$  is an abelian sheaf isomorphism  $E \rightarrow E'$  satisfying certain commutative diagrams. Two such morphisms compose in an obvious way. Therefore butterflies from  $A^\bullet$  to  $B^\bullet$  form a groupoid denoted by  $\mathbf{B}(A^\bullet, B^\bullet)$ . A butterfly is *flippable* or *reversible* if both diagonals of (1.1) are extensions.

For more about crossed modules and butterflies in the abelian context, we refer the reader to [17, §12] and [3, §8].

## 1.2 $(A, B)$ -torsors

Let  $A \rightarrow B$  be a morphism of, not necessarily abelian, sheaves. An  $(A, B)$ -torsor is a pair  $(L, x)$ , where  $L$  is an  $A$ -torsor and  $x : L \rightarrow B$  is an  $A$ -equivariant morphism of sheaves (see [10]). A morphism between two pairs  $(L, x)$  and  $(K, y)$  is a morphism of sheaves  $F : L \rightarrow K$  compatible with the action of  $A$  such that the diagram

$$\begin{array}{ccc}
 L & \xrightarrow{F} & K \\
 \downarrow x & & \downarrow y \\
 & = & \\
 & B &
 \end{array}$$

commutes.  $(A, B)$ -torsors form a category denoted by  $\text{TORS}(A, B)$ .

## 1.3 $(\mathcal{A}, \mathcal{B})$ -torsors

Let  $\mathcal{A}$  be a gr-stack, not necessarily Picard. A stack  $\mathcal{P}$  in groupoids is an (right)  $\mathcal{A}$ -torsor if there exists a morphism of stacks

$$m : \mathcal{P} \times \mathcal{A} \longrightarrow \mathcal{A}$$

compatible with the group laws in  $\mathcal{A}$ , and the morphism

$$(\text{pr}, m) : \mathcal{P} \times \mathcal{A} \longrightarrow \mathcal{P} \times \mathcal{P}$$

is an equivalence, and for all  $U \in \mathcal{S}$ ,  $\mathcal{P}_U$  is not empty. [5, §6.1]

Let  $\mathcal{A} \rightarrow \mathcal{B}$  be a morphism of gr-stacks. An  $(\mathcal{A}, \mathcal{B})$ -torsor is a pair  $(\mathcal{L}, x)$ , where  $\mathcal{L}$  is an  $\mathcal{A}$ -torsor, and  $x : \mathcal{L} \rightarrow \mathcal{B}$  is an  $\mathcal{A}$ -equivariant morphism of stacks [1, §6.1], [3, §6.3.4]. A 1-morphism of  $(\mathcal{A}, \mathcal{B})$ -torsors is a pair

$$(F, \mu) : (\mathcal{L}, x) \longrightarrow (\mathcal{K}, y),$$

where  $F : \mathcal{L} \rightarrow \mathcal{K}$  is a morphism of stacks such that

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{F} & \mathcal{K} \\ & \searrow x & \swarrow y \\ & & \mathcal{B} \end{array} \quad \Downarrow \sigma_F$$

and  $\mu$  is a natural transformation of stacks

$$\begin{array}{ccc} \mathcal{L} \times \mathcal{A} & \xrightarrow{F \times 1} & \mathcal{K} \times \mathcal{A} \\ \downarrow & \Downarrow \mu & \downarrow \\ \mathcal{L} & \xrightarrow{F} & \mathcal{K} \end{array}$$

expressing the compatibility of  $F$  with the torsor structure. A 2-morphism of  $(\mathcal{A}, \mathcal{B})$ -torsors  $(F, \mu) \Rightarrow (G, \nu)$  is given by a natural transformation  $\phi : F \Rightarrow G$  satisfying the conditions given in [3, §6.3.4].  $(\mathcal{A}, \mathcal{B})$ -torsors form a 2-stack denoted by  $\text{TORS}(\mathcal{A}, \mathcal{B})$ .

## 1.4 Abelian Sheaves and Picard Stacks

We recall the following results of Aldrovandi and Noohi from [3]. They are going to be referred *sans cesse* throughout the paper.

**Theorem.** [3, Theorem 8.3.1] *For any two length 2 complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , there is an equivalence of groupoids*

$$\text{Hom}(A^\bullet, B^\bullet) \xrightarrow{\sim} \mathbf{B}(A^\bullet, B^\bullet),$$

where  $\text{Hom}(A^\bullet, B^\bullet)$  is the groupoid of additive functors between the Picard stacks associated to  $A^\bullet$  and  $B^\bullet$ .

**Proposition.** [3, Proposition 8.3.2] *Let  $\mathcal{A}$  be a Picard stack. Then there exists a length 2 complex of abelian sheaves  $A^\bullet : A^{-1} \rightarrow A^0$  such that  $\mathcal{A}$  is equivalent to Picard stack  $[A^{-1} \rightarrow A^0]^\sim$ .*

Let  $\mathbf{C}^{[-1,0]}(\mathbf{S})$  denote the bicategory of morphisms of abelian sheaves over  $\mathbf{S}$  with commutative squares as 1-morphisms and homotopies as 2-morphisms. Let  $\mathbf{PIC}(\mathbf{S})$  denote the 2-category of Picard stacks over  $\mathbf{S}$  with 1-morphisms being additive functors and 2-morphisms being natural 2-transformations. Putting the above results together, they prove:

**Theorem.** [3, Proposition 8.4.3] *The functor*

$$\mathbf{C}^{[-1,0]}(\mathbf{S}) \longrightarrow \mathbf{Pic}(\mathbf{S})$$

*defined by sending a morphism of abelian sheaves  $A^\bullet : [A_1 \rightarrow A_2]$  to its associated Picard stack  $[A_1 \rightarrow A_2]^\sim$  is a biequivalence of bicategories.*

*Remark 1.2.* In the same paper, the authors also prove these facts in the non-abelian context by not assuming that stacks and sheaves are necessarily Picard or abelian.

## 1.5 Tricategories

Even though the language of bicategories and tricategories is going to be extensively used, we are not going to remind here in full detail bicategories or tricategories. Just for motivation, a 3-category can be thought as the category of 2-categories with 2-functors or weak 2-functors in the sense of Bénabou [4] and a tricategory as a weakened version of a 3-category. However, we want to recall the triequivalence since the proof of the Theorem 6.4 will follow its definition. For more about bicategories and tricategories, we refer the reader to [11], [12], [16], and [4].

**Definition 1.3.** [16] A trihomomorphism of tricategories  $T : \mathfrak{C} \rightarrow \mathfrak{D}$  is called a triequivalence if it induces biequivalences  $T_{X,Y} : \mathfrak{C}(X, Y) \rightarrow \mathfrak{D}(TX, TY)$  of hom-bicategories for all objects  $X, Y$  in  $\mathfrak{C}$  ( $T$  is locally a biequivalence), and every object in  $\mathfrak{D}$  is biequivalent in  $\mathfrak{D}$  to an object of the form  $TX$  where  $X$  is an object in  $\mathfrak{C}$ .

## 2 Picard 2-Stacks as Torsors

In this section, our goal is to give some of the fundamental facts about 2-stacks and torsors that will be needed throughout the paper. Our main references for 2-stacks with structures such as monoidal, group-like, braided, Picard are [6], [7] and for torsors [1], [5]

### 2.1 2-Stacks

**Definition 2.1.** [8, Definition 6.2] A 2-stack  $\mathbb{P}$  is a fibered 2-category in 2-groupoids such that

- for all  $X, Y$  objects in  $\mathbf{S}_U$ ,  $\mathit{Hom}_{\mathbb{P}_U}(X, Y)$  is a stack over  $\mathbf{S}/U$ ;
- 2-descent data is effective for every object in  $\mathbb{P}$ .

In the above definition 2-groupoids are considered in the sense of Breen [7], that is, 1-morphisms are weakly invertible.

**Definition 2.2.** [7, Definition 8.4] A gr-2-stack  $\mathbb{P}$  is a 2-stack with a morphism  $\otimes : \mathbb{P} \times \mathbb{P} \rightarrow \mathbb{P}$  of 2-stacks, an associativity constraint  $\mathfrak{a}$  compatible with  $\otimes$ , a left unit  $\mathfrak{l}$  and a right unit  $\mathfrak{r}$  constraints compatible with  $\mathfrak{a}$ , and an inverse constraint  $\mathfrak{i}$  with respect to  $\otimes$  compatible with units.

A more detailed definition of gr-2-stack can be found in [6]. Next, following [7, §8.4] we add to gr-2-stacks commutativity constraints with an increasing level of strictness.

**Definition 2.3.** A gr-2-stack  $\mathbb{P}$  is said to be

- *braided*, if there exists a functorial natural transformation

$$R_{X,Y} : X \otimes Y \longrightarrow Y \otimes X$$

that satisfy the 2-braiding axioms of Kapranov and Voevodsky [15] together with the additional condition that, in their terminology, the pair of 2-morphisms defining the induced  $Z$ -systems coincide.

- *strongly braided*, if it is braided, the two hexagons given by the 2-braiding are compatible, and for any  $X, Y$  two objects, there exists a functorial 2-morphism

$$X \otimes Y \begin{array}{c} \xrightarrow{1_{X \otimes Y}} \\ \Downarrow S_{X,Y} \\ \xrightarrow{R_{Y,X} R_{X,Y}} \end{array} X \otimes Y . \quad (2.1)$$

- *symmetric*, if it is strongly braided and the following whiskerings coincide:

$$X \otimes Y \begin{array}{c} \xrightarrow{1_{X \otimes Y}} \\ \Downarrow S_{X,Y} \\ \xrightarrow{R_{Y,X} R_{X,Y}} \end{array} X \otimes Y \xrightarrow{R_{X,Y}} Y \otimes X , \quad (2.2)$$

$$X \otimes Y \xrightarrow{R_{X,Y}} Y \otimes X \begin{array}{c} \xrightarrow{1_{Y,X}} \\ \Downarrow S_{Y,X} \\ \xrightarrow{R_{X,Y} R_{Y,X}} \end{array} Y \otimes X . \quad (2.3)$$

- *Picard*, if it is symmetric and for any object  $X$ , there exists a functorial 2-morphism

$$X \otimes X \begin{array}{c} \xrightarrow{1_{X,X}} \\ \Downarrow S_X \\ \xrightarrow{R_{X,X}} \end{array} X \otimes X \quad (2.4)$$

additive in  $X$  (i.e. there is a relation between  $S_{X \otimes Y}$ ,  $S_X$ , and  $S_Y$ ) such that  $S_{X,X} = S_X * S_X$ .

Further down in the paper, we will be talking about the 3-category of Picard 2-stacks which requires the concept of morphism of Picard 2-stacks. Following Breen [7], we will call such a morphism additive 2-functor. It will be a cartesian 2-functor between the underlying fibered 2-categories compatible with the monoidal, braided, and Picard structures carried by the 2-categories. The compatibility with monoidal structure is already known. In Gordon, Power, Street [11], a monoidal 2-category is defined as a one-object tricategory. More in detail, one can think of a monoidal 2-category as the hom-2-category of a one-object tricategory, whose associativity and unit constraints hold up to 2-isomorphisms and whose modifications are invertible. Then the trihomomorphism [11, Definition 3.1] between such tricategories will be the right definition of morphism between monoidal 2-categories. For the compatibility with the rest of the structures, we refer the reader to the author's thesis [21].

Here is a technical result that we will use several times in our proofs.

**Lemma 2.4.** *Let  $\mathbb{P}$  be a Picard 2-stack and  $A, B$  be two abelian sheaves with additive 2-functors  $\phi : A \longrightarrow \mathbb{P}$  and  $\psi : B \longrightarrow \mathbb{P}$ . Then  $A \times_{\mathbb{P}} B$  is a Picard stack.*



*Proof.* The fibered category  $A \times_{\mathbb{P}} B$  with fibers  $(A \times_{\mathbb{P}} B)|_U$  consisting of

- objects  $(a, f, b)$ , where  $a \in A(U)$ ,  $b \in B(U)$ , and  $f : \phi(a) \rightarrow \psi(b)$  is a 1-morphism in  $\mathbb{P}_U$ ;
- morphisms  $(a, f, \alpha, g, b)$ , where  $\phi(a) \begin{array}{c} \xrightarrow{f} \\ \Downarrow \alpha \\ \xrightarrow{g} \end{array} \psi(b)$  is a 2-morphism in  $\mathbb{P}_U$ ;

is a prestack since for any  $U \in \mathbf{S}$ , 1-morphisms of  $\mathbb{P}$  form a stack over  $\mathbf{S}/U$ . It is in fact a stack.

Let  $((U_i \rightarrow U), (a_i, f_i, b_i), \alpha_{i,j})_{i,j \in I}$  be a descent datum with  $(U_i \rightarrow U)_{i \in I}$  a covering of  $U$ ,  $(a_i, f_i, b_i)$  an object in  $(A \times_{\mathbb{P}} B)_{U_i}$  and  $\alpha_{i,j}$  a 1-morphism in  $(A \times_{\mathbb{P}} B)_{U_{ij}}$  between  $(a_j, f_j, b_j)|_{U_{ij}}$  and  $(a_i, f_i, b_i)|_{U_{ij}}$ . Since  $a_i|_{U_{ij}} = a_j|_{U_{ij}}$ ,  $b_i|_{U_{ij}} = b_j|_{U_{ij}}$  and  $A$  and  $B$  are sheaves, there exist  $a \in A(U)$  and  $b \in B(U)$  such that  $a|_{U_i} = a_i$  and  $b|_{U_i} = b_i$ . Then the collection  $((U_i \rightarrow U), f_i, \alpha_{i,j})_{i,j \in I}$  satisfies the descent in  $\text{Hom}(\phi(a), \psi(b))$ , which is effective since  $\mathbb{P}$  is a Picard 2-stack. That is, there exists  $f \in \text{Hom}(\phi(a), \psi(b))$  and  $\beta_i : f|_{U_i} \Rightarrow f_i$  compatible with  $\alpha_{i,j}$  such that for all  $i \in I$ ,  $(a_i, f|_{U_i}, \beta_i, f_i, b_i)$  is a morphism from  $(a, f, b)|_{U_i}$  to  $(a_i, f_i, b_i)$ . Thus, the descent  $((U_i \rightarrow U), (a_i, f_i, b_i), \alpha_{i,j})_{i,j \in I}$  is effective.

Next, we show that  $A \times_{\mathbb{P}} B$  is Picard. First, let us recall the notation from Definition 2.2.  $\otimes_{\mathbb{P}}$  is the monoidal operation,  $\mathbf{a}$ ,  $\mathbf{l}$ ,  $\mathbf{r}$ ,  $\mathbf{i}$ ,  $R_{-,-}$ ,  $S_{-,-}$ , and  $S_{-}$  are respectively associativity, left unit, right unit, inverse, braiding, symmetry, and Picard constraints. The unnamed arrows in the diagrams below are structural equivalences resulting from additive 2-functors  $\phi$  and  $\psi$ .

*monoidal structure:* The multiplication is defined as


$$(a_1, f_1, b_1) \otimes (a_2, f_2, b_2) := (a_1 + a_2, f_1 f_2, b_1 + b_2),$$

where  $f_1 f_2$  is the morphism that makes the diagram

$$\begin{array}{ccc} \phi(a_1) \otimes_{\mathbb{P}} \phi(a_2) & \xrightarrow{f_1 \otimes_{\mathbb{P}} f_2} & \psi(b_1) \otimes_{\mathbb{P}} \psi(b_2) \\ \downarrow & \nearrow N_m & \downarrow \\ \phi(a_1 + a_2) & \xrightarrow{f_1 f_2} & \psi(b_1 + b_2) \end{array}$$

commute up to a 2-isomorphism  $N_m$ .

For any three objects  $(a_i, f_i, b_i)$  for  $i = 1, 2, 3$ , the associator is given by the morphism  $(a_1 + a_2 + a_3, f_1(f_2 f_3), \alpha_{f_1, f_2, f_3}, (f_1 f_2) f_3, b_1 + b_2 + b_3)$ , where  $\alpha_{f_1, f_2, f_3}$  is defined as the 2-isomorphism of the bottom face that makes the following cube commutative (we ignored  $\otimes_{\mathbb{P}}$  for compactness).



The other 2-isomorphisms of the cube are, the left and right 2-isomorphisms represent the compatibility of the additive 2-functors  $\psi$  and  $\phi$  with the associativity constraint (see Data HTD5 in [11]), the back and front ones are of the form  $N_m$ , the top one is given by the associativity constraint  $\mathfrak{a}$  of  $\mathbb{P}$  on the 1-morphisms.

The object  $I := (0_A, e, 0_B)$ , where  $0_A$  (resp.  $0_B$ ) is the unit object in  $A$  (resp. in  $B$ ) and  $e$  is defined by the 2-commutative diagram

$$\begin{array}{ccc}
1_{\mathbb{P}} & \xrightarrow{=} & 1_{\mathbb{P}} \\
\downarrow & \Downarrow N_u & \downarrow \\
\phi(0_A) & \xrightarrow{e} & \psi(0_B)
\end{array}$$

is the unit in the fibered product  $A \times_{\mathbb{P}} B$ .  $I$  comes with the functorial morphisms  $l_{(a,f,b)} := (a, ef, L_f, f, b)$  and  $r_{(a,f,b)} := (a, fe, R_f, f, b)$ , where  $L_f$  is defined as the 2-isomorphism of the front face that makes the diagram commute (similar diagram for  $R_f$ ).

$$\begin{array}{ccc}
& \phi(0_A) \otimes_{\mathbb{P}} \phi(a) & \xrightarrow{e \otimes_{\mathbb{P}} f} & \psi(0_B) \otimes_{\mathbb{P}} \psi(b) \\
& \swarrow & \uparrow & \swarrow \\
\phi(0_A + a) & \xrightarrow{ef} & \psi(0_B + b) & \\
\downarrow & & \downarrow & \\
= & \downarrow & \Downarrow L_f & \downarrow \\
& 1_{\mathbb{P}} \otimes_{\mathbb{P}} \phi(a) & \xrightarrow{1_{\mathbb{P}} \otimes_{\mathbb{P}} f} & 1_{\mathbb{P}} \otimes_{\mathbb{P}} \psi(b) \\
& \swarrow & \downarrow & \swarrow \\
\phi(a) & \xrightarrow{f} & \psi(b) & \\
& & = & 
\end{array}$$

The other 2-isomorphisms of the cube are, the left and right 2-isomorphisms represent the compatibility of the additive 2-functors  $\psi$  and  $\phi$  with the unit constraint (see Data HTD6 in [11]), the top and bottom ones are of the form  $N_m$ , and the back one is of the form  $N_u$ .

*braiding*: The morphism between  $(a_1, f_1, b_1) \otimes (a_2, f_2, b_2)$  and  $(a_2, f_2, b_2) \otimes (a_1, f_1, b_1)$  is given by  $(a_1 + a_2, f_1 f_2, \beta_{f_1, f_2}, f_2 f_1, b_1 + b_2)$ , where  $\beta_{f_1, f_2}$  is the 2-isomorphism of the bottom face of the commutative cube.

$$\begin{array}{ccccc}
 & \phi(a_1) \otimes_{\mathbb{P}} \phi(a_2) & \xrightarrow{f_1 \otimes_{\mathbb{P}} f_2} & \psi(b_1) \otimes_{\mathbb{P}} \psi(b_2) & \\
 & \swarrow R & \downarrow & \swarrow R & \\
 \phi(a_2) \otimes_{\mathbb{P}} \phi(a_1) & \xrightarrow{f_2 \otimes_{\mathbb{P}} f_1} & \psi(b_2) \otimes_{\mathbb{P}} \psi(b_1) & & \\
 \downarrow & \downarrow & \downarrow & \downarrow & \\
 & \phi(a_1 + a_2) & \xrightarrow{f_1 f_2} & \psi(b_1 + b_2) & \\
 & \swarrow = & \Downarrow \beta_{f_1, f_2} & \swarrow = & \\
 \phi(a_1 + a_2) & \xrightarrow{f_2 f_1} & \psi(b_1 + b_2) & & 
 \end{array} \tag{2.5}$$

The other 2-isomorphisms of the cube are, the left and right 2-isomorphisms represent the compatibility of the additive 2-functors  $\psi$  and  $\phi$  with the braiding structure [21], the front and back ones are of the form  $N_m$ , and the top one represents the compatibility of  $R_{-, -}$  with  $\mathbb{P}$ .

*group like*: Inverse of an object  $(a, f, b)$  is defined as  $(-a, g, -b)$ , where there exists a 2-isomorphism  $\gamma : fg \Rightarrow e$  defined as the 2-morphism of the front face that makes the cube commutative.

$$\begin{array}{ccccc}
 & \phi(a) \otimes_{\mathbb{P}} \phi(-a) & \xrightarrow{f \otimes_{\mathbb{P}} g} & \psi(b) \otimes_{\mathbb{P}} \psi(-b) & \\
 & \swarrow & \downarrow & \swarrow & \\
 \phi(a + (-a)) & \xrightarrow{fg} & \psi(b + (-b)) & & \\
 \downarrow & \downarrow & \downarrow & \downarrow & \\
 & 1_{\mathbb{P}} & \xrightarrow{\Downarrow \gamma} & 1_{\mathbb{P}} & \\
 \downarrow = & \swarrow & \downarrow = & \swarrow & \\
 \phi(0_A) & \xrightarrow{e} & \psi(0_B) & & 
 \end{array}$$

The other 2-isomorphisms of the cube are, the left and right 2-isomorphisms represent the compatibility of the additive 2-functors  $\psi$  and  $\phi$  with the inverse object constraint [21], the top (resp. bottom) one is of the form  $N_m$  (resp.  $N_u$ ), the back one is the inverse object constraint  $i$ .

*symmetry*: We have to verify that the 2-morphism of the bottom face of the diagram obtained by concatenation of the appropriate two cubes of the form (2.5) is identity. This follows from the fact that, 2-morphism of the top face of the concatenated cube pastes to identity with the help of the structural 2-morphisms of type (2.1).

*Picard*: The morphism from  $(a, f, b) \otimes (a, f, b)$  to itself is identity because the 2-morphism of the top face of the diagram (2.5) becomes identity when it is pasted with (2.4).

The compatibility conditions for each structure are trivially satisfied.  $\square$

## 2.2 Picard 2-Stack Associated to a Complex

An immediate example of a Picard 2-stack is the Picard 2-stack associated to a complex of abelian sheaves which is in a sense the only example (see Lemma 6.3). It is already explained in [17] and in [3] how to associate a 2-groupoid to a length 3 complex. However, this 2-groupoid is not a 2-stack. It is not even a 2-prestack (i.e. 1-morphisms only form a prestack but not a stack and 2-descent data are not effective). Therefore to obtain a 2-stack one has to apply the stackification twice. Instead, we are going to use a torsor model for associated stacks. It is more geometric, intuitive, and can be found in [1] for the abelian case, and in [3] for the non-abelian case.

Consider  $A^\bullet$  a complex of abelian sheaves. Let  $\mathcal{A}$  be the associated Picard stack, that is  $[A^{-2} \rightarrow A^{-1}]^\sim \simeq \text{TORS}(A^{-2}, A^{-1})$  and let  $\Lambda_A : \mathcal{A} \rightarrow A^0$  be an additive functor of Picard stacks, where  $A^0$  is considered as a discrete stack (no non-trivial morphisms). It associates to an object  $(L, s)$  in  $\text{TORS}(A^{-2}, A^{-1})$  an element  $\lambda_A(s)$  in  $A^0$ .

We consider  $\text{TORS}(\mathcal{A}, A^0)$  consisting of pairs  $(\mathcal{L}, s)$ , where  $\mathcal{L}$  is an  $\mathcal{A}$ -torsor and  $s : \mathcal{L} \rightarrow A^0$  is an  $\mathcal{A}$ -equivariant map with respect to  $\Lambda_A$ . A morphism between any two pairs is given by another pair  $(F, \gamma)$

$$(F, \gamma) : (\mathcal{L}_1, s_1) \longrightarrow (\mathcal{L}_2, s_2),$$

where  $F$  is a  $\mathcal{A}$ -torsor morphism compatible with the torsor structure up to  $\gamma$ .  $F$  also fits into the commutative diagram.

$$\begin{array}{ccc} \mathcal{L}_1 & \xrightarrow{F} & \mathcal{L}_2 \\ & \searrow s_1 & \swarrow s_2 \\ & & A^0 \end{array} \quad \begin{array}{c} \\ \\ = \end{array}$$

A 2-morphism

$$\begin{array}{ccc} & \xrightarrow{(F, \gamma)} & \\ (\mathcal{L}_1, s_1) & \Downarrow \theta & (\mathcal{L}_2, s_2) \\ & \xrightarrow{(G, \delta)} & \end{array}$$

is given by a natural transformation  $\theta : F \Rightarrow G$  that makes the diagram

$$\begin{array}{ccc} \mathcal{L}_1 \times \mathcal{A} & \xrightarrow{F \times 1} & \mathcal{L}_2 \times \mathcal{A} \\ \downarrow \gamma & \Downarrow \theta \times 1 & \downarrow \\ \mathcal{L}_1 & \xrightarrow{F} & \mathcal{L}_2 \\ \downarrow \delta & \Downarrow \theta & \downarrow \\ \mathcal{L}_1 & \xrightarrow{G} & \mathcal{L}_2 \end{array}$$

commute. It is an immediate result of the following proposition that the 2-stack  $\text{TORS}(\mathcal{A}, A^0)$ , which we have just constructed is Picard.

**Proposition 2.5.** For any  $\mathcal{A} \rightarrow \mathcal{B}$  morphism of Picard stacks,  $\text{TORS}(\mathcal{A}, \mathcal{B})$  is a Picard 2-stack.

*Proof.* From [3, §6.3.4], it follows that  $\text{TORS}(\mathcal{A}, \mathcal{B})$  is a 2-stack. Its group-like structure is defined in [5, §4.5] and *Picardness* is relatively easy to verify.  $\square$

**Definition 2.6.** For any complex of abelian sheaves  $A^\bullet$ , we define the Picard 2-stack associated to  $A^\bullet$  as  $\text{TORS}(\mathcal{A}, A^0)$ .

### 2.3 Homotopy Exact Sequence

Let  $\text{TORS}(\mathcal{A}, A^0)$  be the associated Picard 2-stack to  $A^\bullet$ , then there is a sequence of Picard 2-stacks

$$\mathcal{A} \xrightarrow{\Lambda_A} A^0 \xrightarrow{\pi_A} 2\wp(A^\bullet), \quad (2.6)$$

where  $A^0$  is considered as discrete Picard 2-stack (no non-trivial 1-morphisms and 2-morphisms). The morphism  $\pi_A$  assigns to an element  $a$  of  $A^0(U)$  the pair  $(\mathcal{A}, a)$ , where  $a$  is identified with the morphism  $\mathcal{A} \rightarrow A^0$  sending  $1_{\mathcal{A}} = (A^{-2}, \delta_A)$  to  $a$ . (2.6) is homotopy exact in the sense that  $\mathcal{A}$  satisfies the pullback diagram.

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & 0 \\ \Lambda_A \downarrow & \nearrow & \downarrow \\ A^0 & \xrightarrow{\pi_A} & 2\wp(A^\bullet) \end{array} \quad (2.7)$$

Since  $\mathcal{A}$  is the Picard stack associated to the morphism of abelian sheaves  $\delta_A : A^{-2} \rightarrow A^{-1}$ , it fits into the commutative pullback square of Picard stacks (see the proof of non-abelian version of Proposition 8.3.2 in [3]).

$$\begin{array}{ccc} A^{-2} & \longrightarrow & 0 \\ \delta_A \downarrow & \nearrow & \downarrow \\ A^{-1} & \xrightarrow{\pi_{\mathcal{A}}} & \mathcal{A} \end{array} \quad (2.8)$$

Then pasting the diagrams 2.7 and 2.8 at  $\mathcal{A}$ , we obtain

$$\begin{array}{ccccc}
A^{-2} & \longrightarrow & 0 & & \\
\delta_A \downarrow & \nearrow & \downarrow & & \\
A^{-1} & \xrightarrow{\pi_{\mathcal{A}}} & \mathcal{A} & \longrightarrow & 0 \\
\lambda_A \searrow & & \Lambda_A \downarrow & \nearrow & \downarrow \\
& & A^0 & \xrightarrow{\pi_A} & 2\wp(A^\bullet)
\end{array} \tag{2.9}$$

## 2.4 The 3-category of Picard 2-Stacks

Picard 2-stacks over  $\mathbf{S}$  form an obvious 3-category which we denote by  $2\text{Pic}(\mathbf{S})$ .  $2\text{Pic}(\mathbf{S})$  has a hom-2-groupoid consisting of additive 2-functors, weakly invertible natural 2-transformations, and strict modifications. For any two Picard 2-stacks  $\mathbb{P}$  and  $\mathbb{Q}$ , it is denoted by  $\text{Hom}(\mathbb{P}, \mathbb{Q})$ . If  $\mathbb{P}$  and  $\mathbb{Q}$  are Picard 2-stacks associated to complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , then the hom-2-groupoid will be denoted as  $\text{Hom}(A^\bullet, B^\bullet)$ .

## 3 The 3-category of Complexes of Abelian Sheaves

We start with a definition of a 3-category  $C^{[-2,0]}(\mathbf{S})$  of complexes of abelian sheaves over  $\mathbf{S}$ . We end with an explicit construction of a trihomomorphism  $2\wp$  between  $C^{[-2,0]}(\mathbf{S})$  and the 3-category  $2\text{Pic}(\mathbf{S})$  of Picard 2-stacks over  $\mathbf{S}$ .

### 3.1 Definition of $C^{[-2,0]}(\mathbf{S})$

Although the 3-category of complexes is very well known, in order to setup our notation and terminology, we will describe it explicitly. Its objects are length 3 complexes of abelian sheaves placed in degrees  $[-2, 0]$ . For a pair of objects  $A^\bullet, B^\bullet$ , the hom-2-groupoid  $\text{Hom}_{C^{[-2,0]}(\mathbf{S})}(A^\bullet, B^\bullet)$  is defined as follows:

- A 1-morphism  $f^\bullet : A^\bullet \rightarrow B^\bullet$  is a degree 0 map given by strictly commutative squares.

$$\begin{array}{ccccc}
A^{-2} & \xrightarrow{\delta_A} & A^{-1} & \xrightarrow{\lambda_A} & A^0 \\
\downarrow f^{-2} & & \downarrow f^{-1} & & \downarrow f^0 \\
B^{-2} & \xrightarrow{\delta_B} & B^{-1} & \xrightarrow{\lambda_B} & B^0
\end{array} \tag{3.1}$$

- A 2-morphism  $s^\bullet : f^\bullet \Rightarrow g^\bullet$  is a homotopy map given by the diagram

$$\begin{array}{ccccc}
A^{-2} & \xrightarrow{\delta_A} & A^{-1} & \xrightarrow{\lambda_A} & A^0 \\
\downarrow g^{-2} & & \downarrow g^{-1} & & \downarrow g^0 \\
\downarrow f^{-2} & \searrow s^{-1} & \downarrow f^{-1} & \searrow s^0 & \downarrow f^0 \\
B^{-2} & \xrightarrow{\delta_B} & B^{-1} & \xrightarrow{\lambda_B} & B^0
\end{array}
\tag{3.2}$$

satisfying the relations

$$g^0 - f^0 = \lambda_B \circ s^0, \quad g^{-1} - f^{-1} = \delta_B \circ s^{-1} + s^0 \circ \lambda_A, \quad g^{-2} - f^{-2} = s^{-1} \circ \delta_A.$$

- A 3-morphism  $v^\bullet : s^\bullet \Rightarrow t^\bullet$  is a homotopy map between homotopies  $s^\bullet$  and  $t^\bullet$  given by the diagram

$$\begin{array}{ccccc}
A^{-2} & \xrightarrow{\delta_A} & A^{-1} & \xrightarrow{\lambda_A} & A^0 \\
\downarrow g^{-2} & & \downarrow g^{-1} & & \downarrow g^0 \\
\downarrow f^{-2} & \searrow s^{-1} & \downarrow f^{-1} & \searrow s^0 & \downarrow f^0 \\
B^{-2} & \xrightarrow{\delta_B} & B^{-1} & \xrightarrow{\lambda_B} & B^0
\end{array}
\tag{3.3}$$

satisfying the relations

$$s^0 - t^0 = \delta_B \circ v, \quad s^{-1} - t^{-1} = -v \circ \lambda_A.$$

*Remark 3.1.* In fact, the hom-2-groupoid  $\text{Hom}_{\mathcal{C}^{[-2,0]}(\mathcal{S})}(A^\bullet, B^\bullet)$  is the 2-groupoid associated to  $\tau^{\leq 0}(\text{Hom}^\bullet(A^\bullet, B^\bullet))$ , the smooth truncation of the hom complex  $\text{Hom}^\bullet(A^\bullet, B^\bullet)$ , that is to the complex

$$\text{Hom}^{-2}(A^\bullet, B^\bullet) \longrightarrow \text{Hom}^{-1}(A^\bullet, B^\bullet) \longrightarrow Z^0(\text{Hom}^0(A^\bullet, B^\bullet))$$

of abelian groups, where for  $i = 1, 2$  the elements of  $\text{Hom}^{-i}(A^\bullet, B^\bullet)$  are morphisms of complexes from  $A^\bullet$  to  $B^\bullet$  of degree  $-i$ , and where  $Z^0(\text{Hom}^0(A^\bullet, B^\bullet))$  is the abelian group of cocycles.

## 3.2 Abelian Sheaves and Picard 2-Stacks

**Lemma 3.2.** *There is a trihomomorphism*

$$2\wp : \mathcal{C}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{Pic}(\mathcal{S}) \tag{3.4}$$

between the 3-category  $\mathcal{C}^{[-2,0]}(\mathcal{S})$  of complexes of abelian sheaves and the 3-category  $2\text{Pic}(\mathcal{S})$  of Picard 2-stacks.

*Proof.* We will give a step by step construction of the trihomomorphism and leave the verification of the axioms to the reader.

- Using the notations in section 2.2, given a complex  $A^\bullet$ , we define  $2\wp(A^\bullet)$  as the associated Picard 2-stack, that is  $2\wp(A^\bullet) := \text{TORS}(\mathcal{A}, A^0)$ .

- For any morphism  $f^\bullet : A^\bullet \rightarrow B^\bullet$  of complexes (see diagram (3.1)), there exists a commutative square of Picard stacks

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{\Lambda_A} & A^0 \\
 F \downarrow & = & \downarrow f^0 \\
 \mathcal{B} & \xrightarrow{\Lambda_B} & B^0
 \end{array} \tag{3.5}$$

where  $F$  is induced by  $f^{\bullet < 0} : A^{\bullet < 0} \rightarrow B^{\bullet < 0}$ . From the square (3.5), we construct a 1-morphism  $2\varphi(f^\bullet)$  in  $2\text{PIC}(\mathbf{S})$

$$2\varphi(f^\bullet) : \text{TORS}(\mathcal{A}, A^0) \longrightarrow \text{TORS}(\mathcal{B}, B^0)$$

that sends an  $(\mathcal{A}, A^0)$ -torsor  $(\mathcal{L}, x)$  to  $(\mathcal{L} \wedge^{\mathcal{A}} \mathcal{B}, f^0 \circ x + \Lambda_B)$ . For details, the reader can refer to [1, §6.1].

- For any 2-morphism  $s^\bullet : f^\bullet \Rightarrow g^\bullet$  of complexes (see diagram 3.2), there exists a diagram of Picard stacks

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{\Lambda_A} & A^0 \\
 \begin{array}{c} \downarrow G \\ \downarrow F \end{array} & \begin{array}{c} \searrow \hat{s}^0 \\ \downarrow \end{array} & \begin{array}{c} \downarrow g^0 \\ \downarrow f^0 \end{array} \\
 \mathcal{B} & \xrightarrow{\Lambda_B} & B^0
 \end{array} \tag{3.6}$$

such that for any  $(L, a)$  in  $\mathcal{A}$ , we have the relation

$$G(L, a) - F(L, a) = \hat{s}^0 \circ \Lambda_A(L, a) \quad \text{with} \quad \hat{s}^0(a) = (B^{-2}, s^0(a)).$$

From the relation, we construct a natural 2-transformation  $\theta$

$$\begin{array}{ccc}
 \text{TORS}(\mathcal{A}, A^0) & \xrightarrow{2\varphi(f^\bullet)} & \text{TORS}(\mathcal{B}, B^0) \\
 & \Downarrow \theta & \\
 \text{TORS}(\mathcal{A}, A^0) & \xrightarrow{2\varphi(g^\bullet)} & \text{TORS}(\mathcal{B}, B^0)
 \end{array}$$

in  $2\text{PIC}(\mathbf{S})$  that assigns to any object  $(\mathcal{L}, x)$  in  $\text{TORS}(\mathcal{A}, A^0)$  a 1-morphism  $\theta_{(\mathcal{L}, x)}$

$$\theta_{(\mathcal{L}, x)} : (\mathcal{L} \wedge_{2\varphi(f^\bullet)}^{\mathcal{A}} \mathcal{B}, x_F) \longrightarrow (\mathcal{L} \wedge_{2\varphi(g^\bullet)}^{\mathcal{A}} \mathcal{B}, x_G) \tag{3.7}$$

in  $\text{TORS}(\mathcal{B}, B^0)$ , where  $x_F = f_0 \circ x + \Lambda_B$  and  $x_G = g_0 \circ x + \Lambda_B$ . The morphism (3.7) is defined by sending  $(l, b)$  to  $(l, b - s^0 \circ x(l))$ .



- For any 3-morphism  $v^\bullet : s^\bullet \Rightarrow t^\bullet$  of complexes (see diagram 3.3), there exists a modification  $\Gamma$

$$\begin{array}{ccc} & \xrightarrow{2\varphi(f^\bullet)} & \\ \text{TORS}(\mathcal{A}, A^0) & \begin{array}{c} \theta \Downarrow \\ \Downarrow \Gamma \\ \Downarrow \phi \end{array} & \text{TORS}(\mathcal{B}, B^0) \\ & \xrightarrow{2\varphi(f^\bullet)} & \end{array}$$

in  $2\text{PIC}(\mathcal{S})$  that assigns to any  $(\mathcal{L}, x)$  object of  $\text{TORS}(\mathcal{A}, A^0)$  a natural 2-transformation  $\Gamma_{(\mathcal{L}, x)}$

$$\begin{array}{ccc} & \xrightarrow{\theta_{(\mathcal{L}, x)}} & \\ (\mathcal{L} \wedge_F^{\mathcal{A}} \mathcal{B}, x_F) & \begin{array}{c} \Downarrow \Gamma_{(\mathcal{L}, x)} \end{array} & (\mathcal{L} \wedge_G^{\mathcal{A}} \mathcal{B}, x_G) \\ & \xrightarrow{\phi_{(\mathcal{L}, x)}} & \end{array}$$

in  $\text{TORS}(\mathcal{B}, B^0)$ , where  $\theta_{(\mathcal{L}, x)}$ ,  $\phi_{(\mathcal{L}, x)}$  are of the form (3.7). The natural 2-transformation  $\Gamma_{(\mathcal{L}, x)}$  is defined by assigning to any object  $(l, b)$  in  $(\mathcal{L} \wedge_F^{\mathcal{A}} \mathcal{B}, x_F)$  a morphism

$$\Gamma_{(\mathcal{L}, x)}(l, b) : (l, b - s^0 \circ x(l)) \longrightarrow (l, b - t^0 \circ x(l))$$

in  $(\mathcal{L} \wedge_G^{\mathcal{A}} \mathcal{B}, x_G)$  given by the triple  $(id_l, 1_{\mathcal{A}}, \beta)$  with  $\beta$  being the isomorphism

$$b - s^0 \circ x(l) \longrightarrow b - s^0 \circ x(l) + \delta_B \circ v \circ x(l),$$

and  $id_l$  the identity of  $l$  in  $\mathcal{L}$ , and  $1_{\mathcal{A}}$  the unit element in  $\mathcal{A}$ .

□

## 4 Weak Morphisms of Complexes of Abelian Sheaves

We fix two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ . We define  $\text{Frac}(A^\bullet, B^\bullet)$  a weakened analog of the hom-2-groupoid  $\text{Hom}_{\mathcal{C}[-2,0](\mathcal{S})}(A^\bullet, B^\bullet)$ . We also prove that  $\text{Frac}(A^\bullet, B^\bullet)$  is a bigroupoid.

### 4.1 Definition of $\text{Frac}(A^\bullet, B^\bullet)$

$\text{Frac}(A^\bullet, B^\bullet)$  is a consists of objects, 1-morphisms, and 2-morphisms such that

- An object is an ordered triple  $(q, M^\bullet, p)$ , called fraction

$$(q, M^\bullet, p) : A^\bullet \xleftarrow{q} M^\bullet \xrightarrow{p} B^\bullet$$

with  $M^\bullet$  a complex of abelian sheaves,  $p$  a morphism of complexes, and  $q$  a quasi-isomorphism.

- A 1-morphism from the fraction  $(q_1, M_1^\bullet, p_1)$  to the fraction  $(q_2, M_2^\bullet, p_2)$  is an ordered triple  $(r, K^\bullet, s)$  with  $K^\bullet$  a complex of abelian sheaves,  $r$  and  $s$  quasi-isomorphisms making the diagram

$$\begin{array}{ccccc}
 & & M_1^\bullet & & \\
 & q_1 \nearrow & \uparrow s & \searrow p_1 & \\
 A^\bullet & \xleftarrow{q} & K^\bullet & \xrightarrow{p} & B^\bullet \\
 & q_2 \searrow & \downarrow r & \nearrow p_2 & \\
 & & M_2^\bullet & & 
 \end{array}
 \tag{4.1}$$

commutative.

- A 2-morphism from the 1-morphism  $(r_1, K_1^\bullet, s_1)$  to the 1-morphism  $(r_2, K_2^\bullet, s_2)$  is an isomorphism  $t^\bullet : K_1^\bullet \rightarrow K_2^\bullet$  of complexes of abelian sheaves such that the diagram that we will call “diamond”

$$\begin{array}{ccccc}
 & & M_1^\bullet & & \\
 & q_1 \nearrow & K_1^\bullet & \searrow p_1 & \\
 A^\bullet & \xrightarrow{q_1} & K_1^\bullet & \xrightarrow{p_1} & B^\bullet \\
 & q_2 \searrow & \downarrow r_1 & \nearrow p_2 & \\
 & & M_2^\bullet & & 
 \end{array}
 \begin{array}{c}
 \nearrow s_1 \\
 \dashrightarrow t^\bullet \\
 \searrow s_2
 \end{array}
 \tag{4.2}$$

commutes.

*Remark 4.1.* For reasons of clarity, we will represent the above 2-morphism by the following planar commutative diagram

$$\begin{array}{ccccc}
 & & M_1^\bullet & & \\
 & q_1 \nearrow & \uparrow s_1 & \searrow p_1 & \\
 A^\bullet & \xrightarrow{q_1} & K_1^\bullet & \xrightarrow{p_1} & B^\bullet \\
 & q_2 \searrow & \downarrow r_1 & \nearrow p_2 & \\
 & & M_2^\bullet & & 
 \end{array}
 \begin{array}{c}
 \dashrightarrow t^\bullet \\
 \searrow s_2
 \end{array}$$

where we have ignored the maps from  $K^\bullet$ 's to  $A^\bullet$  and  $B^\bullet$ .


*Remark 4.2.* From the definition of 2-morphisms, it is immediate that all 2-morphisms are isomorphisms.

## 4.2 $\text{Frac}(A^\bullet, B^\bullet)$ is a bigroupoid


**Proposition 4.3.** *Let  $A^\bullet$  and  $B^\bullet$  be two complexes of abelian sheaves. Then  $\text{Frac}(A^\bullet, B^\bullet)$  is a bigroupoid.*

*Proof.* We will describe the necessary data to define the bigroupoid without verifying that they satisfy the required axioms.

- For any two composable morphisms  $(r_1, K_1^\bullet, s_1) : (q_1, M_1^\bullet, p_1) \rightarrow (q_2, M_2^\bullet, p_2)$  and  $(r_2, K_2^\bullet, s_2) : (q_2, M_2^\bullet, p_2) \rightarrow (q_3, M_3^\bullet, p_3)$  shown by the diagram




the composition is defined by the pullback diagram.




That is the composition is the triple  $(r_2 \circ \text{pr}_2, K_1^\bullet \times_{M_2^\bullet} K_2^\bullet, s_1 \circ \text{pr}_1)$ .

- For two 2-morphisms  $t_1 : (r_1, K_1^\bullet, s_1) \Rightarrow (r_2, K_2^\bullet, s_2)$  and  $t_2 : (r_2, K_2^\bullet, s_2) \Rightarrow (r_3, K_3^\bullet, s_3)$  shown by the diagram




the vertical composition is defined by  $t_2^\bullet \circ t_1^\bullet$ .

- For two 2-morphisms  $t^\bullet : (r_1, K_1^\bullet, s_1) \Rightarrow (r_2, K_2^\bullet, s_2)$  and  $u^\bullet : (r'_1, L_1^\bullet, s'_1) \Rightarrow (r'_2, L_2^\bullet, s'_2)$  shown by the diagram




the horizontal composition is given by the natural morphism  $K_1^\bullet \times_{M_2^\bullet} L_1^\bullet \rightarrow K_2^\bullet \times_{M_2^\bullet} L_2^\bullet$  between the pullbacks of pairs  $(r_1, s'_1)$  and  $(r_2, s'_2)$  over  $M_2^\bullet$ .

Any three composable 1-morphisms  $(r_1, K_1^\bullet, s_1)$ ,  $(r_2, K_2^\bullet, s_2)$ , and  $(r_3, K_3^\bullet, s_3)$  can be pictured as a sequence of three fractions



simply by ignoring the maps to  $A^\bullet$  and  $B^\bullet$ . They can be composed in two different ways, either first by pulling back over  $M_2^\bullet$  then over  $M_3^\bullet$  or vice versa. The resulting fractions will be  $(r, (K_1^\bullet \times_{M_2^\bullet} K_2^\bullet) \times_{M_3^\bullet} K_3^\bullet, s)$  and  $(r', K_1^\bullet \times_{M_2^\bullet} (K_2^\bullet \times_{M_3^\bullet} K_3^\bullet), s')$ , respectively, where  $r$  and  $r'$  (resp.  $s$  and  $s'$ ) are equal to  $r_3$  (resp.  $s_1$ ) composed with appropriate projection maps. The 2-isomorphism between these fractions is given by the natural isomorphism between the pullbacks. Thus, the associativity of composition of 1-morphisms is weak.

We also observe that 1-morphisms are weakly invertible. Let  $(r, K^\bullet, s)$  be a 1-morphism from  $(q_1, M_1^\bullet, p_1)$  to  $(q_2, M_2^\bullet, p_2)$ , then  $(s, K^\bullet, r)$  is a weak inverse of  $(r, K^\bullet, s)$  in the sense that the composition  $(r \circ \text{pr}, K^\bullet \times_{M_2^\bullet} K^\bullet, r \circ \text{pr})$  is equivalent to the identity, that is there is a natural 2-transformation  $\theta : r \circ \text{pr} \Rightarrow \text{id} \circ (r \circ \text{pr})$  as shown in the below diagram.



Thus,  $\text{Frac}(A^\bullet, B^\bullet)$  is a bigroupoid.  $\square$

*Remark 4.4.* In the terminology of [2], what we have called fractions are called in the non-abelian context weak morphisms of 2-crossed modules or butterflies of gr-stacks or bats of sheaves.

## 5 Biequivalence of $\text{Frac}(A^\bullet, B^\bullet)$ and $\text{Hom}(A^\bullet, B^\bullet)$

Fix again two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ . In this Section, we prove that the bigroupoid  $\text{Frac}(A^\bullet, B^\bullet)$  of fractions defined in Section 4 is biequivalent to the 2-groupoid  $\text{Hom}(A^\bullet, B^\bullet)$  of additive 2-functors from  $2\varphi(A^\bullet)$  to  $2\varphi(B^\bullet)$  defined in Section 2.4.

### 5.1 Morphisms of Picard 2-Stacks as Fractions

**Lemma 5.1.** *A morphism  $f : A^\bullet \rightarrow B^\bullet$  is a quasi-isomorphism if and only if*

$$2\varphi(f) : 2\varphi(A^\bullet) \longrightarrow 2\varphi(B^\bullet)$$

*is a biequivalence.*

*Proof.* Given  $f : A^\bullet \rightarrow B^\bullet$  a morphism of complexes, we know how to induce a morphism of Picard 2-stacks (see construction of trihomomorphism  $2\varphi(f)$ ). It is also known that a 2-stack (not necessarily Picard) can be seen as a 2-gerbe over its own  $\pi_0$  bounded by the stack  $\mathcal{A}ut(\mathbb{I})$  of automorphisms of identity [7, §8.1]. In particular, the Picard 2-stacks  $\text{TORS}(\mathcal{A}, A^0)$  and  $\text{TORS}(\mathcal{B}, B^0)$  are 2-gerbes over their own  $\pi_0$  bounded by  $\mathcal{A}ut(\mathbb{I}_{2\varphi(A^\bullet)}) \simeq [A^{-2} \rightarrow \ker(\delta_A)]^\sim$  and  $\mathcal{A}ut(\mathbb{I}_{2\varphi(B^\bullet)}) \simeq [B^{-2} \rightarrow \ker(\delta_B)]^\sim$ , respectively. Furthermore, if  $f$  is a quasi-isomorphism, then  $H^{-i}(A^\bullet) \simeq H^{-i}(B^\bullet)$  for  $i = 0, 1, 2$  and thus,  $\pi_i(2\varphi(A^\bullet)) \simeq \pi_i(2\varphi(B^\bullet))$  for  $i = 0, 1, 2$ . So  $\text{TORS}(\mathcal{A}, A^0)$  and  $\text{TORS}(\mathcal{B}, B^0)$  are 2-gerbes with equivalent bands. Therefore they are equivalent.  $\square$

Given an additive 2-functor  $F$  in  $\text{Hom}(A^\bullet, B^\bullet)$ , we will show in the next lemma that there is a corresponding object in  $\text{Frac}(A^\bullet, B^\bullet)$ .

**Lemma 5.2.** *For any additive 2-functor  $F : 2\varphi(A^\bullet) \rightarrow 2\varphi(B^\bullet)$ , there exists a fraction  $(q, M^\bullet, p)$  such that  $F \circ 2\varphi(q) \simeq 2\varphi(p)$ .*

*Proof.* From the sequences

$$\mathcal{A} \xrightarrow{\Lambda_A} A^0 \xrightarrow{\pi_A} 2\wp(A^\bullet) \quad \text{and} \quad \mathcal{B} \xrightarrow{\Lambda_B} B^0 \xrightarrow{\pi_B} 2\wp(B^\bullet),$$

we can construct the commutative diagram

$$\begin{array}{ccccc}
 & & \mathcal{A} \times \mathcal{B} & & \\
 & \swarrow & \downarrow \mu_F & \searrow & \\
 \mathcal{A} & & & & \mathcal{B} \\
 \downarrow \Lambda_A & \searrow \nu_F & & \swarrow \xi_F & \downarrow \Lambda_B \\
 & & \mathcal{E}_F & & \\
 \swarrow \text{pr}_1 & & & & \searrow \text{pr}_2 \\
 A^0 & & & & B^0 \\
 \downarrow \pi_A & \searrow F \circ \pi_A & & & \downarrow \pi_B \\
 2\wp(A^\bullet) & \xrightarrow{F} & & & 2\wp(B^\bullet)
 \end{array} \tag{5.1}$$

where  $\mathcal{E}_F := A^0 \times_{F,B} B^0$ . It follows from the commutativity of the above diagram that  $\mu_F = (\Lambda_A, \Lambda_B)$ . The sequence

$$\mathcal{B} \xrightarrow{\xi_F} \mathcal{E}_F \xrightarrow{\text{pr}_1} A^0 \tag{5.2}$$

is homotopy exact since it is the pullback of the exact sequence  $\mathcal{B} \rightarrow B^0 \rightarrow 2\wp(B^\bullet)$ . From Lemma 2.4, it follows that  $\mathcal{E}_F$  is a Picard stack. Therefore by [3, Proposition 8.3.2], there exists a length 2 complex  $E^\bullet = [\delta_E : E_F^{-1} \rightarrow E_F^0]$  of abelian sheaves such that the associated Picard stack  $\text{TORS}(E_F^{-1}, E_F^0)$  is equivalent to  $\mathcal{E}_F$ . Then by [3, Theorem 8.3.1], there exists a butterfly representing  $\mu_F$

$$\begin{array}{ccc}
 A^{-2} \times B^{-2} & & E_F^{-1} \\
 \downarrow \delta_A \times \delta_B & \searrow \kappa & \downarrow \delta_E \\
 & & P_F \\
 & \swarrow \rho & \searrow j \\
 A^{-1} \times B^{-1} & & E_F^0 \\
 \downarrow \pi_{\mathcal{A}} \times \pi_{\mathcal{B}} & & \downarrow \pi_{\mathcal{E}_F} \\
 \mathcal{A} \times \mathcal{B} & \xrightarrow{\mu_F} & \mathcal{E}_F
 \end{array} \tag{5.3}$$

with  $P_F \simeq (A^{-1} \times B^{-1}) \times_{\mathcal{E}_F} E_F^0$ . From a different perspective, this butterfly can be seen as

$$\begin{array}{ccccc}
 0 & \longrightarrow & E_F^{-1} & \xrightarrow{\delta_E} & E_F^0 \\
 \downarrow & & \downarrow \iota & & \downarrow \text{id} \\
 A^{-2} \times B^{-2} & \xrightarrow{\kappa} & P_F & \xrightarrow{j} & E_F^0 \\
 \downarrow \text{id} & & \downarrow \rho & & \downarrow \\
 A^{-2} \times B^{-2} & \xrightarrow{\delta_A \times \delta_B} & A^{-1} \times B^{-1} & \longrightarrow & 0
 \end{array} \tag{5.4}$$

where each column is an exact sequence of abelian sheaves. The only non-trivial sequence is the second column and its exactness follows from the definition of a butterfly (1.1). So we have a short exact sequence of complexes of abelian sheaves

$$0 \longrightarrow E_F^\bullet \longrightarrow M_F^\bullet \longrightarrow A^{\bullet < 0} \times B^{\bullet < 0} \longrightarrow 0, \quad (5.5)$$

where

$$\begin{aligned} M_F^\bullet &:= A^{-2} \times B^{-2} \longrightarrow P_F \longrightarrow E_F^0, \\ E_F^\bullet &:= 0 \longrightarrow E_F^{-1} \longrightarrow E_F^0, \\ A^{\bullet < 0} \times B^{\bullet < 0} &:= A^{-2} \times B^{-2} \longrightarrow A^{-1} \times B^{-1} \longrightarrow 0. \end{aligned} \quad (5.6)$$

From the lower part of the diagram (5.4) and the definition of  $P_F$ , we deduce that there are morphisms of complexes

$$\begin{array}{ccccc} & & A^{-2} \times B^{-2} & & \\ & \swarrow \text{pr}_1 & \downarrow \kappa & \searrow \text{pr}_2 & \\ & A^{-2} & & & B^{-2} \\ & \downarrow \delta_A & \downarrow P_F & & \downarrow \delta_B \\ & A^{-1} & & & B^{-1} \\ & \downarrow \lambda_A & \downarrow j & & \downarrow \lambda_B \\ & A^0 & & & B^0 \end{array} \quad (5.7)$$

$$\begin{array}{ccc} & M_F^\bullet & \\ & \swarrow q & \searrow p \\ & A^\bullet & B^\bullet \end{array}$$

We claim that  $q$  is a quasi-isomorphism, that is

$$H^{-2}(M_F^\bullet) \simeq \ker(\delta_A), \quad H^{-1}(M_F^\bullet) \simeq \ker(\lambda_A)/\text{im}(\delta_A), \quad H^0(M_F^\bullet) \simeq \text{coker}(\lambda_A).$$

Indeed, from the exact sequence (5.5), we obtain the long exact sequence of homology sheaves

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{-2}(M_F^\bullet) & \longrightarrow & H^{-2}(A^{\bullet < 0}) \times H^{-2}(B^{\bullet < 0}) & \longrightarrow & H^{-1}(E_F^\bullet) \\ & & & & \searrow \partial & & \downarrow \\ & & H^{-1}(M_F^\bullet) & \longrightarrow & H^{-1}(A^{\bullet < 0}) \times H^{-1}(B^{\bullet < 0}) & \longrightarrow & H^0(E_F^\bullet) \\ & & & & \longrightarrow & & H^0(M_F^\bullet) \longrightarrow 0 \end{array} \quad (5.8)$$

On the other hand, by [3, Proposition 6.2.6] applied to the exact sequence (5.2), we get a long exact sequence of homotopy groups

$$0 \longrightarrow \pi_1(\mathcal{B}) \longrightarrow \pi_1(\mathcal{E}_F) \longrightarrow \pi_1(A^0) \longrightarrow \pi_0(\mathcal{B}) \longrightarrow \pi_0(\mathcal{E}_F) \longrightarrow \pi_0(A^0) \longrightarrow 0. \quad (5.9)$$

Since  $\pi_1(A^0) = H^{-1}(A^0) = 0$  and  $\pi_0(A^0) = H^0(A^0) = A^0$ , it follows from (5.9) that we have an isomorphism

$$H^{-2}(B^{\bullet < 0}) \xrightarrow{\simeq} H^{-1}(E_F^\bullet) \quad (5.10)$$

and an exact sequence

$$0 \longrightarrow H^{-1}(B^{\bullet < 0}) \longrightarrow H^0(E_F^\bullet) \longrightarrow A^0 \longrightarrow 0. \quad (5.11)$$

(5.10) implies that  $\partial = 0$  in (5.8). Therefore from (5.8) again, we obtain a short exact sequence

$$0 \longrightarrow H^{-2}(M_F^\bullet) \longrightarrow H^{-2}(A^{\bullet < 0}) \times H^{-2}(B^{\bullet < 0}) \longrightarrow H^{-1}(E_F^\bullet) \longrightarrow 0$$

from which we deduce that  $H^{-2}(M_F^\bullet) \simeq H^{-2}(A^{\bullet < 0}) = \ker(\delta_A)$ .

Now, apply the snake lemma to the short exact sequence (5.11) and to

$$0 \longrightarrow H^{-1}(B^{\bullet < 0}) \longrightarrow H^{-1}(A^{\bullet < 0}) \times H^{-1}(B^{\bullet < 0}) \longrightarrow H^{-1}(A^{\bullet < 0}) \longrightarrow 0$$

in order to get the dashed exact sequence

$$\begin{array}{ccccccc}
& & 0 & \longrightarrow & H^{-1}(M_F^\bullet) & \longrightarrow & \ker(\lambda_A)/\text{im}(\delta_A) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & H^{-1}(B^{\bullet < 0}) & \longrightarrow & H^{-1}(A^{\bullet < 0}) \times H^{-1}(B^{\bullet < 0}) & \longrightarrow & H^{-1}(A^{\bullet < 0}) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & H^{-1}(B^{\bullet < 0}) & \longrightarrow & H^0(E_F^\bullet) & \longrightarrow & A_0 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
& & 0 & \longrightarrow & H^0(M_F^\bullet) & \longrightarrow & \text{coker}(\lambda_A) & \longrightarrow & 0
\end{array}$$

(Dashed lines indicate the snake lemma construction connecting the rows.)

from which it follows  $H^{-1}(M_F^\bullet) \simeq \ker(\lambda_A)/\text{im}(\delta_A)$ , and  $H^0(M_F^\bullet) \simeq \text{coker}(\lambda_A)$  as wanted.

We end this proof by showing that  $F \circ 2\wp(q) \simeq 2\wp(p)$ . (5.7) induces a diagram of Picard 2-stacks



$$\begin{array}{ccc}
& 2\wp(M_F^\bullet) & \\
2\wp(q) \swarrow & & \searrow 2\wp(p) \\
2\wp(A^\bullet) & \xrightarrow{F} & 2\wp(B^\bullet)
\end{array} \quad . \quad (5.12)$$

We claim that (5.12) commutes up to a natural 2-transformation. To show that, it is enough to look at  $2\wp(M_F^\bullet)$  locally. Given  $U \in \mathbf{S}$ ,  $2\wp(M_F^\bullet)_U$  is the 2-groupoid associated to the complex of abelian groups (for the definition of the 2-groupoid associated to a complex see [3] or [17])

$$A^{-2}(U) \times B^{-2}(U) \xrightarrow{\delta} P_F(U) \xrightarrow{\lambda} E_F^0(U)$$

Then, an object of  $2\wp(M_F^\bullet)_U$  is an element  $e$  of  $E_F^0(U)$ . Since  $\mathcal{E}_F := A^0 \times_{F,B} B^0 \simeq \text{TORS}(E_F^{-1}, E_F^0)$ ,  $e$  can be taken as  $(a, f, b)$ , where  $a \in A^0(U)$ ,  $b \in B^0(U)$ , and  $f : F(a) \rightarrow b$  is a 1-morphism in  $2\wp(B^\bullet)_U$ .

A 1-morphism of  $2\wp(M_F^\bullet)_U$  from  $e_1$  to  $e_2$  is given by an element  $p$  of  $P_F(U)$  such that  $\lambda(p) + e_1 = e_2$  in  $E_F^0(U)$ . We can again take  $\lambda(p)$ ,  $e_1$ , and  $e_2$  as  $(a, f, b)$ ,  $(a_1, f_1, b_1)$ , and  $(a_2, f_2, b_2)$ , respectively. Therefore, the addition in  $E_F^0(U)$  should be replaced by the monoidal operation on  $\mathcal{E}_F$  between the triples, that is  $(a, f, b) \otimes_{\mathcal{E}_F} (a_1, f_1, b_1) = (a_2, f_2, b_2)$ . This monoidal operation is described in the proof of the technical Lemma 2.4. It creates a diagram commutative up to a 2-isomorphism in  $2\text{PIC}(B^\bullet)_U$  that defines  $f_2$ .

$$\begin{array}{ccc}
F(a_2) & \xrightarrow{f_2} & b_2 \\
\downarrow \simeq & \theta \not\rightarrow & \downarrow \simeq \\
F(a) \otimes_B F(a_1) & \xrightarrow{f \otimes_B f_1} & b \otimes_B b_1
\end{array}$$

The collection  $(f, \theta)$  gives the natural 2-transformation between  $2\wp(q) \circ F$  and  $2\wp(p)$ .

*Remark 5.3.* Since  $q$  is a quasi-isomorphism in  $C^{[-2,0]}(\mathbf{S})$ , the technical lemma 5.1 implies that  $2\wp(q)$  is a biequivalence in  $2\text{PIC}(\mathbf{S})$ . Therefore, by choosing an inverse of  $2\wp(q)$  up to a natural 2-transformation we can write  $F$  as  $F \simeq 2\wp(p) \circ 2\wp(q)^{-1}$ .

□

## 5.2 Hom-categories of $\text{Frac}(A^\bullet, B^\bullet)$ and $\text{Hom}(A^\bullet, B^\bullet)$

In the next two lemmas, we are going to explore the relation between 1-morphisms (resp. 2-morphisms) of  $\text{Frac}(A^\bullet, B^\bullet)$  and natural 2-transformations (resp. modifications) of Picard 2-stacks.

Suppose we have a natural 2-transformation  $\theta$

$$\begin{array}{ccc}
2\wp(A^\bullet) & \begin{array}{c} \xrightarrow{F} \\ \Downarrow \theta \\ \xrightarrow{G} \end{array} & 2\wp(B^\bullet)
\end{array} \quad (5.13)$$

between the two additive 2-functors  $F, G : 2\wp(A^\bullet) \rightarrow 2\wp(B^\bullet)$ . By Lemma 5.2, we know that there are fractions  $(q_F, M_F^\bullet, p_F)$  and  $(q_G, M_G^\bullet, p_G)$  associated to  $F$  and  $G$ .

**Lemma 5.4.** *For any natural 2-transformation  $\theta$  as in (5.13), there is a 1-morphism in  $\text{Frac}(A^\bullet, B^\bullet)$  between the fractions  $(q_F, M_F^\bullet, p_F)$  and  $(q_G, M_G^\bullet, p_G)$ .*

*Proof.* For  $F$  and  $G$ , we have the following diagrams similar to (5.1)

$$\begin{array}{ccc}
 & \mathcal{A} \times \mathcal{B} & \\
 & \swarrow \quad \searrow & \\
 \mathcal{A} & & \mathcal{B} \\
 \downarrow \Lambda_A & \nu_F \searrow & \swarrow \xi_F \\
 & \mathcal{E}_F & \\
 \downarrow \Lambda_B & & \\
 A^0 & & B^0 \\
 \downarrow \pi_A & F \circ \pi_A \searrow & \downarrow \pi_B \\
 2\wp(A^\bullet) & \xrightarrow{F} & 2\wp(B^\bullet)
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \mathcal{A} \times \mathcal{B} & \\
 & \swarrow \quad \searrow & \\
 \mathcal{A} & & \mathcal{B} \\
 \downarrow \Lambda_A & \nu_G \searrow & \swarrow \xi_G \\
 & \mathcal{E}_G & \\
 \downarrow \Lambda_B & & \\
 A^0 & & B^0 \\
 \downarrow \pi_A & G \circ \pi_A \searrow & \downarrow \pi_B \\
 2\wp(A^\bullet) & \xrightarrow{G} & 2\wp(B^\bullet)
 \end{array}$$

where  $\mathcal{E}_F := A^0 \times_{F,B} B^0$  and  $\mathcal{E}_G := A^0 \times_{G,B} B^0$  are Picard stacks by Lemma 2.4. Therefore by [3, Proposition 8.3.2], there exist  $E_F^{-1} \rightarrow E_F^0$  and  $E_G^{-1} \rightarrow E_G^0$  morphisms of abelian sheaves such that the Picard stack associated to them are respectively  $\mathcal{E}_F$  and  $\mathcal{E}_G$ . The natural 2-transformation  $\theta : F \Rightarrow G$  induces an equivalence  $H : \mathcal{E}_G \rightarrow \mathcal{E}_F$  of Picard stacks defined as follows:

- For any  $(a, g, b)$  object of  $(\mathcal{E}_G)_U$ ,  $H((a, g, b)) := (a, f, b)$ , where  $f$  fits into the commutative diagram

$$\begin{array}{ccc}
 F(a) & \xrightarrow{f} & b \\
 \downarrow \theta_a & = & \downarrow \sim \\
 G(a) & \xrightarrow{g} & b
 \end{array}$$

- For any  $(a, g, \sigma, g', b)$  morphism of  $(\mathcal{E}_G)_U$ ,  $H((a, g, \sigma, g', b)) := (a, f, \tau, f', b)$ , where  $\tau$  is defined by the following whiskering.

$$F(a) \xrightarrow{\theta_a} G(a) \begin{array}{c} \xrightarrow{g} \\ \Downarrow \sigma \\ \xrightarrow{g'} \end{array} b$$

By [3, Theorem 8.3.1],  $H$  corresponds to a butterfly  $[E_G^\bullet, N, E_F^\bullet]$ . Since  $H$  is an equivalence, this butterfly is flippable.

We compose  $H$  and  $\mu_G$  by composing their corresponding butterflies.

$$\begin{array}{ccccc}
A^{-2} \times B^{-2} & & & & E_F^{-1} \\
\downarrow \delta_A \times \delta_B & \searrow \kappa' & & \swarrow i' & \downarrow \delta_E \\
& & P_G \times \begin{matrix} E_G^{-1} \\ E_G^0 \end{matrix} & N & \\
& & \swarrow \rho' & \searrow j' & \\
A^{-1} \times B^{-1} & & & & E_F^0 \\
\downarrow \pi_{\mathcal{A}} \times \pi_{\mathcal{B}} & & & & \downarrow \pi_{\mathcal{E}_F} \\
\mathcal{A} \times \mathcal{B} & \xrightarrow{H \circ \mu_G} & & & \mathcal{E}_F
\end{array}$$

There is also a direct morphism  $\mu_F$  from  $\mathcal{A} \times \mathcal{B}$  to  $\mathcal{E}_F$ .  $\mu_F$  is equivalent to  $H \circ \mu_G$  since they both map an object of  $\mathcal{A} \times \mathcal{B}$  to an object in  $\mathcal{E}_F$  which is isomorphic to the unit object in  $2\wp(B^\bullet)$ . Then by [3, Theorem 8.3.1], there exists an isomorphism  $k$  between the corresponding butterflies of  $\mu_F$  and  $H \circ \mu_G$ , that is the dotted arrow in the diagram below such that all regions commute.

$$\begin{array}{ccccc}
A^{-2} \times B^{-2} & \xrightarrow{\kappa'} & P_G \times \begin{matrix} E_G^{-1} \\ E_G^0 \end{matrix} & N & \xleftarrow{i'} & E_F^{-1} & (5.14) \\
\downarrow \delta_A \times \delta_B & \searrow \kappa & \downarrow k & \swarrow i & \searrow j' & \downarrow \delta_E & \\
& & P_F & & & E_F^0 & \\
& \swarrow \rho' & \downarrow \rho & \searrow j & & \downarrow \pi_{\mathcal{E}_F} & \\
A^{-1} \times B^{-1} & & & & & & \\
\downarrow \pi_{\mathcal{A}} \times \pi_{\mathcal{B}} & & & & & & \\
\mathcal{A} \times \mathcal{B} & \xrightarrow{H \circ \mu_G} & & & & \mathcal{E}_F & \\
& \xrightarrow{\mu_F} & & & & & 
\end{array}$$

Let  $M_F^\bullet : A^{-2} \times B^{-2} \rightarrow P_F \rightarrow E_F^0$  and  $M_G^\bullet : A^{-2} \times B^{-2} \rightarrow P_G \rightarrow E_G^0$ . We claim that, there exists a complex  $K^\bullet$  with quasi-isomorphisms  $r_F$  and  $r_G$  such that all regions in the diagram

$$\begin{array}{ccccc}
& & M_F^\bullet & & \\
& \swarrow q_F & \uparrow r_F & \searrow p_F & \\
A^\bullet & \xleftarrow{q} & K^\bullet & \xrightarrow{p} & B^\bullet \\
& \swarrow q_G & \downarrow r_G & \searrow p_G & \\
& & M_G^\bullet & & 
\end{array} \quad (5.15)$$

commute.

*Proof of the claim:* Let  $K^\bullet : A^{-2} \times B^{-2} \rightarrow P_G \times_{E_G^0} N \rightarrow N$  and define  $r_F$  by the composition

$$\begin{array}{ccccc}
K^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N & \longrightarrow & N & (5.16) \\
\downarrow r_F & \parallel & & \downarrow \text{quotient} & & \downarrow \text{quotient} & \\
M_F^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0}^{E_G^{-1}} N & \longrightarrow & N/E_G^{-1} & \\
& \parallel & & \downarrow & & \downarrow & \\
& A^{-2} \times B^{-2} & \longrightarrow & P_F & \longrightarrow & E_F^0 & 
\end{array}$$

and  $r_G$  by the diagram

$$\begin{array}{ccccc}
K^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N & \longrightarrow & N & (5.17) \\
\downarrow r_G & \parallel & & \downarrow & & \downarrow & \\
M_G^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G & \longrightarrow & E_G^0 & 
\end{array}$$

The commutativity of the diagram (5.16) follows from composition of butterflies. Since  $P_G \times_{E_G^0}^{E_G^{-1}} N \simeq P_F$  and the butterfly  $[E_G^\bullet, N, E_F^\bullet]$  is flippable,  $r_F$  is a quasi-isomorphism. The diagram (5.17) commutes because its left square is a pullback. This implies that  $r_G$  is a quasi-isomorphism.

It remains to show that  $q_F \circ r_F = q_G \circ r_G$ , that is in the diagram below each column closes to a commutative square.

$$\begin{array}{ccccccc}
A^\bullet & A^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & \\
q_F \uparrow & \uparrow & & \uparrow & & \uparrow & \\
M_F^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_F & \longrightarrow & E_F^0 & \\
r_F \uparrow & \parallel & & \uparrow & & \uparrow & \\
K^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N & \longrightarrow & N & \\
r_G \downarrow & \parallel & & \downarrow & & \downarrow & \\
M_G^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G & \longrightarrow & E_G^0 & \\
q_G \downarrow & \downarrow & & \downarrow & & \downarrow & \\
A^\bullet & A^{-2} & \longrightarrow & A^{-1} & \longrightarrow & A^0 & 
\end{array}$$

It is obvious for the first column. The commutativity of the triangles

$$\begin{array}{ccc}
P_G \times_{E_G^0}^{E_G^{-1}} N & \xrightarrow{k} & P_F \\
& \searrow \rho' & \downarrow \rho \\
& & A^{-1} \times B^{-1}
\end{array}
\qquad
\begin{array}{ccc}
\mathcal{E}_G & \xrightarrow{H} & \mathcal{E}_F \\
& \searrow \text{pr}_1 & \swarrow \text{pr}_2 \\
& & A^0
\end{array}$$

imply that the middle and last columns close to a commutative square, respectively (the first triangle is extracted from diagram (5.14)).

In the same way, we also show that  $p_F \circ r_F = p_G \circ r_G$ .  $\square$

Now, suppose we have a modification  $\Gamma$

$$2\wp(A^\bullet) \begin{array}{c} \xrightarrow{F} \\ \theta \Downarrow \Gamma \Downarrow \phi \\ \xrightarrow{G} \end{array} 2\wp(B^\bullet) \quad (5.18)$$

between two natural 2-transformations  $\theta, \phi : F \Rightarrow G$ . We have proved in Lemmas 5.2 and 5.4 that both  $\theta$  and  $\phi$  correspond to a 1-morphism in  $\text{Frac}(A^\bullet, B^\bullet)$ .

**Lemma 5.5.** *Given a modification  $\Gamma$  as in (5.18), there exists a 2-morphism between the two 1-morphisms corresponding to  $\theta$  and  $\phi$ .*

*Proof.* Using the same notations as in Lemma 5.4, we construct a diagram of Picard stacks

$$\mathcal{E}_G \begin{array}{c} \xrightarrow{H_\theta} \\ \Downarrow T \\ \xrightarrow{H_\phi} \end{array} \mathcal{E}_F,$$

where  $T$  is a natural transformation. For any object  $(a, g, b)$  in  $\mathcal{E}_G$ ,  $T_{(a,g,b)}$  is a morphism in  $\mathcal{E}_F$  defined by

$$F(a) \begin{array}{c} \xrightarrow{f_\theta} \\ \Downarrow 1_g * \Gamma_a \\ \xrightarrow{f_\phi} \end{array} b,$$

where

$$F(a) \begin{array}{c} \xrightarrow{\theta_a} \\ \Downarrow \Gamma_a \\ \xrightarrow{\phi_a} \end{array} G(a),$$

and  $H_\theta(a, g, b) = (a, f_\theta, b)$ ,  $H_\phi(a, g, b) = (a, f_\phi, b)$ . By [3, Theorem 5.3.6], the natural transformation  $T$  corresponds to an isomorphism  $t$  between the centers of the butterflies associated to  $H_\theta$  and  $H_\phi$ .

$$\begin{array}{ccccc} E_G^0 & \xrightarrow{\kappa_\theta} & N_\theta & \xleftarrow{\iota_\theta} & E_F^{-1} \\ & \searrow \kappa_\phi & \uparrow & \swarrow \iota_\phi & \\ & & N_\phi & & \\ \delta_{E_G} \downarrow & \rho_\theta \swarrow & \downarrow t & \searrow \rho_\theta & \delta_{E_F} \downarrow \\ E_G^0 & \xrightarrow{\rho_\phi} & N_\phi & \xrightarrow{\rho_\theta} & E_F^0 \\ \pi_{\mathcal{E}_G} \downarrow & & \downarrow & & \downarrow \pi_{\mathcal{E}_F} \\ \mathcal{E}_G & \begin{array}{c} \xrightarrow{H_\theta} \\ \Downarrow T \\ \xrightarrow{H_\phi} \end{array} & & & \mathcal{E}_F \end{array} \quad (5.19)$$

$t$  induces an isomorphism of complexes  $t^\bullet$ .

$$\begin{array}{ccccc}
K_\phi^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N_\phi & \longrightarrow & N_\phi \\
t^\bullet \downarrow & \parallel & & \downarrow \text{id} \times t & & \downarrow t \\
K_\theta^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N_\theta & \longrightarrow & N_\theta
\end{array}$$

The proof finishes by showing that all the regions in the diagram (4.2) commute. The only regions, whose commutativity are non-trivial, are the triangles in the middle sharing an edge marked by the isomorphism  $t^\bullet$ . They commute as well since in the diagram below

$$\begin{array}{ccccc}
M_G^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G & \longrightarrow & E_G^0 \\
r_{G,\phi} \uparrow & \parallel & & \uparrow \text{pr}_1 & & \uparrow \rho_\phi \\
K_\phi^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N_\phi & \longrightarrow & N_\phi \\
t^\bullet \downarrow & \parallel & & \downarrow \text{id} \times t & & \downarrow t \\
K_\theta^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G \times_{E_G^0} N_\theta & \longrightarrow & N_\theta \\
r_{G,\theta} \downarrow & \parallel & & \downarrow \text{pr}_1 & & \downarrow \rho_\theta \\
M_G^\bullet & A^{-2} \times B^{-2} & \longrightarrow & P_G & \longrightarrow & E_G^0
\end{array}$$

each column closes to a commutative triangle. This is immediate for the first two columns. The triangle formed by the last column commutes as well, since it is a piece of the commutative diagram (5.19).  $\square$

For any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , the proofs of Lemmas 5.2 and 5.4 define us a 2-functor

$$2\wp_{(A^\bullet, B^\bullet)} : \text{Frac}(A^\bullet, B^\bullet) \longrightarrow \text{Hom}(A^\bullet, B^\bullet) \quad (5.20)$$

between the bigroupoid  $\text{Frac}(A^\bullet, B^\bullet)$  and the 2-groupoid  $\text{Hom}(A^\bullet, B^\bullet)$  of additive 2-functors between  $2\wp(A^\bullet)$  and  $2\wp(B^\bullet)$  considered as a bigroupoid. In fact, we have proved:

**Theorem 5.6.** *For any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ ,  $2\wp_{(A^\bullet, B^\bullet)}$  is a biequivalence of bigroupoids.*

## 6 The Tricategory of Complexes of Abelian Sheaves

After proving in Section 5 that for any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ ,  $\text{Frac}(A^\bullet, B^\bullet)$  is biequivalent as a bigroupoid to  $\text{Hom}(A^\bullet, B^\bullet)$ , it is clear that the trihomomorphism  $2\wp$  3.4 defined in Section 3.2 cannot be a triequivalence. To attain the triequivalence, we need to consider at least a tricategory with same objects as  $\mathbb{C}^{[-2,0]}(\mathbb{S})$  and with hom-bicategories of the form  $\text{Frac}(A^\bullet, B^\bullet)$ . Furthermore, there is the question of essential surjectivity which we deal with in this section.

### 6.1 Definition of $\mathbb{T}^{[-2,0]}(\mathbb{S})$

We define the tricategory  $\mathbb{T}^{[-2,0]}(\mathbb{S})$  promised at the beginning of the section.

**Definition-Proposition 6.1.**  $\mathbb{T}^{[-2,0]}(\mathcal{S})$  with objects complexes of abelian sheaves, and hom-bigroupoids  $\text{Frac}(A^\bullet, B^\bullet)$ , for any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ , is a tricategory.

*Proof.* We have to verify that  $\mathbb{T}^{[-2,0]}(\mathcal{S})$  has the data given in [12, Definition 3.3.1].

- Objects are complexes of abelian sheaves.
- For any two complexes of abelian sheaves  $A^\bullet$  and  $B^\bullet$ ,  $\text{Frac}(A^\bullet, B^\bullet)$  is the hom-bicategory.
- For any three complexes of abelian sheaves  $A^\bullet$ ,  $B^\bullet$ , and  $C^\bullet$ , the composition is given by the weak functor

$$\otimes_T : \text{Frac}(A^\bullet, B^\bullet) \times \text{Frac}(B^\bullet, C^\bullet) \longrightarrow \text{Frac}(A^\bullet, C^\bullet),$$

which is defined on

1. objects, by

$$\begin{array}{c} M_1^\bullet \\ \swarrow q_1 \quad \searrow p_1 \\ A^\bullet \quad \quad B^\bullet \end{array} \otimes_T \begin{array}{c} M_2^\bullet \\ \swarrow q_2 \quad \searrow p_2 \\ B^\bullet \quad \quad C^\bullet \end{array} = \begin{array}{c} M_1^\bullet \times_{B^\bullet} M_2^\bullet \\ \swarrow q_1 \circ p_{r_1} \quad \searrow p_2 \circ p_{r_2} \\ A^\bullet \quad \quad \quad C^\bullet \end{array}$$

2. 1-morphisms, by

$$\begin{array}{c} M_1^\bullet \\ \swarrow q_1 \quad \uparrow s_1 \quad \searrow p_1 \\ A^\bullet \leftarrow x_1 - K^\bullet \rightarrow y_1 B^\bullet \\ \swarrow q'_1 \quad \downarrow r_1 \quad \searrow p'_1 \\ N_1^\bullet \end{array} \otimes_T \begin{array}{c} M_2^\bullet \\ \swarrow q_2 \quad \uparrow s_2 \quad \searrow p_2 \\ B^\bullet \leftarrow x_2 - L^\bullet \rightarrow y_2 C^\bullet \\ \swarrow q'_2 \quad \downarrow r_2 \quad \searrow p'_2 \\ N_2^\bullet \end{array} = \begin{array}{c} M_1^\bullet \times_{B^\bullet} M_2^\bullet \\ \swarrow q_1 \circ p_{r_1} \quad \uparrow s_1 \times s_2 \quad \searrow p_2 \circ p_{r_2} \\ A^\bullet \leftarrow x_1 \circ p_{r_1} - K^\bullet \times_{B^\bullet} L^\bullet \rightarrow y_1 \circ p_{r_2} C^\bullet \\ \swarrow q'_1 \circ p_{r_1} \quad \downarrow r_1 \times r_2 \quad \searrow p'_2 \circ p_{r_2} \\ N_1^\bullet \times_{B^\bullet} N_2^\bullet \end{array}$$

3. 2-morphisms, by

$$\begin{array}{c} M_1^\bullet \\ \swarrow q_1 \quad \uparrow \quad \searrow p_1 \\ A^\bullet \quad K_1^\bullet \rightarrow K_2^\bullet \quad B^\bullet \\ \swarrow q'_1 \quad \downarrow \quad \searrow p'_1 \\ N_1^\bullet \end{array} \otimes_T \begin{array}{c} M_2^\bullet \\ \swarrow q_2 \quad \uparrow \quad \searrow p_2 \\ B^\bullet \quad L_1^\bullet \rightarrow L_2^\bullet \quad C^\bullet \\ \swarrow q'_2 \quad \downarrow \quad \searrow p'_2 \\ N_2^\bullet \end{array} = \begin{array}{c} M_1^\bullet \times_{B^\bullet} M_2^\bullet \\ \swarrow q_1 \circ p_{r_1} \quad \uparrow \quad \searrow p_2 \circ p_{r_2} \\ A^\bullet \quad K_1^\bullet \times_{B^\bullet} L_1^\bullet \rightarrow K_2^\bullet \times_{B^\bullet} L_2^\bullet \quad C^\bullet \\ \swarrow q'_1 \circ p_{r_1} \quad \downarrow \quad \searrow p'_2 \circ p_{r_2} \\ N_1^\bullet \times_{B^\bullet} N_2^\bullet \end{array}$$

We leave defining the rest of the data as well as verifying that they satisfy the axioms to the reader.  $\square$

The trihomomorphism (3.4) extends to a trihomomorphism

$$2\wp : \mathbb{T}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{Pic}(\mathcal{S}) \tag{6.1}$$

on the tricategory  $\mathbb{T}^{[-2,0]}(\mathcal{S})$  as follows<sup>1</sup>: On objects, it is defined as explained in Section 3.2. On 1-,2-,3-morphisms, by the biequivalence  $2\wp_{(A^\bullet, B^\bullet)}$ , where  $A^\bullet$  and  $B^\bullet$  are any two complexes of abelian sheaves.

Theorem 5.6 implies that (6.1) is already fully faithful in the appropriate sense. In order to prove the triequivalence, one needs to show that it is essentially surjective, as well.

The essential surjectivity depends on the following Lemma, which is similar to in [9, Lemme 1.4.3].

**Lemma 6.2.** *For any set  $E$ , denote by  $\mathbb{Z}(E)$  the free abelian group generated by  $E$ . Let  $\mathbb{C}$  be a Picard 2-category and  $F_0 : E \rightarrow \mathbb{C}$  be a set map. Then  $F_0$  extends to an additive 2-functor  $F : \mathbb{Z}(E) \rightarrow \mathbb{C}$  where  $\mathbb{Z}(E)$  is considered as a 2-category (trivially Picard).*

The proof of the above Lemma will be given in author's thesis [21]. Now, we can show that (6.1) is essentially surjective.

**Lemma 6.3.** *Let  $\mathbb{P}$  be a Picard 2-stack, then there exists a complex of abelian sheaves  $A^\bullet$  such that  $2\wp(A^\bullet)$  is biequivalent to  $\mathbb{P}$ .*

*Proof.* There is a construction analogous to the skeleton of categories. Consider the 2-stack  $2\text{sk}(\mathbb{P})$  obtained from  $\mathbb{P}$  by choosing one representative from each equivalence class of objects. It is obvious that the inclusion  $i : 2\text{sk}(\mathbb{P}) \rightarrow \mathbb{P}$  is a biequivalence. It is also easy to show that  $\text{Ob}(2\text{sk}(\mathbb{P}))$  is a presheaf of sets. Let  $A^0$  be the abelian sheaf over  $\mathcal{S}$  associated to the presheaf  $\{U \rightarrow \mathbb{Z}(2\text{sk}(\mathbb{P}_U))\}$ , where  $\mathbb{Z}(2\text{sk}(\mathbb{P}_U))$  is the free abelian group associated to the set  $\text{Ob}(2\text{sk}(\mathbb{P}_U))$ . By Lemma 6.2, the inclusion  $i$  extends to

$$\pi_{\mathbb{P}} : A^0 \longrightarrow \mathbb{P}$$

an essentially surjective additive 2-functor on  $A^0$ .

Define  $\mathcal{A}$  by the pullback diagram

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & 0 \\ \Lambda_A \downarrow & \nearrow & \downarrow \\ A^0 & \xrightarrow{\pi_{\mathbb{P}}} & \mathbb{P} \end{array} \tag{6.2}$$

of morphisms of Picard 2-stacks, which is similar to (2.7). Then, the sequence of Picard 2-stacks

$$\mathcal{A} \longrightarrow A^0 \longrightarrow \mathbb{P}$$

is exact sequence in the sense of Section 2.3.

On the other hand, from Lemma 2.4, it follows that  $\mathcal{A}$  is a Picard stack. Therefore by [3, Proposition 8.3.2], there exists a morphism of abelian sheaves  $\delta_A : A^{-2} \rightarrow A^{-1}$ , where  $A^{-2}$  is defined by the pullback diagram

---

<sup>1</sup>We commit an abuse of notation by calling both functors (3.4) and (6.1) by  $2\wp$ .



$$\begin{array}{ccc}
A^{-2} & \longrightarrow & 0 \\
\delta_A \downarrow & \nearrow & \downarrow \\
A^{-1} & \xrightarrow{\pi_{\mathcal{A}}} & \mathcal{A}
\end{array} \tag{6.3}$$

and  $\mathcal{A} := \text{TORS}(A^{-2}, A^{-1})$ .

Now putting the diagrams (6.2) and (6.3) together,

$$\begin{array}{ccccc}
A^{-2} & \longrightarrow & 0 & & \\
\delta_A \downarrow & \nearrow & \downarrow & & \\
A^{-1} & \xrightarrow{\pi_{\mathcal{A}}} & \mathcal{A} & \longrightarrow & 0 \\
& \searrow \lambda_A & \downarrow \Lambda_A & \nearrow & \downarrow \\
& & A^0 & \xrightarrow{\pi_{\mathbb{P}}} & \mathbb{P}
\end{array} \tag{6.4}$$

we have a diagram of Picard 2-stacks. It implies that  $A^\bullet : A^{-2} \xrightarrow{\delta_A} A^{-1} \xrightarrow{\lambda_A} A^0$  is a complex.

The Picard 2-stack associated to  $A^\bullet$ , that is  $2\varphi(A^\bullet) := \text{TORS}(\mathcal{A}, A^0)$ , verifies by definition the above diagram (see 2.9).

The biequivalence  $2\varphi(A^\bullet) \simeq \mathbb{P}$  is almost immediate. Essential surjectivity follows from the definition of  $\pi_{\mathbb{P}}$  and equivalence of hom-categories from the fact that  $A^0$  and  $0$  pull back to  $\mathcal{A}$  over  $2\varphi(A^\bullet)$  and to  $\mathbb{P}$ .  $\square$

## 6.2 Main Theorem

Considering  $2\text{PIC}(\mathcal{S})$  as a tricategory, our main result follows from Theorem 5.6 and Lemma 6.3.

**Theorem 6.4.** *The trihomomorphism (6.1) is a triequivalence.*

An immediate consequence of the Theorem 6.4, which was also the motivation for this paper, is the following.

Let  $2\text{PIC}^{\text{bb}}(\mathcal{S})$  denote the category of Picard 2-stacks obtained from  $2\text{PIC}(\mathcal{S})$  by ignoring the modifications and taking as morphisms the equivalence classes of additive 2-functors. Let  $\text{D}^{[-2,0]}(\mathcal{S})$  be the subcategory of the derived category of category of complexes of abelian sheaves  $A^\bullet$  over  $\mathcal{S}$  with  $H^{-i}(A^\bullet) \neq 0$  for  $i = 0, 1, 2$ . We deduce from Theorem 6.4 the following, which generalizes Deligne's result [9, Proposition 1.4.15] from Picard stacks to Picard 2-stacks.

**Corollary 6.5.** *The functor (6.1) induces an equivalence*

$$2\varphi^{\text{bb}} : \text{D}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\text{PIC}^{\text{bb}}(\mathcal{S}) \tag{6.5}$$

of categories.

*Proof.* It is enough to observe from the calculations in Section 4 that  $\pi_0(\text{Frac}(A^\bullet, B^\bullet)) \simeq \text{Hom}_{\mathcal{D}^{[-2,0]}(\mathcal{S})}(A^\bullet, B^\bullet)$ . Since the objects of  $\mathcal{D}^{[-2,0]}(\mathcal{S})$  are same as the objects of  $\mathcal{T}^{[-2,0]}(\mathcal{S})$ , the essential surjectivity follows from the Lemma 6.3.  $\square$

## 7 Stackification

We want to conclude with an informal discussion of stack versions of some of our results. We will assume that all structures are strict unless otherwise stated. Throughout the paper, we dealt with 2- and 3-categories and their weakened versions bi- and tricategories. They can be stackified.

2-stacks over a site are well known [7]. The collection of 2-stacks over  $\mathcal{S}$ , denoted by  $2\text{STACK}(\mathcal{S})$ , comprise a 3-category structure. We can consider the fibered 3-category  $2\mathfrak{STACK}(\mathcal{S})$ , whose fiber over  $U$  is the 3-category  $2\text{STACK}(\mathcal{S}/U)$  of 2-stacks over  $\mathcal{S}/U$ . In [7, Remark 1.12], Breen claims that  $2\mathfrak{STACK}(\mathcal{S})$  is a 3-stack. Hirschowitz and Simpson in [14], generalize this result to weak  $n$ -stacks.

**Theorem.** [14, Théorème 20.5] *The weak  $(n + 1)$ -prestack of weak  $n$ -stacks  $nW\mathfrak{STACK}(\mathcal{S})$  is a weak  $(n + 1)$ -stack over  $\mathcal{S}$ .*

We can use the above facts to deduce that the 3-prestack of Picard 2-stacks  $2\mathfrak{PIC}(\mathcal{S})$  with fibers  $2\text{PIC}(\mathcal{S}/U)$  over  $U$  is a 3-stack.

**Claim.**  $\mathfrak{Hom}(A^\bullet, B^\bullet)$  fibered over  $\mathcal{S}$  in 2-groupoids is a 2-stack where for any  $U \in \mathcal{S}$ , the 2-groupoid  $\text{Hom}(A^\bullet_U, B^\bullet_U)$  of additive 2-functors from  $2\wp(A^\bullet)_U$  to  $2\wp(B^\bullet)_U$  defines the fiber over  $U$ .

We have also fibered analogs for each hom-bicategory  $\text{Frac}(A^\bullet, B^\bullet)$  and for  $\mathcal{T}^{[-2,0]}(\mathcal{S})$ . It follows from the above claim and Theorem 5.6 that the bi-prestack  $\mathbb{F}\text{rac}(A^\bullet, B^\bullet)$  of fractions from  $A^\bullet$  to  $B^\bullet$  with fibers defined by  $\mathbb{F}\text{rac}(A^\bullet_U, B^\bullet_U)$  is a bistack. Then, once an appropriate notion of 3-descent has been specified and all descent data are shown to be effective, we conclude by the characterization proposition [14, Proposition 10.2] for  $n$ -stacks that the tri-prestack of complexes  $\mathfrak{T}^{[-2,0]}(\mathcal{S})$  with fibers  $\mathcal{T}^{[-2,0]}(\mathcal{S}/U)$  is a tristack. The characterization proposition cited above briefly says that  $\mathfrak{P}$  is an  $n$ -stack over  $\mathcal{S}$  if and only if all descent data are effective and for any  $X, Y$  objects of  $\mathfrak{P}_U$ ,  $\text{Hom}_{\mathfrak{P}_U}(X, Y)$  is an  $n - 1$  stack over  $\mathcal{S}/U$ .

*Remark 7.1.* The characterization proposition in [14, Proposition 10.2] is originally enounced for Segal  $n$ -categories,  $n$ -prestacks, and  $n$ -stacks. But again in the same paper, it has been remarked that the proposition holds for non Segal structures [14, §20] where in this case, the weak structure is assumed to be the one defined by Tamsamani. Its definition can be found in [19] and [20]. However, we are being very informal and not discussing here the connection of the weak structure of our categories, pre-stacks and, stacks with the ones mentioned above.

Finally, we define the trihomomorphism of tristacks by localizing the triequivalence (6.1).

$$\mathfrak{T}^{[-2,0]}(\mathcal{S}) \longrightarrow 2\mathfrak{PIC}(\mathcal{S}), \quad (7.1)$$

where  $2\mathfrak{PIC}(\mathcal{S})$  is considered naturally as a tristack. We deduce then its stack analog

**Theorem 7.2.** (7.1) *is a triequivalence of tristacks.*

## References

- [1] Ettore Aldrovandi. 2-gerbes bound by complexes of  $gr$ -stacks, and cohomology. *J. Pure Appl. Algebra*, 212(5):994–1038, 2008.
- [2] Ettore Aldrovandi and Behrang Noohi. Butterflies III: Higher butterflies and higher gr-satcks. *In preparation*.
- [3] Ettore Aldrovandi and Behrang Noohi. Butterflies I: Morphisms of 2-group stacks. *Advances in Mathematics*, 221(3):687 – 773, 2009.
- [4] Jean Bénabou. Introduction to bicategories. In *Reports of the Midwest Category Seminar*, pages 1–77. Springer, Berlin, 1967.
- [5] Lawrence Breen. Bitorseurs et cohomologie non abélienne. In *The Grothendieck Festschrift, Vol. I*, volume 86 of *Progr. Math.*, pages 401–476. Birkhäuser Boston, Boston, MA, 1990.
- [6] Lawrence Breen. Théorie de Schreier supérieure. *Ann. Sci. École Norm. Sup. (4)*, 25(5):465–514, 1992.
- [7] Lawrence Breen. On the classification of 2-gerbes and 2-stacks. *Astérisque*, (225):160, 1994.
- [8] Lawrence Breen. Notes on 1- and 2-gerbes, 2006.
- [9] Pierre Deligne. La formule de dualité globale, 1973. SGA 4 III, Exposé XVIII.
- [10] Pierre Deligne. Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, Proc. Sympos. Pure Math., XXXIII, pages 247–289. Amer. Math. Soc., Providence, R.I., 1979.
- [11] R. Gordon, A. J. Power, and Ross Street. Coherence for tricategories. *Mem. Amer. Math. Soc.*, 117(558):vi+81, 1995.
- [12] Nick Gurski. An algebraic theory of tricategories. *PhD Thesis*, 2006.
- [13] Monique Hakim. *Topos annelés et schémas relatifs*. Springer-Verlag, Berlin, 1972. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 64*.
- [14] Andre Hirschowitz and Carlos Simpson. Descente pour les  $n$ -champs (descent for  $n$ -stacks), 1998.
- [15] M. M. Kapranov and V. A. Voevodsky. 2-categories and Zamolodchikov tetrahedra equations. In *Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park, PA, 1991)*, volume 56 of *Proc. Sympos. Pure Math.*, pages 177–259. Amer. Math. Soc., Providence, RI, 1994.
- [16] Stephen Lack. Bicat is not triequivalent to Gray. *Theory Appl. Categ.*, 18:No. 1, 1–3 (electronic), 2007.

- [17] Behrang Noohi. On weak maps between 2-groups, 2005.
- [18] Behrang Noohi. Notes on 2-groupoids, 2-groups and crossed modules. *Homology, Homotopy Appl.*, 9(1):75–106 (electronic), 2007.
- [19] Carlos Simpson. A closed model structure for  $n$ -categories, internal *hom*,  $n$ -stacks and generalized seifert-van kampen, 1997.
- [20] Zouhair Tamsamani. Sur des notions de  $n$ -catégorie et  $n$ -groupoïde non strictes via des ensembles multi-simpliciaux. *K-Theory*, 16(1):51–99, 1999.
- [21] Ahmet E. Tatar. On the picard 2-stacks. *PhD Thesis, In preparation*.