

STABLE BIRATIONAL EQUIVALENCE AND GEOMETRIC CHEVALLEY-WARNING

XIA LIAO

ABSTRACT. We propose a ‘geometric Chevalley-Warning’ conjecture, that is a motivic extension of the Chevalley-Warning theorem in number theory. It is equivalent to a particular case of a recent conjecture of F. Brown and O.Schnetz. In this paper, we show the conjecture is true for linear hyperplane arrangements, quadratic and singular cubic hypersurfaces of any dimension, and cubic surfaces in \mathbb{P}^3 . The last section is devoted to verifying the conjecture for certain special kinds of hypersurfaces of any dimension. As a by-product, we obtain information on the Grothendieck classes of the affine ‘Potts model’ hypersurfaces considered in [AM].

1. INTRODUCTION

Let \mathbb{F}_q be the finite field of q elements with q a prime power. The Chevalley-Warning theorem states that the number of solutions in \mathbb{F}_q of a system of polynomial equations with n variables is divisible by q , provided that the sum of the degrees of these polynomials is less than n . In [BS], §3.3, F. Brown and O. Schnetz conjecture that a similar statement holds in the Grothendieck ring of varieties over a C_1 field k . They conjecture that the class of an affine k -variety defined by equations satisfying the same degree condition should be a multiple of the class \mathbb{L} of the affine line \mathbb{A}_k^1 . Even the ‘geometric’ case, i.e., when k is an algebraically closed field, appears to be open. We propose the following variant of this conjecture:

Conjecture 1 (Geometric Chevalley-Warning). *Let f_1, \dots, f_l be homogeneous polynomials in $k[x_0, \dots, x_n]$ such that $\sum_{i=1}^l \deg(f_i) < n + 1$, where k is an algebraically closed field of characteristic 0. Then $[Z(f_1, \dots, f_l)] \equiv 1 \pmod{\mathbb{L}}$ in $K_0(\text{Var}_k)$, where $Z(f_1, \dots, f_l)$ denotes the set of zeros of f_1, \dots, f_l in \mathbb{P}^n .*

Over a field k as in this statement, Conjecture 1 is equivalent to the conjecture of Brown and Schnetz. Indeed, let $X = Z(f_1, \dots, f_l) \subseteq \mathbb{P}^n$, then $[X] \cdot (\mathbb{L} - 1) + 1$ is the class of the zero-locus of f_1, \dots, f_l in \mathbb{A}^{n+1} ; the Brown-Schnetz conjecture would imply that this class is $\equiv 0 \pmod{\mathbb{L}}$, and this is equivalent to $[X] \equiv 1 \pmod{\mathbb{L}}$.

In this paper, we show that Conjecture 1 is true for hyperplane arrangements, for quadratic hypersurfaces of any dimension, for cubic surfaces in \mathbb{P}^3 , and for singular cubic hypersurfaces in any dimension. Along the way, we also establish a result which settles some special cases of the conjecture in higher dimensions. The hypothesis that k is an algebraically closed field of characteristic zero is used in our proofs, and it is underlying the contextual remarks that follow in this introduction.

We note that the statement of Conjecture 1 (for any l) is equivalent to the case of hypersurfaces ($l = 1$). Indeed, we have the equality $[Z(f_1, f_2)] = [Z(f_1)] + [Z(f_2)] - [Z(f_1 f_2)]$

Keywords and phrases. Chevalley-Warning theorem, rationality, Grothendieck group.
2010 Mathematics Subject Classification. 14E08, 14N25, 14Q10.

in $K_0(\text{Var})$, and the condition on degrees is satisfied by polynomials on one side of the equation whenever it is satisfied by polynomials on the other side of the equation. It follows that the conjecture is true for $Z(f_1, f_2)$ as long as it is true for the hypersurfaces $Z(f_1)$, $Z(f_2)$, and $Z(f_1 f_2)$. The same type of considerations applies to the zero set of any finite numbers of polynomial equations.

When the variety in consideration is a hypersurface, the condition on degree asked by the geometric Chevalley-Warning conjecture becomes $\deg(f) < n + 1$. This condition is reminiscent of results concerning the “weakened rationality” of varieties. Recall that a variety is rationally chain connected if two general points on the variety can be joined by a chain of rational curves. It is known that a smooth hypersurface of degree d in \mathbb{P}^n is rationally chain connected if and only if $d < n + 1$ [KMM92]. Moreover, if we fix the degree of the hypersurface, and make the dimension of the ambient projective space large enough, then it is proved that a general such hypersurface is unirational [HV92].

Introducing the notion of \mathbb{L} -rationality, Conjecture 1 admits an equivalent reformulation.

Definition 1.1. [AM11] A variety is \mathbb{L} -rational if its class in $K_0(\text{Var})$ is 1 modulo \mathbb{L} .

Conjecture 2. *Every hypersurface of degree $< n + 1$ in \mathbb{P}^n is \mathbb{L} -rational.*

Conjecture 2 postulates that \mathbb{L} -rationality behaves in a sense as rational chain connectedness does. For instance, we know that neither cubic nor quartic smooth threefolds in \mathbb{P}^4 are *rational*, [IM71], [CG72], while they would be both \mathbb{L} -rational and rationally chain connected according to Conjecture 2 and previous discussion.

The notion of \mathbb{L} -rationality is motivated by stable rationality. We recall that two nonsingular irreducible varieties X and Y are “stably birational” if $X \times \mathbb{P}^k$ is birational to $Y \times \mathbb{P}^l$ for some k and l . We say that a nonsingular, complete irreducible variety is ‘stably rational’ if it is stably birational to projective space. For nonsingular varieties, \mathbb{L} -rationality and stable rationality are equivalent. The argument is the following. Recall that the ideal generated by \mathbb{L} in $K_0(\text{Var})$ has a concrete meaning in stably birational geometry. Denote by $\mathbb{Z}[SB]$ the monoid ring generated by stably birational classes of smooth complete irreducible varieties. Then M. Larsen and V. Lunts prove in [LL03] that there exists a surjective homomorphism $\Psi_{SB} : K_0(\text{Var}) \rightarrow \mathbb{Z}[SB]$, mapping the class of a smooth complete variety in $K_0(\text{Var})$ to its class in $\mathbb{Z}[SB]$, and the kernel of this homomorphism is precisely (\mathbb{L}) . Thus, a smooth projective variety is stably birational to projective space precisely when its class in $K_0(\text{Var})$ is 1 modulo \mathbb{L} .

The reader should note that e.g., every *cone* is \mathbb{L} -rational; cf. Lemma 2.4. Also, according to the result we recalled above, a smooth projective rational variety is \mathbb{L} -rational. However, *singular* rational varieties may well not be \mathbb{L} -rational. For example, if the normalization morphism of an irreducible rational curve is not set theoretically injective, then the curve itself is not \mathbb{L} -rational. Thus, ‘most’ singular rational curves are not \mathbb{L} -rational. Examples in higher dimension may be obtained by applying Lemma 3.3. While all varieties considered in this paper are ruled or rational, it is by no means obvious a priori that they should be \mathbb{L} -rational as prescribed by conjecture 1, and as we prove below.

We wrap up this discussion by noting that rationally chain connectedness admits a description analogous to the description of stable rationality we just recalled. In [KR], B. Kahn and R. Sujatha construct a category of pure birational motives by localizing the category of pure motives with respect to certain classes of birational morphisms. They prove ([KR], §3.1) that if X is a rationally chain connected smooth projective F -variety, then $h^\circ(X) = 1$ in $Mot_{rat}^\circ(F, \mathbb{Q})$. Thus, $h^\circ(X)$ plays for rational connectedness a role analogous to the role played by the class of X in $\mathbb{Z}[SB] \cong K_0(\text{Var})/(\mathbb{L})$ for stable rationality.

2. A FEW SIMPLE CASES OF THE CONJECTURE

In this section we verify that the conjecture is true when the degrees of the homogeneous polynomials defining the variety are low. Namely, we will show the following results:

Proposition 2.1. *If X is the union of n or fewer hyperplanes in \mathbb{P}^n , then X is \mathbb{L} -rational.*

Proposition 2.2. *Any quadratic hypersurface in \mathbb{P}^n ($n > 1$) is \mathbb{L} -rational.*

Proposition 2.1 can be proved in a way similar to the reduction of the varieties in Conjecture 1 to hypersurfaces. In fact, the equation of X can be written as $f_1 \dots f_l$ where l is the numbers of hyperplanes and the f_i 's are all linear equations. Then $[X] = Z[f_1 \dots f_{l-1}] + Z[f_l] - Z[f_1 \dots f_{l-1}, f_l]$. The notation $Z[\dots]$ indicates the set of common zeros of the equations appearing in the bracket, separated by commas, as mentioned in Conjecture 1. The last term is the class of the union of $l - 1$ hyperplanes in \mathbb{P}^{n-1} . By induction, all terms on the right side of the equation are equivalent to 1 modulo \mathbb{L} , so is $[X]$.

To prove Proposition 2.2, we observe that any singular quadratic hypersurface is a cone. According to the following lemma and its corollary, the singular case can be taken care of generally, and we are left to consider the class of a nonsingular quadratic hypersurface.

Lemma 2.3. *Let Z be the join of varieties X and Y , which is obtained by connecting pairs of points from X and Y by \mathbb{P}^1 and assuming these rational curves only meet at points of X or Y . If either X or Y is \mathbb{L} -rational, then Z is also \mathbb{L} -rational.*

Proof. Taking out X and Y from the variety Z , we get a bundle over $X \times Y$ whose fiber is \mathbb{P}^1 taken out 2 points. Thus the class of Z in $K_0(\text{Var})$ is $[X] \cdot [Y] \cdot (\mathbb{L} - 1) + [X] + [Y] = [X] \cdot [Y] \cdot \mathbb{L} - ([X] - 1) \cdot ([Y] - 1) + 1$. \square

Corollary 2.4. *If the projective variety $X' \subset \mathbb{P}^m$ is a cone over another projective variety $X \subset \mathbb{P}^n$, $n < m$, then X' is \mathbb{L} -rational.*

Remark 2.5. The union of n or fewer hyperplanes in \mathbb{P}^n is a cone (over an arbitrary point in the set of the intersection). Thus we get a new proof of Propostion 2.1.

A special case of the result by Larsen-Lunts mentioned in the introduction gives an effective treatment of the Chevalley-Warning problem for nonsingular quadratic hypersurfaces, as

Lemma 2.6. *Every rational smooth complete variety is \mathbb{L} -rational.*

Proof. A rational smooth complete variety has the same stable birational class as a point. Thus the difference of its class in $K_0(\text{Var})$ and 1 is in the ideal generated by \mathbb{L} . [LL03] \square

The previous lemmas settle the Chevalley-Warning problem for quadratic hypersurfaces. However, we can give another proof for the nonsingular one avoiding using Lemma 2.6. Let Q be a nonsingular quadratic hypersurface. The projection from the ‘‘north pole’’ gives a nice birational map between Q and \mathbb{A}^n which allows us to stratify Q .

Proof of Proposition 2.2 in the nonsingular case. First, let's fix some notations. Let Q_n be the nonsingular quadratic hypersurface in \mathbb{P}^{n+1} defined by the equation $X_0^2 + X_1^2 + \dots + X_{n+1}^2 = 0$ and Y_n be the affine variety defined by $\sum_{i=1}^{n+1} y_i^2 = 1$ in \mathbb{A}^{n+1} .

Projecting from the point $P = (0, 0, \dots, 1)$, we can establish a birational map between Y_n and \mathbb{A}^n . The formula is given by:

$$x_i = -\frac{y_i}{y_{n+1} - 1}$$

Here, the x_i 's are coordinates of the affine space \mathbb{A}^n .

The inverse rational map from \mathbb{A}^n to Q_n is given by the formula:

$$y_i = \frac{2x_i}{\sum_{i=1}^n x_i^2 + 1} \quad y_{n+1} = \frac{\sum_{i=1}^n x_i^2 - 1}{\sum_{i=1}^n x_i^2 + 1}$$

From this description, it is easy to see the closed set $Z(\sum x_i^2 + 1)$ is not in the image of the polar projection. Then we have the following relation in $K_0(Var)$:

$$(1) \quad [Y_n] - [Z(\sum_{i=1}^n y_i^2)] = [\mathbb{A}^n] - [Y_{n-1}]$$

In addition to this relation, we also have the trivial relation

$$(2) \quad [Q_n] = [Q_{n-1}] + [Y_n]$$

Because the variety $Z(\sum_{i=1}^n y_i^2)$ is a cone, by the proof of 2.4, we get $Z(\sum_{i=1}^n y_i^2)$ is equal to $1 + (\mathbb{L} - 1)[Q_{n-2}]$ in $K_0(Var)$. So we can replace our first equation by:

$$(3) \quad [Y_n] - (1 + (\mathbb{L} - 1)[Q_{n-2}]) = [\mathbb{A}^n] - [Y_{n-1}]$$

With the last two equations and the simple cases $[Q_1] = \mathbb{L} + 1$, $[Y_0] = 2$, $[Y_1] = \mathbb{L} - 1$, we can conclude by an induction on dimension that $[Q_n] \equiv 1 \pmod{\mathbb{L}}$ when $n > 0$ and $[Y_n] \equiv 0 \pmod{\mathbb{L}}$ when $n > 1$. \square

3. CUBIC HYPERSURFACES

The next easiest case to consider is the variety defined by a cubic equation in \mathbb{P}^3 . We have the following theorem:

Theorem 3.1. *Any cubic surface in \mathbb{P}^3 is \mathbb{L} -rational.*

In fact, we can prove something more:

Theorem 3.2. *Any singular cubic hypersurface in \mathbb{P}^n ($n \geq 3$) is \mathbb{L} -rational.*

The following lemma helps to analyze singular cubic hypersurfaces.

Lemma 3.3. *Let $X \subseteq \mathbb{P}^n$ have equation $F = x_n f_k(x_0, \dots, x_{n-1}) + f_{k+1}(x_0, \dots, x_{n-1}) = 0$ where f_k and f_{k+1} are homogeneous polynomials of degree k and $k + 1$ respectively. X is \mathbb{L} -rational if and only if the variety in \mathbb{P}^{n-1} defined by $f_k = 0$ is \mathbb{L} -rational.*

Proof of Lemma 3.3. On the hypersurface $Z(F)$, the equation $f_k = 0$ defines a cone over the point $(0, \dots, 1)$. By Corollary 2.4, this subvariety of the hypersurface $Z(F)$ is \mathbb{L} -rational. On the other hand, we have the isomorphism between the affine open set $f_k \neq 0$ in \mathbb{P}^{n-1} and $Z(F) - Z(f_k)$ provided by $(x_0, \dots, x_{n-1}) \rightarrow (x_0, \dots, x_{n-1}, -\frac{f_{k+1}}{f_k})$. We see the \mathbb{L} -rationality of X is equivalent to the condition that the class of the affine open set $f_k \neq 0$ is in the ideal generated by \mathbb{L} . This happens if and only if the hypersurface $f_k = 0$ in \mathbb{P}^{n-1} is \mathbb{L} -rational. \square

Proof of theorem 3.2. For a singular cubic hypersurface, we can assume one of its singular point is $(0 : \dots : 0 : 1)$ by a change of projective coordinates. We keep using the notation of the previous lemma. The equation of the cubic surface is written as $x_n f_2(x_0, \dots, x_{n-1}) + f_3(x_0, \dots, x_{n-1}) = 0$ or $f_3(x_0, \dots, x_{n-1}) = 0$, depending on the singular point is a double point or a triple point. Then the \mathbb{L} -rationality of the singular cubic hypersurface follows immediately from lemma 3.3, proposition 2.2 or corollary 2.4. \square

Proof of theorem 3.1. A nonsingular cubic surface arises from the blow-up of \mathbb{P}^2 at 6 general points. In particular, it is rational. The \mathbb{L} -rationality follows from Lemma 2.6. The singular case has been done by theorem 3.2. \square

Remark 3.4. One can also approach theorem 3.1 by the classification of cubic surfaces given in [BW79]. That is, one can directly compute their class in $K_0(\text{Var})$ according to the standard equations given in this reference. This however leads to a lengthy computation.

Remark 3.5. The criterion we derived in lemma 3.3 can be applied to give another proof of proposition 2.2. Since as long as the rank of the quadratic form is greater or equal to 2, the equation can be written as $F = x_0x_1 + x_2^2 + \dots$

Corollary 3.6. *If a singular quartic hypersurface in \mathbb{P}^4 has a triple point, then it is \mathbb{L} -rational.*

Proof. Assuming the triple point is $(0 : 0 : 0 : 0 : 1)$, then the equation of the quartic hypersurface can be written as $F = x_4f_3(x_0, x_1, x_2, x_3) + g_4(x_0, x_1, x_2, x_3)$. The \mathbb{L} -rationality follows immediately from lemma 3.3 and theorem 3.1. \square

4. \mathbb{L} -RATIONALITY OF HIGHER DIMENSIONAL VARIETIES

Theorem 4.1. *If the equation of a hypersurface of degree n in \mathbb{P}^m ($m \geq n \geq 4$) can be written as $F = x_n \dots x_4 f_3(x_0, x_1, x_2, x_3) + \sum_{i=5}^n x_n \dots x_i g_{i-1}(x_0, \dots, x_{i-2}) + g_n(x_0, \dots, x_{n-1})$, then this hypersurface is \mathbb{L} -rational.*

Proof. When $m > n$, not all coordinates of \mathbb{P}^m appear in F . In this case F defines a cone in \mathbb{P}^m and the \mathbb{L} -rationality of this hypersurface follows from lemma 2.4. When $m = n$, Rewrite the polynomial as

$$F = x_n [x_{n-1} \dots x_4 f_3(x_0, x_1, x_2, x_3) + \sum_{i=5}^{n-1} x_{n-1} \dots x_i g_{i-1}(x_0, \dots, x_{i-2}) + g_{n-1}(x_0, \dots, x_{n-2})] + g_n(x_0, \dots, x_{n-1}).$$

Then the proof follows by induction, lemma 3.3 and corollary 3.6. \square

Theorem 4.2. *If the equation of the hypersurface of degree at most n in \mathbb{P}^n ($n \geq 4$) has degree 1 in all variables except at most 4 variables, then this hypersurface is \mathbb{L} -rational.*

Proof. Suppose the 4 possibly non-linear variables are x_0, x_1, x_2 , and x_3 . Proceed the proof by an induction on n . When $n = 4$, the equation of the hypersurface is either $x_4 f_k(x_0, x_1, x_2, x_3) + f_{k+1}(x_0, x_1, x_2, x_3) = 0$ ($k \leq 3$) or $f_k(x_0, x_1, x_2, x_3) = 0$ ($k \leq 4$). Since we have checked the \mathbb{L} -rationality of cubic surfaces, quadratic hypersurfaces, the \mathbb{L} -rationality of such hypersurfaces are guaranteed by lemma 3.3 or corollary 2.4. When $n > 4$, if the equation of the hypersurface is written only in terms of x_0, x_1, x_2, x_3 , then it is a cone. Otherwise, let x_n be one of its linear variables. Then the equation of the hypersurface is $x_n f_k(x_0, \dots, x_{n-1}) + f_{k+1}(x_0, \dots, x_{n-1})$ where f_k is again linear in all variables except at most 4 variables. So the \mathbb{L} -rationality follows from lemma 3.3 and the induction hypothesis. \square

Remark 4.3. Theorem 4.2 generalizes Corollary 3.3 from [AM11], since the equations of the “graph hypersurfaces” considered there are linear in all variables, they must be \mathbb{L} -rational by Theorem 4.2.

Example 4.4 (Affine Potts model hypersurface). The previous theorem also gives an easy way to calculate the modulo \mathbb{L} class of the affine Potts model hypersurfaces appearing in [AM] definition (2.2) equation (2.5). The equations for such affine hypersurfaces are

$$Z_G(q, t) = \sum_{G' \subseteq G} q^{k(G')} \prod_{e \in E(G')} t_e,$$

where G' is a subgraph of G , $k(G')$ and $E(G')$ are the number of connected components and the set of edges of the graph G' respectively. Fix q in this equation and denote by n the number of edges of the graph G , then this equation defines the affine Potts model hypersurface in \mathbb{A}^n . Its class in $K_0(\text{Var})$ is congruent to 1 modulo \mathbb{L} if n is odd and congruent to -1 modulo \mathbb{L} when n is even.

The calculation goes as follows. These hypersurfaces are defined by inhomogeneous polynomials of degree n in \mathbb{A}^n , linear in all variables, where n is the number of the edges of the graph in consideration. Homogenize the equation and write it as $F = 0$, then clearly F is a homogeneous polynomial of degree n linear in all variables except the variable x_0 introduced in homogenizing. Now the class of the affine Potts model hypersurface is $[Z(F)] - [Z(x_0, x_1 x_2 \dots x_n)]$. $Z(F)$ is \mathbb{L} -rational by the previous theorem, and now we are left to calculate $[Z(x_0, x_1 x_2 \dots x_n)]$ which is the class of the union of n hyperplanes in \mathbb{P}^{n-1} .

Let x_1, \dots, x_n be the projective coordinates of \mathbb{P}^{n-1} . Consider the complement of $Z(x_1 x_2 \dots x_n)$, which can be explicitly expressed as points $(x_1 : \dots : x_n)$ such that all projective coordinates are nonzero. We see this affine open set is isomorphic to the $(n-1)$ -fold cartesian product of $(\mathbb{A}^1 - \{pt\})$. So the class of the union of n hyperplanes in \mathbb{P}^{n-1} equals $[\mathbb{P}^{n-1}] - (\mathbb{L} - 1)^{n-1}$. Taking into account that $[\mathbb{P}^{n-1}] = 1 + \mathbb{L} + \dots + \mathbb{L}^{n-1}$, we see this class is $1 + (-1)^n$ modulo \mathbb{L} . We conclude the class of the affine Potts model hypersurface is \mathbb{L} -rational when n is odd and congruent to -1 modulo \mathbb{L} when n is even.

REFERENCES

- [AM] Paolo Aluffi and Matilde Marcolli. A motivic approach to phase transitions in Potts models. arXiv:1102.3462.
- [AM11] Paolo Aluffi and Matilde Marcolli. Graph hypersurfaces and a dichotomy in the Grothendieck ring. *Lett. Math. Phys.*, 95(3):223–232, 2011.
- [BS] Francis Brown and Oliver Schnetz. A K3 in ϕ^4 . arXiv:1006.4064.
- [BW79] J. W. Bruce and C. T. C. Wall. On the classification of cubic surfaces. *J. London Math. Soc. (2)*, 19(2):245–256, 1979.
- [CG72] C. Herbert Clemens and Phillip A. Griffiths. The intermediate Jacobian of the cubic threefold. *Ann. of Math. (2)*, 95:281–356, 1972.
- [HV92] Kapil H. Paranjape and V. Srinivas. Unirationality of the general complete intersection of small multidegree. *Astérisque*, 211:241–248, 1992.
- [IM71] V. A. Iskovskih and Ju. I. Manin. Three-dimensional quartics and counterexamples to the Lüroth problem. *Mat. Sb. (N.S.)*, 86(128):140–166, 1971.
- [KMM92] János Kollár, Yoichi Miyaoka, and Shigefumi Mori. Rationally connected varieties. *J. Algebraic Geom.*, 1(3):429–448, 1992.
- [KR] Bruno Kahn and R. Sujatha. birational geometry and localisation of categories. arXiv:0805.3753.
- [LL03] Michael Larsen and Valery A. Lunts. Motivic measures and stable birational geometry. *Mosc. Math. J.*, 3(1):85–95, 259, 2003.

MATHEMATICS DEPARTMENT, FLORIDA STATE UNIVERSITY, TALLAHASSEE FL 32306, U.S.A.
E-mail address: `xliao@math.fsu.edu`