

# Generalized Lantern Relations and Planar Line Arrangements

Eriko Hironaka\*

June 30, 2011

## Abstract

In this paper, we show that to each planar line arrangement defined over the real numbers, for which no two lines are parallel, one can write down a corresponding relation on Dehn twists that can be read off from the combinatorics and relative locations of intersections. This gives an alternate proof of Wajnryb's generalized lantern relations, and of Endo, Mark and Horn-Morris' daisy relations.

## 1 Introduction

The study of hyperplane arrangements has a rich history in the realms of topology, algebraic geometry, and analysis (see, for example, [O-T]). Several authors have studied the topology of planar line arrangements using braid monodromy (see, for example, [MT], [C-S], [Cor]). While easy to draw, the deformation theory of real planar line arrangements holds many mysteries. For example, there are topologically distinct real line arrangements with equivalent combinatorics [ARCM] (see also, [Ryb]); and by the Sylvester-Gallai theorem [ES] there are planar line arrangements defined over complex numbers, whose combinatorics cannot be duplicated by a real line arrangement, for example, the lines through the 9 flexes of a smooth cubic plane curve.

In this paper, we use the theory of line arrangements to generate relations on Dehn twists in the mapping class group  $MCG(S)$  of an oriented surface  $S$  of finite type. The lantern relation on Dehn twists is of interest because it is one of five simple to state relations that generate all relations in the Dehn-Lickorish-Humphreys presentation of  $MCG(S)$  [Waj1] (see also [Lic], [Hum], [Bir], [Mat]). The lantern relation also plays an important role in J. Harer's proof that the abelianization of  $MCG(S)$  is trivial if  $S$  is a closed surface of genus  $g \geq 3$  [Har] (see also [FM], Sec. 5.1.2).

To state the lantern relations, we first fix notation. Let  $S$  be an oriented surface of finite type. If  $S$  is closed, the mapping class group  $MCG(S)$  is the group of isotopy classes of self-homeomorphisms of  $S$ . If  $S$  has boundary components, then the definition of  $MCG(S)$  is the same, except that we impose the condition that all maps fix the boundary of  $S$  pointwise. For a compact annulus  $A$ ,

---

\*This work was partially supported by a grant from the Simons Foundation (#209171to Eriko Hironaka).

$MCG(A)$  is isomorphic to  $\mathbb{Z}$  and is generated by a right or left Dehn twist around its core curve. As illustrated in Figure 1, a right Dehn twist takes a path on  $A$  transverse to the core curve to a path that wraps once around the core curve turning in the right hand direction (a left Dehn twist correspondingly turns in the left direction) as it passes through  $c$ . The Dehn twist can also be thought of as being obtained by rotating one of the boundary components by  $360^\circ$  while leaving the other boundary component fixed. Each simple closed curve  $c$  on  $S$  determines a right Dehn




Figure 1: Right Dehn twist.

twist on an annulus neighborhood of  $c$ , and this Dehn twist extends by the identity to all of  $S$ . The isotopy class  $\delta_c$  of this map is the (*right*) *Dehn twist centered at  $c$*  and is an element of  $MCG(S)$ .

The original statement and proof of the lantern relation appears in Dehn's 1938 paper [Deh] and relates a product of three interior Dehn twists to four boundary twists on a genus zero surface with four boundary components. The relation was little known until it was rediscovered by D. Johnson [Joh]. B. Wajnryb gave the following generalized version in [Waj2] (Lemma 1.17).

**Theorem 1** *Let  $S_{0,n+1}^c \subset S$  be a surface of genus zero with  $n+1$  boundary components  $d_0, d_1, \dots, d_n$ . There is a collection of simple closed curves  $a_{i,j}, 1 \leq i < j \leq n$  in the interior of  $S_{0,n+1}^c$ , so that*

- (i) *for each  $i, j$ ,  $a_{i,j}$  separates  $d_i \cup d_j$  from the rest of the boundary components, and*
- (ii) *there is a relation on Dehn twists*

$$\partial_0(\partial_1, \dots, \partial_n)^{n-2} = \alpha_{1,2}\alpha_{1,3}, \dots, \alpha_{1,n}, \alpha_{2,3}, \dots, \alpha_{2,n}, \dots, \alpha_{n-2,n-1}, \alpha_{n-2,n}, \alpha_{n-1,n}, \quad (1)$$

*where  $\alpha_{i,j} = \delta_{a_{i,j}}$  is the right Dehn twist around  $a_{i,j}$ , and  $\partial_i = \delta_{d_i}$  is the right Dehn centered at a curve parallel to the boundary components  $d_i$ .*

We now restate Theorem 1 in terms of line arrangements in  $\mathbb{R}^2$ .

**Theorem 2** *Let  $\mathcal{L}$  be a union of  $n \geq 3$  distinct lines in the  $(x, y)$ -plane over the reals with distinct slopes and no slope parallel to the  $y$ -axis. Let  $\mathcal{I} = \{p_1, \dots, p_s\}$  be the intersection points on  $\mathcal{L}$  numbered by largest to smallest  $x$ -coordinate. For each  $L \in \mathcal{L}$ ,  $m_L$  is the number of points in  $\mathcal{I} \cap L$ . Let  $S_{0,n+1}^c$  be a surface of genus zero and  $n+1$  boundary components  $d_L$ , one for each  $L \in \mathcal{L}$  and an extra boundary component  $d_0$ . Then there are simple closed curve  $a_{p_k}, k = 1, \dots, s$  on  $S_{0,n+1}^c$  so that the following holds:*

(i) each  $a_{p_k}$  separates

$$\bigcup_{p_k \in L \in \mathcal{L}} d_L$$

from the rest of the boundary curves; and

(ii) the Dehn twists  $\partial_L = \delta_{d_L}$  and  $\alpha_{p_k} = \delta_{a_{p_k}}$  satisfy

$$\partial_0 \prod_{L \in \mathcal{L}} \partial_L^{m_L-1} = \alpha_{p_s} \cdots \alpha_{p_1}. \quad (2)$$

**Remark 1.** In Equation (2), the terms on the left side commute, while the ones on the right typically don't. Thus, the ordering of  $p_1, \dots, p_s$  matters, and reflects the subtle combinatorics of line arrangements. The curves  $a_{p_k}$  can be found explicitly (see Section 2.2, Lemma 3).

**Remark 2.** The relations in  $\text{MCG}(S_{0,n+1}^c)$  give rise to relations on  $\text{MCG}(S)$  for any surface  $S$  admitting an embedding  $S_{0,n+1}^c \hookrightarrow S$ .




Figure 2: Three lines in general position, and curves defining associated lantern relation drawn two ways

When  $n = 3$ , Theorem 1 gives the standard Lantern relation

$$\partial_0 \partial_1 \partial_2 \partial_3 = \alpha_{1,2} \alpha_{1,3} \alpha_{2,3}.$$

The core curves for these Dehn twists and the corresponding line arrangements are shown in Figure 2. The diagram to the right is the motivation for the name of this relation.

Here is the main idea of our proof of Theorem 2. First consider a great ball  $B \subset \mathbb{C}^2$  containing all the points of intersection of  $\mathcal{L}$ . Let  $\mathbb{C}\mathbb{P}^2$  be the projective compactification of  $\mathbb{C}^2$ . Then the complement of  $B$  in  $\mathbb{C}\mathbb{P}^2$  is a neighborhood of the “line at infinity” or  $L_\infty = \mathbb{C}\mathbb{P}^2 \setminus \mathbb{C}^2$ . Thus the monodromy of  $\rho$  restricted to the exterior of  $B$  depends only on the way  $\mathcal{L}$  intersects  $L_\infty$ . If no lines in  $\mathcal{L}$  are parallel to each other, then without changing any slopes, and hence the topology of  $\mathbb{C}\mathbb{P}^2 \setminus B$ , it is possible to move the lines in  $\mathcal{L}$  to obtain a new configuration  $\mathcal{T}$  where all lines meet at a single point. Since moving the lines in the arrangement in this way does not change the monodromy, we get an equality between the two monodromies defined by  $\mathcal{L}$  and  $\mathcal{T}$ . Thus, Theorem 2 follows from an analysis of the monodromy of line arrangements on compactified fibers of a generic projection. The monodromy is reinterpreted as point pushing maps, and we keep track of twisting on the boundary components of the compactified fibers using the complex coordinate system on  $\mathbb{C}^2$ .

This paper is organized as follows. In Section 2.1 we recall the Moishezon-Teicher braid monodromy representation of a free group associated to a planar line arrangement. We refine the representation so that its image is the mapping class group of compactified fibers in Section 2.2. In Section 2.3, we prove Theorem 2 using deformations of line arrangements and give further variations of the lantern relation, including the daisy relation Theorem 7.

**Acknowledgments:** The author is grateful to J. Mortada and D. Margalit for helpful discussions and comments.

## 2 Real line arrangements and relations on Dehn twists

In this section, we recall properties of line arrangements  $\mathcal{L}$  in the complex plane defined by real equations and the monodromy on generic fibers under linear projections

$$\mathbb{C}^2 \setminus \mathcal{L} \rightarrow \mathbb{C}.$$

A key ingredient to our analysis is B. Moishezon and M. Teicher description of the monodromy as elements of the pure braid group (see, for example, [MT] and [Hir]). We generalize this braid monodromy by studying the action of the monodromy not only on generic fibers of  $\rho$ , but also on their compactifications as genus zero surfaces with boundary. This leads to a proof of Theorem 2.

The ideas in this section can be generalized to more arbitrary plane curves. An investigation of the topology of plane curve complements using such general projections appears in work of O. Zariski and E. van Kampen [Kam]. We leave this as a topic for future study.

### 2.1 Braid monodromy defined by planar line arrangements over the reals

In this section we recall the braid monodromy associated to a real line arrangement. For convenience we choose Euclidean coordinates  $(x, y)$  for  $\mathbb{C}^2$  so that no line is parallel to the  $y$ -axis, and no two intersection points have the same  $x$  coordinate. For  $i = 1, \dots, n$ , let  $L_i$  be the zero set of a linear equation in  $x$  and  $y$  with real coefficients:

$$L_i : y = m_i x + c_i \quad m_i, c_i \in \mathbb{R},$$

and assume that the lines are ordered so that the slopes satisfy:

$$m_1 > m_2 > \dots > m_n.$$

Let  $\mathcal{I} = \mathcal{I}(\mathcal{L}) = \{p_1, \dots, p_s\} \in \mathbb{C}^2$  be the collection of intersections points of  $\mathcal{L}$  ordered so that the  $x$ -coordinates are strictly decreasing.

Let  $\rho : \mathbb{C}^2 \rightarrow \mathbb{C}$  be the projection of  $\mathbb{C}^2$  onto  $\mathbb{C}$  given by  $\rho(x, y) = x$ . For each  $x \in \mathbb{C}$ , let

$$F_x = \rho^{-1}(x) \setminus \mathcal{L}.$$

The  $y$  coordinate allows us to uniformly identify  $F_x$  with the complement in  $\mathbb{C}$  of  $n$  points  $L_i(x)$ , where  $L_i(x)$  is the unique point in

$$\rho^{-1}(x) \cap L_i.$$

Let  $x_0 \in \mathbb{R}$  be greater than any point in  $\rho(\mathcal{I})$ . Then there is a natural map

$$\gamma : [0, 1] \rightarrow \mathbb{C} \setminus \rho(\mathcal{I})$$

from paths based at  $x_0$  to a braid on  $n$  strands in  $\mathbb{C}$  parameterized by

$$\{L_i(\gamma(t)) : i = 1, \dots, n\}.$$

Since two homotopic paths give rise to isotopic braids, and composition of paths corresponds to composition of braids, this defines a homomorphism

$$\beta : \pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), x_0) \rightarrow B(S^2, n + 1)$$

from the fundamental group to the spherical braid group on  $n + 1$  strands.

The (*braid*) *monodromy* of  $(\mathbb{C}^2, \mathcal{L})$  with respect to the projection  $\rho$  and basepoint  $x_0$  is the homomorphism

$$\sigma_{\mathcal{L}} : \pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), x_0) \rightarrow \text{MCG}(F_{x_0}), \quad (3)$$

given by the composition of  $\beta$  and the braid representation

$$B(S^2, n + 1) \rightarrow \text{MCG}(S_{0, n+1}) = \text{MCG}(F_{x_0}),$$

from the braid group to the mapping class group on a genus zero surface with  $n + 1$  punctures..

We now study the image of simple generators of  $\pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), x_0)$  in  $\text{MCG}(F_{x_0})$ .




Figure 3: Simple loop generators for  $\pi_1(\mathbb{C} \setminus \rho(\mathcal{I}))$ .

By a simple loop in  $\pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), x_0)$ , we mean a path of the form  $g = f_p g_p f_p^{-1}$ , where

$$\begin{aligned} g_p : [0, 1] &\rightarrow \mathbb{C} \setminus \rho(\mathcal{I}) \\ t &\mapsto p + \epsilon_p e^{2\pi i t} \end{aligned}$$

and  $f_p$  is a path from  $x_0$  to  $p + \epsilon$  whose image is in the upper half plane except at its endpoints. Since  $\pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), x_0)$  is generated by simple loops, it is enough to understand the monodromy in the image of these elements.

In order to describe the monodromy of a simple loop  $L_p = f_p g_p f_p^{-1}$ , we study how  $F_x^c$  is transformed as  $x$  follows each of the segments of  $L_p$ .




Figure 4: Monodromy defined by  $g_p$  with the real part of  $\mathcal{L}$  drawn in.

First we look at  $g_p$ . Let  $L_{j_1}, L_{j_2}, \dots, L_{j_k}$  be the lines in  $\mathcal{L}$  that pass through  $p$ . We can assume by a translation of coordinates that  $p = (0, 0)$ , and  $L_{j_r}$  is defined by an equation of the form

$$y = m_r x$$

where  $m_1 > m_2 > \dots > m_k$ . Then as  $t$  varies in  $[0, 1]$ , the intersection of  $L_{j_r}$  with  $F_{g_p(t)}$  is given by

$$L_{j_r}(t) = (\epsilon_p e^{2\pi i t}, m_r \epsilon_p e^{2\pi i t}).$$

The other lines in  $\mathcal{L}$  locally can be thought of as having constant slope, hence their intersections with  $F_{g_p(t)}$  retain their order and stay outside a circle on  $F_{g_p(t)}$  enclosing  $L_{j_1}(g_p(t)), \dots, L_{j_k}(g_p(t))$  (see Figure 4). Let  $a_p^{\text{loc}} \subset F_{p+\epsilon}$  be this circle. The restriction of  $\rho$  to  $\mathbb{C}^2 \setminus \mathcal{L}$  defines a trivial bundle over the image of  $f_p$ . Thus  $a_p^{\text{loc}}$  determines a simple closed curve  $a_p$  on  $F_{x_0}$  separating  $L_{j_1}(x_0), \dots, L_{j_k}(x_0)$  from the rest of the  $L_j(x_0)$ .

Next we notice that lifting over  $f_p$  defines a mapping class on  $F_{x_0}$ . This is because there is a canonical identification of  $F_{x_0}$  and  $F_x$  for any  $x \in \mathbb{R} \setminus \rho(\mathcal{I})$  given by the natural ordering of  $\mathcal{L} \cap \rho^{-1}(x)$  by the size of the  $y$ -coordinate from largest to smallest. Thus  $f_p$  determines a braid  $\beta_p$  on  $n$  strands. We have shown the following.

**Lemma 3** *The image  $\sigma_{\mathcal{L}}(g)$  in  $MCG(F_{x_0})$ , where  $g = f_p g_p f_p^{-1}$  is given by*

$$\beta_p \cdot \sigma_p \cdot \beta_p^{-1},$$

where  $\sigma$  is a right Dehn twist around the pullback of  $a_p$ .

## 2.2 Monodromy on compactified fibers

In this section, we define the monodromy representation of  $\pi_1(F_{x_0}, y_0)$  into  $\text{MCG}(F_{x_0}^c)$ , where  $F_{x_0}^c$  is a compactification of  $F_{x_0}$  as a compact surface with boundary.

As before choose coordinates for  $\mathbb{C}^2$ , and let  $\mathcal{L} = \cup_{i=1}^n L_i$  be a planar line arrangement defined over the reals with distinct slopes. Assume all points of intersection  $\mathcal{I}$  have distinct  $x$ -coordinates. Let  $\epsilon > 0$  be such that the  $\epsilon$  radius disks  $B_\epsilon(p)$  around the points  $p \in \rho(\mathcal{I})$  are disjoint. Let  $\delta > 0$  be such that the  $\delta$  radius tubular neighborhoods  $N_\delta(L_i)$  around  $L_i$  are disjoint in the complement of

$$B_\delta(\rho(\mathcal{I})) = \cup_{p \in \rho(\mathcal{I})} \rho^{-1}(B_\delta(p)).$$

Let  $D$  be a disk in  $\mathbb{C}$  containing all points of  $\rho(\mathcal{I})$  in its interior, and having  $x_0$  on its boundary. Let  $N_\infty$  be the product of a disk centered at the origin of  $\mathbb{C}$  so that  $\mathbb{C} \times N_\infty$  contains  $\mathcal{L} \cap \rho^{-1}(D)$ .

For each  $x \in \mathbb{C} \setminus \rho(\mathcal{I})$ , let

$$F_x^c = \rho^{-1}(x) \cap N_\infty \setminus \subset F_x \setminus N_\epsilon(L_i).$$

For each  $x \in D$  and  $i = 1, \dots, n$ , let

$$d_i(x) = \partial N_\epsilon(L_i) \cap \rho^{-1}(x),$$

and let

$$d_\infty(x) = \partial N_\infty \cap \rho^{-1}(x).$$

We are now ready to define the monodromy on the compactified fibers

$$\sigma_{\mathcal{L}}^c : \pi_1(F_{x_0}, y_0) \rightarrow \text{MCG}(F_{x_0}^c).$$

Let  $\eta$  be the inclusion homomorphism

$$\eta : \text{MCG}(F_{x_0}^c) \rightarrow \text{MCG}(F_{x_0}).$$

Then we have a commutative diagram

$$\begin{array}{ccc} \pi_1(\mathbb{C} \setminus \rho(\mathcal{I}), y_0) & \xrightarrow{\sigma_{\mathcal{L}}^c} & \text{MCG}(F_{x_0}^c) \\ & \searrow \sigma_{\mathcal{L}} & \downarrow \eta \\ & & \text{MCG}(F_{x_0}). \end{array}$$

The kernel of  $\eta$  is generated by Dehn twists centered at the boundary components of  $F_{x_0}^c$  (Theorem 3.18, [FM]). Thus, in order to describe  $\sigma_{\mathcal{L}}^c$ , we need to understand what twists occur near boundary components in the monodromy associated to the paths  $g_p$  and  $f_p$  defined in Section 2.1.

Consider the simplest case when  $L \subset \mathbb{C}^2$  is a single line defined by

$$y = mx.$$

Let  $N_\delta(L)$  be the tubular neighborhood

$$N_\delta(L) = \{(x, L(x) + y) : |y| < \delta\}.$$

Then  $N_\delta(L) \cap F_{g(t)}$  is a disk centered at  $L(g(t))$  of radius  $\delta$ . The boundary  $\partial N_\delta(L)$  is a trivial bundle over  $\mathbb{C} \setminus B_\delta(\rho(\mathcal{L}))$  with trivialization defined by the framing of  $\mathbb{C}$  by real and complex coordinates.

Now assume that there are several lines  $L_{j_1}, \dots, L_{j_k}$  meeting at  $p$ , and that  $\rho(p) + \epsilon$  is the basepoint for  $g_p$  a loop of going clockwise around a circle of radius epsilon around  $p$ . Let  $L$  be a line through  $p$  with slope equal to the average of those of  $L_{j_1}, \dots, L_{j_k}$ , and let  $\epsilon > 0$  be such that  $N_\epsilon(L) \cap F_{\rho(p)+\epsilon}$  contains  $d_{j_1}(g_p(0)), \dots, d_{j_k}(g_p(0))$ , but no other boundary components of  $F_{g_p(0)}$ . Let

$$d_N(g_p(0)) = \partial N_\epsilon(L) \cap F_{g_p(0)}.$$

Then looking back at Figure 4, we see that the points  $L_{j_1}(g_p(t)), \dots, L_{j_k}(g_p(t))$  rotate as a group  $360^\circ$  in the counterclockwise direction as  $t$  ranges in  $[0, 1]$ . The corresponding mapping class on the bounded portion of  $F_{g_p(0)}$  enclosed by  $d_N$  can be thought of as being obtained by simultaneously point pushing the inner boundary components  $d_{j_1}(g_p(0))$  in a clockwise direction.

Figure 5 and Figure 6 illustrates the Dehn twist  $\partial_{d_N}$  centered at a simple closed curve parallel to  $d_N$  and the monodromy  $\sigma_{\mathcal{L}}^c(g_p)$  in the case when  $\mathcal{L}$  is a union of 4 lines meeting at a single point  $p$ . The middle picture illustrates the fiber  $F_{g_p(0.5)}$  half way around the circle traversed by  $g_p$ . From




Figure 5: The mapping class  $\partial_{d_N}$ .




Figure 6: The monodromy defined by  $g_p$ .

the effect of the mapping class on relative curves, we see the equality

$$\sigma_{\mathcal{L}}^c(g_p) = (\partial_{d_1} \partial_{d_2} \partial_{d_3} \partial_{d_4})^{-1} \partial_{d_N}.$$

More generally we have the following lemma.

**Lemma 4** *Let  $L_{j_1}, \dots, L_{j_k}$  be the lines meeting at  $p$ , and let*

$$\begin{aligned} g_p : [0, 1] &\rightarrow \mathbb{C} \setminus \mathcal{L} \\ t &\mapsto \rho(p) + \epsilon e^{2\pi i t}. \end{aligned}$$

*Then the monodromy on  $F_{g_p(0)}^c$  defined by  $g_p$  is given by*

$$\sigma_{\mathcal{L}}^c(g_p) = (\partial_{d_{j_1}} \cdots \partial_{d_{j_k}})^{-1} \partial_{d_N}.$$

### 2.3 Deformations of line arrangements

To finish our proof we analyze the effect of deforming a line arrangement.

Let

$$\mathcal{L} = \bigcup_{i=1}^n L_i$$

be a finite union of real lines in the Euclidean plane,  $\mathbb{R}^2$  with no two lines parallel. Let  $\mathcal{T}$  be the complexified real line arrangement with all  $n$  lines intersecting at a single point  $p_0$ . Let  $\rho : \mathbb{C}^2 \rightarrow \mathbb{C}$  be a generic projection, and let  $D \subset \mathbb{C}$  be a disk of radius  $r$  centered at the origin containing  $\rho(\mathcal{I})$  and  $\rho(p_0)$  in its interior. Let

$$\begin{aligned} \gamma : [0, 1] &\rightarrow \mathbb{C} \\ t &\mapsto r e^{2\pi i t}. \end{aligned}$$

**Lemma 5** *The monodromies  $\sigma_{\mathcal{L}}(\gamma)$  and  $\sigma_{\mathcal{T}}(\gamma)$  are the same.*

**Proof.** Identify  $\mathbb{C}^2$  with  $\mathbb{C}\mathbb{P}^2 \setminus L_\infty$ , where  $\bar{L}_\infty$  is a projective line. Identify the points in  $\bar{L}_\infty$  with slopes of lines in  $\mathbb{C}^2$ . The complement

$$N_\infty = \mathbb{C}\mathbb{P}^2 \setminus \rho^{-1}(D)$$

is naturally identified with a disk bundle over  $\bar{L}_\infty$  punctured at a point. Since the restriction of  $\rho$  to  $N_\infty \cap \bar{\mathcal{T}}$  and to  $N_\infty \cap \bar{\mathcal{L}}$  are the same, the monodromy over the boundary path  $\gamma$  of  $D$  is the same.  $\square$

**Proof of Theorem 2.** By Lemma 5,  $\sigma_{\mathcal{L}}(\gamma) = \sigma_{\mathcal{T}}(\gamma)$ . Figure —refpathhomotopy-fig gives and illustration of two equivalent representations of the homotopy type of  $\gamma$ . Let




Figure 7: Two representatives of  $\gamma$  in  $\pi_1(\mathbb{C} \setminus \rho(\mathcal{I}))$ .

By Lemma 3 and Lemma 4, we have

$$\sigma_{\mathcal{T}}(\gamma) = (\partial_{d_1} \cdots \partial_{d_n})^{-1} \partial_{d_\infty}.$$

Let  $p_1, \dots, p_s$  be the elements of  $\mathcal{I}$  numbered by decreasing  $x$ -coordinate. Then for each  $i = 1, \dots, s$ , we have

$$\sigma_{\mathcal{L}}(f_{p_i} g_{p_i} f_{p_i}) = (\partial_{d_{j_1}} \circ \partial_{d_{j_k}})^{-1} \alpha_{p_i}$$

where

$$\alpha_{p_i} = \partial_{f_{p_i}^* d_{N_{p_i}}}$$

is the pullback of  $d_{N_{p_i}}$  along the path  $f_{p_i}$ . Thus,

$$\sigma_{\mathcal{L}}(\gamma) = (\partial_{d_1}^{m_1} \circ \partial_{d_s}^{m_s})^{-1} \alpha_{p_s} \circ \alpha_{p_1},$$

where  $m_i$  is the number of elements in  $\mathcal{I} \cap L_i$ . □

To show that Theorem 1 follows from Theorem 2, we need to show that the ordering given in Equation (1) can be obtained by a union of lines  $\mathcal{L}$  satisfying the conditions. To do this, we start with a union of lines  $\mathcal{T}$  intersecting in a single point. Let  $L_1$  be the line in  $\mathcal{T}$  with largest slope. Translate  $L_1$  in the positive  $x$  direction without changing its slope. The translated line  $L'_1$  intersect the other lines in simple points starting with  $L_2$  and ending with  $L_n$ . Continue for each line from highest to lowest slope, making sure with each time that the shifting  $L$  creates new intersections lying to the left of all previously created ones.

More generally, we can deform the lines through a single point  $\mathcal{T}$  to one in general position  $\mathcal{L}$  so that the only condition on the resulting ordering on the pairs of lines is the following. A pair  $(i, j)$  must precede  $(i, j + 1)$  for each  $1 \leq i < j \leq n$ . Thus, we obtain another restatement of Theorem 1.

**Theorem 6** *Let  $\{p_1, \dots, p_s\}$  be an ordering of the pairs  $(i, j)$ ,  $1 \leq i < j \leq n$ , so that for all  $i$ , the sequence*

$$(i, i + 1), (i, i + 2) \dots, (i, n)$$

*is strictly decreasing. Then there a lantern relation of the form*

$$\partial_0(\partial_1 \cdots \partial_n)^{n-2} = \alpha_{p_1} \cdots \alpha_{p_s}.$$

### 3 Applications

Although it is known that all relations on the Dehn-Lickorish-Humphreys generators can be obtained from the braid, chain, lantern and hyperelliptic relations, there are some other nice symmetric relations that come out of line arrangements that are not trivially derived from the four generating ones. We conclude this paper with a sampling.

#### 3.1 Daisy relation

Consider the line arrangements given in Figure 8. As pointed out to me by D. Margalit, this relation was recently also discovered H. Endo, T. Mark, and J. Van Horn-Morris using rational blowdowns of 4 manifolds [EMHM]. We follow their nomenclature and call this the *daisy relation*.

Let  $S_{0,n+1}^c$  denote the compact surface of genus 0 with  $n + 1$  boundary components. Consider the configuration of simple closed curves shown in Figure ???. Let  $d_0, \dots, d_n$  be the boundary components of  $S_{0,n+1}^c$ . Let  $d_1$  be the distinguished boundary component at the center of the arrangement, and let  $d_0, d_2, \dots, d_n$  be the boundary components arranged in a circle (ordered in the clock wise direction around  $d_1$ ). Let  $a_{1,k}$  be a simple closed loop encircling  $d_1$  and  $d_k$ , where




Figure 8: Line arrangement, and associated arrangement of curves (n=6).

$k = 0, 2, 3, \dots, n$ . Let  $\partial_i$  be the Dehn twist centered at  $d_i$ , and let  $\alpha_{1,k}$  be the Dehn twist centered at  $a_{1,k}$ .

**Theorem 7** [*Daisy-relation*] For  $n \geq 3$ , the Dehn twists on  $S_{0,n+1}^c$  satisfy the relation

$$\partial_0 \partial_1^{n-2} \partial_2 \cdots \partial_n = \alpha_{1,0} \alpha_{1,n} \cdots \alpha_{1,2}$$

where  $\partial_i$  is the Dehn twist centered at the boundary component  $d_i$ , and  $\alpha_{1,j}$  is the Dehn twist centered at curves  $a_{1,j}$ .

When  $n = 3$ , Theorem 7 specializes to the usual lantern relation.

**Proof.** We associate the boundary component  $d_i$  with  $L_i$  for  $i = 1, \dots, n$ , and  $d_0$  with the “line at infinity”. Theorem 2 applied to the line arrangement in Figure 8 gives:

$$\partial_0 (\partial_1 \cdots \partial_n)^{-1} = R_{p_n} \cdots R_{p_1}$$

where  $p_1, \dots, p_N$  are the intersection points of the line arrangement  $\mathcal{L}$  ordered by largest to smallest  $x$ -coordinate. For this configuration,  $p_k$  gives rise to

$$R_{p_k} = (\partial_1 \partial_{k+1})^{-1} \alpha_{1,k+1},$$

for  $k = 1, \dots, n-1$ . Noting that the loop that separates  $d_2 \cup \cdots \cup d_n$  from  $d_0 \cup d_1$  can be written as  $a_{1,0}$ , we have

$$R_{p_n} = \partial_2 \cdots \partial_{n-1} \alpha_{1,0}$$

yielding the desired formula. □

**Remark 3.** Let

$$\beta : B(S^2, n+1) \rightarrow \text{MCG}(S_{0,n+1})$$




Figure 9: Alternate drawing of the daisy configuration ( $n=4$ )

be the braid representation from the spherical braid group to the mapping class group. Recall the relation  $R$  in  $B(S^2, n+1)$  given by

$$\begin{aligned}
 (\sigma_1^2)(\sigma_1^{-1}\sigma_2^2\sigma_1)\cdots(\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{n-1}\sigma_n^2\sigma_{n-1}\cdots\sigma_1) &= \sigma_1\cdots\sigma_{n-1}\sigma_n^2\sigma_{n-1}\cdots\sigma_1 \\
 &= 1
 \end{aligned}$$

This induces a relation  $R'$  in  $\text{MCG}(S_{0,n+1})$ . The daisy relation can be considered as the lift of  $R'$  under the inclusion homomorphism  $\eta$ .

### 3.2 Flying saucer relation

As a final example, we consider a configuration of  $n \geq 5$  lines, with  $n-2$  meeting in a single point. There are several ways this can be drawn. We give one example in Figure 10. Other line arrangements satisfying these conditions will give similar relations, but the drawings of the associated curves will be more complicated.




Figure 10: Configuration of lines and associated configuration of curves ( $n=5$ ).

As before, let  $d_0, \dots, d_n$  be the boundary components of  $S_{0,n+1}^c$ . The boundary component  $d_i$  is associated to the line  $L_i$  for  $i=1, \dots, n$ , and  $d_0$  is the boundary component associated to the “line

at infinity". Let  $a_{i,j}$  be the loop encircling  $d_i \cup d_j$  and no other boundary component. Let  $c$  be the loop encircling  $d_3, \dots, d_n$  drawn in Figure 10 (with  $n=5$ ). For convenience  $a_{1,n}$  is drawn as a loop encircling  $d_2 \cup d_n \cup d_0$ . An alternate drawing is given in Figure 10.

**Theorem 8 (Flying saucer relation)** *Let  $\partial_i$  be the right Dehn twist centered at  $d_i$ ,  $\alpha_{i,j}$  the right Dehn twist centered at  $a_{i,j}$ , and  $\beta$  the right Dehn twist centered at  $c$ . Then*

$$\delta_0 \delta_1^{n-2} \delta_2 \cdots \delta_{n-1} \delta_n^{n-2} = \alpha_{n-1,n} \alpha_{n-2,n} \cdots \alpha_{2,n} \beta \alpha_{1,n} \alpha_{1,n-1} \cdots \alpha_{1,2}$$

**Proof.** Theorem 2 applied to the line arrangement in Figure 10 gives the equation

$$\partial_0 (\partial_1 \cdots \partial_n)^{-1} = R_{p_1} \cdots R_{p_{n-2}} R_{p_{n-1}} R_q R_{s_1} \cdots R_{s_{n-2}},$$

where

$$\begin{aligned} R_{p_k} &= (\partial_1 \partial_{k+1})^{-1} \alpha_{1,k+1} \\ R_q &= (\partial_2 \cdots \partial_{n-1})^{-1} \beta \\ R_{s_k} &= (\partial_n \partial_{k+1})^{-1} \alpha_{k+1,n}. \end{aligned}$$

Putting these together yields the desired formula. □

One sees from Figure 10 that the order of  $R_{p_{n-1}}$  and  $R_q$  may be interchanged.




Figure 11: Alternate drawing of the saucer configuration ( $n=5$ )

## References

- [ARCM] B. Artal, J. Ruber, J. Cogolludo, and M. Marco. Topology and combinatorics of real line arrangements. *Comp. Math.* **141** (2005), 1578–1588.
- [Bir] J. Birman. Mapping class groups of surfaces. In *Braids (Santa Cruz, CA, 1986)*, volume 78. Amer. Math. Soc., Providence, RI, 1988.

- [C-S] D. Cohen and A. Suci. Braided monodromy of plane algebraic curves and hyperplane arrangements. *Comm. Math. Helv.* **72** (1997), 285–315.
- [Cor] R. Cordovil. The fundamental group of the complement of the complexification of a real arrangement of hyperplanes. *Adv. App. Math.* **21** (1998).
- [Deh] M. Dehn. Die Gruppe der Abbildungsklassen. *Acta Math.* **69** (1938).
- [EMHM] H. Endo, T. E. Mark, and J. Van Horn-Morris. Monodromy substitutions and rational blowdowns. *J. Topol.* **4** (2011), 227–253.
- [ES] P. Erdős and R. Steinberg. Three point collinearity. *American Mathematical Monthly* **51** (1944), 169–171.
- [FM] B. Farb and D. Margalit. *A Primer on Mapping Class Groups*. Princeton University Press, 2011.
- [Har] J. Harer. The second homology group of the mapping class group. *Invent. Math.* **72** (1983), 221–239.
- [Hir] E. Hironaka. Abelian coverings of the complex projective plane branched along configurations of real lines. *Mem. of the A.M.S.* **105** (1993).
- [Hum] S. Humphries. *Generators of the mapping class group*, volume 722, pages 44–47. Springer, 1979.
- [Joh] D. Johnson. Homeomorphisms of a surface which act trivially on homology. *Proc. of the Amer. Math. Soc.* **75** (1979), 119–125.
- [Lic] W.B.R. Lickorish. On the homeotopy group of a 2-manifold. *Proc. Cambridge Philos. Soc.* **62** (1964), 679–681.
- [Mat] M. Matsumoto. A simple presentation of mapping class groups in terms of Artin groups. *Sugaku Expositions* **15** (2002), 223–236.
- [MT] B. Moishezon and M. Teicher. Braid group technique in complex geometry. I. Line arrangements in  $\mathbf{CP}^2$ . In *Braids (Santa Cruz, CA, 1986)*, volume 78 of *Contemp. Math.*, pages 425–555. Amer. Math. Soc., Providence, RI, 1988.
- [O-T] P. Orlik and H. Terao. *Arrangement of Hyperplanes*, volume 300 of *Grundlehren der math. Wissenschaften*. Springer-Verlag, Berlin, 1992.
- [Ryb] G. Rybnikov. On the fundamental group of the complement of a complex hyperplane arrangement. *DIMACS: Technical Report* (1994), 33–50.
- [Kam] E. van Kampen. On the fundamental group of an algebraic curve. *Am. Jour. Math.* **55** (1933), 255–260.
- [Waj1] B. Wajnryb. A simple presentation for the mapping class group of an orientable surface. *Israel J. Math.* **45** (1983), 1983.
- [Waj2] B. Wajnryb. Mapping class group of a handlebody. *Fund. Math.* **158** (1998), 195–228.