


## MAD 5932-01 Applied Graph Theory Summer 2006 "B" Session

Instructor: Steven Bellenot [bellenot@math.fsu.edu](mailto:bellenot@math.fsu.edu)  
<http://www.math.fsu.edu/~bellenot/class/su06/graph>  
 MTWRF 9:30-10:50 201 LOV



$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Adjacency Matrix

Petersen Graph --  $\{u, v\}$  adjacent to  $\{s, t\} \Leftrightarrow$  the sets are disjoint

$$\begin{bmatrix} 3 & 1 & 1 & 1 & 1 & 1 & -2 & -2 & -2 & -2 \\ 1 & 1 & 1 & 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 & -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & -1 \\ 1 & -1 & 0 & -1 & 0 & -1 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} & -1 \\ 1 & -1 & -1 & 0 & -1 & 0 & \frac{1}{2} & \frac{1}{2} & -\frac{3}{2} & -1 \\ 1 & -1 & -1 & -1 & -1 & -1 & -1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

The **Eigenvectors** (the column vectors) and **Eigenvalues** of the Adjacency Matrix.

Unlike the more theoretical brother MAD5305, MAD5932 Applied Graph Theory is designed more towards using than proving. Here Matlab eigenanalysis of the adjacency matrix is used to determine graph properties. For example, the largest eigenvalue/eigenvector says this graph is 3-regular. Often different graphs have different eigenvalues which will imply they are not isomorphic. Chemists use this feature.