

Regulating flows, topology of foliations and rigidity

Sérgio R. Fenley *

December 11, 2004

1 Introduction

In this article we study the topology of foliations in 3-manifolds as influenced by a transverse pseudo-Anosov flow. A Reebless foliation in a closed 3-manifold lifts to a foliation in the universal cover. It is very natural to ask what is the topology of the leaf space in the universal cover. The basic form of this is to check whether the leaf space is Hausdorff or not. The Reebless condition implies that this leaf space is always a one dimensional manifold [No]. Consequently if it is Hausdorff it will be homeomorphic to the real numbers \mathbf{R} . In this case we call the foliation a \mathbf{R} -covered foliation [Fe3, P14]. An immediate consequence of being \mathbf{R} -covered is that the fundamental group acts on \mathbf{R} and this action can be very useful [P13]. One important question addressed here is: when is a foliation \mathbf{R} -covered?

It turns out that in general it is very hard and subtle to check whether a foliation is \mathbf{R} -covered or not. Fibrations over the circle are always \mathbf{R} -covered and finite depth foliations [Ga1, Ga2] are never \mathbf{R} -covered unless the compact leaves are fibers of fibrations over the circle [Fe1].

For stable or unstable foliations of Anosov flows the question is much more complex and interesting. There is the groundbreaking work of Verjovsky [Ve], who was the first to analyse the topology of the lifts of the stable/unstable foliations of an Anosov flow and its important consequences. Suspension Anosov flows (suspensions) and geodesic flows in the unit tangent bundle of closed surfaces of negative curvature (geodesic flows) have stable/unstable foliations which are \mathbf{R} -covered. Anosov flows in Seifert fibered spaces are completely understood, due to the work of Ghys [Gh]. He proved that the stable foliation of the flow is always \mathbf{R} -covered and this was an essential step to show that such flows are always topologically conjugate to geodesic flows. Anosov flows in graph manifolds are also well understood due to the work of Barbot [Ba1, Ba3, Ba4] and there are many \mathbf{R} -covered examples. Again the \mathbf{R} -covered property for the stable foliation had important consequences. Also there are many non \mathbf{R} -covered examples in graph manifolds. In hyperbolic manifolds there are many examples of \mathbf{R} -covered Anosov foliations obtained by Dehn surgery on closed orbits of suspensions or geodesic flows [Fe3]. This has strong consequences for the geometry of the flows in hyperbolic manifolds, namely the study of the quasigeodesic property for the flow lines – when the stable foliation is \mathbf{R} -covered then the flow is not a quasigeodesic flow [Fe3]. In both graph manifolds and hyperbolic manifolds, the proof of the \mathbf{R} -covered property involves a very tricky analysis - for hyperbolic manifolds this entailed the study of how Dehn surgery modified the topology of the leaf space in the universal cover [Fe3]. The results were obtained under very controlled circumstances and in general Dehn surgery on suspensions may produce non \mathbf{R} -covered Anosov foliations - see [Fe5] for non \mathbf{R} -covered examples where the surgered manifold is hyperbolic.

Another class of \mathbf{R} -covered foliations is that of “slitherings”. A slithering of a manifold over the circle is a fibration of the universal cover of the manifold over the circle which is equivariant under group translations and produces a foliation in the manifold [Th5]. By construction these are all \mathbf{R} -covered. Fibrations over the circle are examples of slitherings, as are stable foliations of a large class of Anosov flows [Th5, Fe3]. Furthermore there are several techniques to construct or replicate slitherings, for instance branched covering techniques [Th5, Ca1]. This makes slitherings very common.

Finally there are also some new constructions by Calegari [Ca1] of \mathbf{R} -covered foliations in hyperbolic 3-manifolds which are not associated to slitherings. Another important fact is that for a general foliation in a closed, hyperbolic 3-manifold, the \mathbf{R} -covered property implies that every leaf in the universal cover has limit set which is the entire sphere at infinity.

On the other hand, the failure of the \mathbf{R} -covered property for the stable/unstable foliations of an Anosov flow in a 3-manifold also has strong consequences for the dynamics of the flow and the topology of the manifold [Fe4, Fe5]. In this case the general structure of such leaf spaces is not yet well understood.

*Research partially supported by NSF grants DMS-9612317 and DMS-0071683.

All of these examples demonstrate the importance of knowing whether a foliation is \mathbf{R} -covered or not. When \mathcal{F} has a compact leaf B , then \mathcal{F} is \mathbf{R} -covered if and only if B is a virtual fiber of M over the circle [Go-Sh]. However when \mathcal{F} has no compact leaves, it is very difficult to analyse the \mathbf{R} -covered property and very little is known beyond the examples mentioned above.

In this article we analyse this question when there is a transverse pseudo-Anosov flow. First we mention various classes of examples where there is a pseudo-Anosov flow transverse to the foliation. 1) If the manifold fibers over the circle with pseudo-Anosov monodromy, then there is a pseudo-Anosov flow transverse to the fibration [Th3, Bl-Ca]. This flow gives substantial information about the manifold and the foliation [Th2, Th4, Ca-Th]. 2) Another example is the situation of finite depth foliations: there are compact leaves, all leaves are properly embedded and the foliation strongly reflects a topological hierarchy of the manifold [Ga1, Ga2]. Gabai showed that if the manifold is orientable, irreducible and has non trivial second homology, then it has Reebless finite depth foliations [Ga1, Ga2]. Work of Gabai and Mosher [Mo2] shows that, as in the fibering case, there is a pseudo-Anosov flow transverse to the foliation (in general the flow is “almost” transverse to the foliation – a technical condition). The stable/unstable laminations obtained by splitting the singular leaves of the stable/unstable singular foliations of the flow are used to prove important results on most Dehn surgeries on knot complements in \mathbf{S}^3 [Mo2]. 3) Work of Roberts [Rob1, Rob2] implies that if γ is a knot in \mathbf{S}^3 with hyperbolic complement, then there are infinitely many Dehn surgeries on γ yielding closed manifolds with foliations transverse to pseudo-Anosov flows. 4) About two years ago, Thurston showed that a foliation induced by a slithering of the manifold over the circle is either reducible or admits a transverse pseudo-Anosov flow [Th5]. 5) We recently extended this result to general \mathbf{R} -covered foliations:

Theorem ([Fe10]) - Let \mathcal{F} be a transversely oriented, \mathbf{R} -covered foliation in M^3 closed, aspherical. Then either there is a $\mathbf{Z} \oplus \mathbf{Z}$ subgroup of $\pi_1(M)$ or there is a pseudo-Anosov flow transverse to \mathcal{F} . In particular the second option occurs if M is hyperbolic.

This result was independently proved by Calegari [Ca2]. 6) Finally Calegari has shown the same result for foliations with one sided branching [Ca3] and minimal foliations [Ca4].

Pseudo-Anosov flows transverse to foliations are amenable to branched cover and cut and paste techniques. This will be explored in this article to analyse 2 classes of examples. See also the wealth of examples created by Cooper, Long and Reid [CLR1, CLR2] and the author [Fe7] for compact immersed surfaces transverse to pseudo-Anosov flows. Thurston proposed that the existence of a pseudo-Anosov flow transverse to a foliation is a very general situation [Th6, Th7].

Let then Φ be a pseudo-Anosov transverse to a foliation \mathcal{F} . The question considered here is: can the transverse pseudo-Anosov flow be used to decide whether \mathcal{F} is \mathbf{R} -covered or not? First consider the following: does an arbitrary flow line in the universal cover intersect every leaf of the lifted foliation? If this happens we say that Φ *regulates* \mathcal{F} [Th5], or that Φ is a *regulating flow*. Clearly if that is the case, then the foliation is \mathbf{R} -covered – points in the given flow line parametrize the leaf space as homeomorphic to \mathbf{R} . Our goal is to analyse the converse: does \mathcal{F} being \mathbf{R} -covered imply that Φ is regulating for \mathcal{F} ? Surprisingly this is not true in general and there are easy counterexamples. However we prove that the converse only fails for some very restricted and well understood situations.

Let $\mathcal{F}^s, \mathcal{F}^u$ be the singular stable/unstable foliations associated to the Anosov flow Φ . A pseudo-Anosov flow is called \mathbf{R} -covered if one of its stable or unstable foliations is \mathbf{R} -covered [Ba1, Fe3]. This implies there are no p -prong orbits and also that both foliations are \mathbf{R} -covered [Ba1, Fe3]. Given a pseudo-Anosov flow Φ transverse to a foliation \mathcal{F} in M , let $\tilde{\Phi}$ be the lifted flow to the universal cover \tilde{M} and let $\tilde{\mathcal{F}}$ be the lifted foliation. We first study basic properties of foliations transverse to pseudo-Anosov flows.

Theorem A - Let Φ be a pseudo-Anosov flow transverse to a foliation \mathcal{F} in M^3 closed. Then \mathcal{F} is Reebless. In \tilde{M} , a given orbit of $\tilde{\Phi}$ intersects a leaf of $\tilde{\mathcal{F}}$ at most once.

It does not follow that \mathcal{F} is taut [Ga1], see constructions in section 3. Then we prove the main result:

Main theorem - Let Φ be a pseudo-Anosov flow transverse to a foliation \mathcal{F} in M^3 closed. Suppose that Φ is not an \mathbf{R} -covered pseudo-Anosov flow. Then Φ is regulating if and only if \mathcal{F} is an \mathbf{R} -covered foliation. There are counterexamples when Φ is an \mathbf{R} -covered pseudo-Anosov flow.

The rough idea is as follows: as \mathcal{F} is \mathbf{R} -covered, there is a unique way to move transversally to $\tilde{\mathcal{F}}$ in \tilde{M} . When Φ is not regulating for \mathcal{F} , this unique transversal direction to $\tilde{\mathcal{F}}$ will force the creation of a unique transversal direction to say the stable foliation $\tilde{\mathcal{F}}^s$ of $\tilde{\Phi}$. This in turn implies that \mathcal{F}^s has to be \mathbf{R} -covered and hence Φ is an \mathbf{R} -covered Anosov flow. The fundamental tool used here is the topological theory of pseudo-Anosov flows as developed in [Fe5, Fe8].

When Φ is an \mathbf{R} -covered Anosov flow, counterexamples to the main theorem can be obtained as follows: let \mathcal{F}

be the stable foliation of Φ – which is \mathbf{R} -covered. Perturb Φ to nearby Φ' which is transverse to \mathcal{F} . Then Φ' will not be regulating for \mathcal{F} . This is explained in detail in section 5.

The main theorem is a useful tool to analyse the topology of foliations:

Theorem B - *There is an infinite family of examples of non \mathbf{R} -covered foliations obtained by branched covers of foliations transverse to suspension Anosov flows and to skewed \mathbf{R} -covered Anosov flows. The resulting foliations are transverse to pseudo-Anosov flows. There are also infinitely many examples obtained by cut and paste operations on fibrations over the circle.*

The skewed condition is explained in detail in section 5.

The main theorem leaves open the situation where Φ is an \mathbf{R} -covered Anosov flow. In this case we prove a strong topological rigidity. We do this by characterizing all the possible foliations transverse to such a flow. A foliation \mathcal{F}_1 is a *blow up* of another foliation \mathcal{F}_2 if \mathcal{F}_1 is obtained by blowing up a collection of leaves of \mathcal{F}_2 into foliated I -bundles. We prove:

Theorem C (rigidity theorem) - *Let Φ be an \mathbf{R} -covered pseudo-Anosov flow transverse to a foliation \mathcal{F} . Then \mathcal{F} is \mathbf{R} -covered. In addition if Φ is not regulating for \mathcal{F} , then \mathcal{F} is topologically conjugate to a blow up of either the stable or unstable foliation of Φ .*

This topological rigidity result says that the only counterexamples to the main theorem are essentially obtained by the construction mentioned above and detailed in section 6. The proof of this rigidity result takes about a third of the article and is left to the end. As usual with rigidity results it uses tools from a variety of sources. Here we need: the structure of \mathbf{R} -covered Anosov foliations, topology of foliations, Gromov negatively curved spaces, quasi-isometries of metric spaces, dynamics and topological conjugacy of flows, amongst others. The proof is achieved by producing a new flow which is topologically conjugate to the original flow and which is tangent to a blow down of the foliation \mathcal{F} .

The results of this article have several applications. In a subsequent article we analyse the atoroidal property for manifolds with pseudo-Anosov flows and prove:

Theorem ([Fe9]) *Let Φ be a pseudo-Anosov flow in M^3 closed. Let $\mathbf{A} = \mathbf{Z} \oplus \mathbf{Z}$ be a subgroup of the fundamental group $\pi_1(M)$. Then \mathbf{A} does not act freely in the orbit space of $\tilde{\Phi}$, unless Φ is topologically conjugate to a suspension Anosov flow. This means that some non trivial element of \mathbf{A} leaves invariant an orbit of $\tilde{\Phi}$. Hence if Φ is not topologically conjugate to a suspension Anosov flow, then \mathbf{A} represents a free homotopy from a closed orbit of Φ to itself.*

Together with the results here one obtains an important relationship with the topology of the underlying manifold:

Theorem ([Fe9]) *Let Φ be a pseudo-Anosov flow transverse to an \mathbf{R} -covered foliation \mathcal{F} in M^3 closed. Then M is atoroidal, unless Φ is a non regulating \mathbf{R} -covered pseudo-Anosov flow. The condition is necessary as simple examples show.*

We can also apply the results here to obtain topological rigidity of pseudo-Anosov flows transverse to foliations. This means characterizing all pseudo-Anosov flows transverse to a given foliation.

Theorem ([Fe11]) *Let Φ be a regulating pseudo-Anosov flow transverse to an \mathbf{R} -covered foliation \mathcal{F} in closed M^3 . Then up to topological conjugacy, Φ is the only regulating pseudo-Anosov flow transverse to \mathcal{F} .*

The article is organized as follows: the next section contains general results about foliations transverse to pseudo-Anosov flows. Section 3 contains a review of the topological theory of pseudo-Anosov flows as needed here. In section 4 we prove the main theorem and in the next section we produce the counterexamples in the case of \mathbf{R} -covered Anosov flows. In section 6 we use the main theorem to study the topology of some classes of foliations. In the last section we show that the counterexamples to the main theorem can only occur in a very restricted setting.

We thank Bill Thurston for enlightening and very enjoyable conversations about foliations and flows. The main result in this article was motivated by a question asked by him. He also suggested that we analyse the branched cover examples of section 7.

2 Foliation transverse to flows

Pseudo-Anosov flows are a generalization of suspension flows of pseudo-Anosov surface homeomorphisms. These flows behave much like Anosov flows, but they may have finitely many singular orbits which are periodic and have




Figure 1: *a. Non Hausdorff behavior in the leaf space of $\widetilde{\mathcal{F}}^s$: (a1) F, L non separated from each other, as seen in \widetilde{M} , (a2) the corresponding picture in the leaf space \mathcal{H}^s ; (b) A singular leaf of $\widetilde{\mathcal{F}}^s$: (b1) as seen in \widetilde{M} , (b2) as seen in \mathcal{H}^s .*

a prescribed behavior.

Let Φ be a flow on a closed, oriented 3-manifold M . We say that Φ is a *pseudo-Anosov flow* if the following are satisfied:

- For each $x \in M$, the flow line $t \rightarrow \Phi(x, t)$ is C^1 , it is not a single point, and the tangent vector bundle $D_t\Phi$ is C^0 .
- There is a finite number of periodic orbits $\{\gamma_i\}$, called *singular orbits*, such that the flow is “topologically” smooth off of the singular orbits (see below).
- The flowlines are tangent to two singular transverse foliations $\mathcal{F}^s, \mathcal{F}^u$ which have smooth leaves off of γ_i and intersect exactly in the flow lines of Φ . These are like Anosov foliations off of the singular orbits. This is the topologically smooth behavior described above. A leaf containing a singularity is homeomorphic to $P \times I/f$ where p is a p -prong in the plane and f is a homeo from $I \times \{1\}$ to $I \times \{0\}$.
- In a stable leaf all orbits are forward asymptotic, in an unstable leaf all orbits are backwards asymptotic.

Notation/definition: The lifted singular foliations to \widetilde{M} are denoted by $\widetilde{\mathcal{F}}^s, \widetilde{\mathcal{F}}^u$. If $x \in M$ let $W^s(x)$ denote the leaf of \mathcal{F}^s containing x . Similarly one defines $W^u(x)$ and in the universal cover $\widetilde{W}^s(x), \widetilde{W}^u(x)$. Similarly if α is an orbit of Φ define $W^s(\alpha)$, etc... Let also $\widetilde{\Phi}$ be the lifted flow to \widetilde{M} .

In figure 1 we highlight the difference between non Hausdorff behavior in the leaf space of $\widetilde{\mathcal{F}}^s$ and the splitting (or branching) of leaves associated to singular orbits of $\widetilde{\Phi}$. In part (a) the leaves F, L of $\widetilde{\mathcal{F}}^s$ are not separated from each other in the leaf space of $\widetilde{\mathcal{F}}^s$. Notice that the sequence F_i converges to F and L . In fig 1 part (b) we sketch a singular leaf S with 3 prongs. Even though S separates \widetilde{M} into 3 or more regions, non Hausdorffness is not involved. The leaves S_i converge only to S . In this article, unless otherwise specified, all pictures of leaves of $\widetilde{\mathcal{F}}^s, \widetilde{\mathcal{F}}^u$ will describe them as subsets of \widetilde{M} , rather than in the leaf space of $\widetilde{\mathcal{F}}^s$.

Let Φ be a pseudo-Anosov flow in a closed 3-manifold M . The following facts and definitions will be needed. We proved in [Fe-Mo] that the orbit space of Φ in \widetilde{M} is homeomorphic to the plane \mathbf{R}^2 . This orbit space is denoted by $\mathcal{O} \cong \widetilde{M}/\widetilde{\Phi}$. Let $\Theta: \widetilde{M} \rightarrow \mathcal{O} \cong \mathbf{R}^2$ be the projection map. For any subset $B \subset \mathcal{O}$, let $B \times \mathbf{R} \subset \widetilde{M}$, be the set $\Theta^{-1}(B)$. As the foliations $\widetilde{\mathcal{F}}^s, \widetilde{\mathcal{F}}^u$ are invariant under $\widetilde{\Phi}$, they induce singular, transverse 1-dim foliations $\widetilde{\mathcal{F}}_{\mathcal{O}}^s, \widetilde{\mathcal{F}}_{\mathcal{O}}^u$ in \mathcal{O} . The singular points of $\widetilde{\mathcal{F}}_{\mathcal{O}}^u$ are the same as those of $\widetilde{\mathcal{F}}_{\mathcal{O}}^s$. If L is a leaf of $\widetilde{\mathcal{F}}^s$ or $\widetilde{\mathcal{F}}^u$, then $\Theta(L) \subset \mathcal{O}$ is a tree which is either homeomorphic to \mathbf{R} if L is regular, or is a union of p -rays all with the same starting point if L has a singular p -prong orbit. In particular every orbit in L disconnects L .

In this section we develop some preliminaries of foliations transverse to pseudo-Anosov flows. Let \mathcal{F} be a foliation transverse to a pseudo-Anosov flow Φ . Then $\mathcal{F}, \mathcal{F}^s, \mathcal{F}^u$ are 3 foliations in M which are transverse to each other everywhere.

Proposition 2.1. *Let Φ be a pseudo-Anosov flow transverse to a foliation \mathcal{F} in M^3 . Then \mathcal{F} is Reebless. In addition every orbit of $\widetilde{\Phi}$ intersects a leaf of \mathcal{F} at most once.*

Proof. Suppose that \mathcal{F} is not Reebless and first consider a Reeb component which is a solid torus V bounded by a torus T . Assume the flow is incoming along T . Since orbits are trapped inside once they enter V , the shadow lemma for pseudo-Anosov flows [Han, Man, Mo1], shows that there is a periodic orbit γ in $V - T$. Then $W^u(\gamma)$ is contained in $V - T$, because the flow is incoming along T . Since γ^n is not null homotopic for any $n \neq 0$ [Mo1], then \tilde{V} is homeomorphic to $D^2 \times \mathbf{R}$. It now follows that $\tilde{W}^u(\gamma)$ is not properly embedded in the universal cover, contradiction [Ga-Oe]. The same proof shows that there cannot be non-orientable Reeb components - that is the quotient of a Reeb component by an orientation reversing involution of V which preserves the meridian. This can be seen by lifting to an orientation double cover of M : the lifted flow is pseudo-Anosov transverse to a Reeb component.

As for the second statement: Since \mathcal{F} is Reebless and transverse to a pseudo-Anosov flow, then leaves of $\tilde{\mathcal{F}}$ are properly embedded topological planes in \tilde{M} , because closed transversals to \mathcal{F} are not null homotopic [No]. Therefore leaves of $\tilde{\mathcal{F}}$ separate \tilde{M} . Since orbits of $\tilde{\Phi}$ are transverse to $\tilde{\mathcal{F}}$, an orbit of $\tilde{\Phi}$ can intersect a leaf of $\tilde{\mathcal{F}}$ at most once. \square

Suppose that Φ is not regulating. Let $\Theta : \tilde{M} \rightarrow \mathcal{O}$ be the projection. There is γ orbit of $\tilde{\Phi}$ not intersecting all leaves of $\tilde{\mathcal{F}}$ and a leaf F of $\tilde{\mathcal{F}}$ not intersecting all orbits of $\tilde{\Phi}$. Hence $\Theta(F)$ is not all of \mathcal{O} . The following result describes the boundary of $\Theta(F) \subset \mathcal{O}$.

If $L \in \tilde{\mathcal{F}}^s$ then a *sector* of L is a component of $\tilde{M} - L$. A *line leaf* of L is the boundary L' of a sector of L . The line leaf L' is *regular* on the side which is a sector of L . It is regular on both sides if and only if L is a regular leaf.

Proposition 2.2. $\partial\Theta(F) \subset \mathcal{O}$ is a disjoint union of line leaves of $\tilde{\mathcal{F}}_{\mathcal{O}}^s, \tilde{\mathcal{F}}_{\mathcal{O}}^u$, all regular on the side containing $\Theta(F)$. Also if $L \in \tilde{\mathcal{F}}^s$ (or $\tilde{\mathcal{F}}^u$), then $F \cap L$ is connected.

Proof. Because Φ is transverse to \mathcal{F} and M is compact, then for $\epsilon > 0$ sufficiently small there is $\eta(\epsilon)$ with $\eta(\epsilon) \rightarrow 0$, when $\epsilon \rightarrow 0$ so that: any orbit of $\tilde{\Phi}$ which comes within ϵ of a point $z \in F \in \tilde{\mathcal{F}}$ will in fact intersect F within $\eta(\epsilon)$ of z . Let $p \in \partial\Theta(F)$ and let $p_i \in \Theta(F)$ with $p_i \rightarrow p$. Let $z \in \tilde{M}$ with $\Theta(z) = p$. Consider $D \subset \tilde{M}$ a small embedded disk, transverse to $\tilde{\Phi}$ with $z \in \text{int}(D)$ and Θ injective in D . Let $w_i \in F$ with $\Theta(w_i) = p_i$. By truncating finitely many terms if necessary, there are unique $z_i \in D$ and $t_i \in \mathbf{R}$ so that $w_i = \tilde{\Phi}_{t_i}(z_i)$. If $|t_i| \not\rightarrow +\infty$ assume up to subsequence that $t_i \rightarrow t_0$. It follows that $w_i \rightarrow \tilde{\Phi}_{t_0}(z)$. But since F is closed in \tilde{M} , then $\tilde{\Phi}_{t_0}(z) \in F$, contradicting $p \notin \Theta(F)$.

Assume then that there is a subsequence $t_i \rightarrow +\infty$. Suppose that the corresponding p_i are all in the closure of the same sector defined by $m = \Theta(\tilde{W}^s(z))$ at p . Let l be the line leaf of m which bounds this sector.

Claim - l is a boundary leaf of the projection of F , that is, $l \subset \partial\Theta(F)$, see fig. 2, b.

Let $v \in \tilde{M}$ with $\Theta(v) \in l$. Choose β a small segment in $\tilde{W}^u(v)$ transverse to the flow $\tilde{\Phi}$ in $\tilde{W}^u(v)$ and with v in the interior of β . For i big enough let $q_i = \beta \cap \tilde{W}^s(w_i)$. The intersection is non void because l is regular on the side containing p_i . There are

$$s_i \in \mathbf{R}, \quad s_i \rightarrow +\infty \quad \text{so that} \quad \tilde{\Phi}_{s_i}(q_i) \in \tilde{W}^s(w_i) \quad \text{and} \quad d(\tilde{\Phi}_{s_i}(q_i), w_i) \rightarrow 0.$$

This implies that there are $\epsilon_i \rightarrow 0$ with $\tilde{\Phi}_{s_i+\epsilon_i}(q_i) \in F$. Hence for i big enough $\Theta(q_i) \in \Theta(F)$. In fact this shows that a segment α in $\tilde{W}^s(q_i)$ transverse to $\tilde{\Phi}$ in $\tilde{W}^s(q_i)$ and going from $\tilde{\Phi}_{s_i}(q_i)$ to w_i projects into $\Theta(F)$. One concludes that

$$\Theta(v) \in \Theta(F) \cup \partial\Theta(F).$$

If $\Theta(v) \in \Theta(F)$ let E be a small disk contained in F with v in the interior. Hence for i big enough there are bounded r_i with $\tilde{\Phi}_{r_i}(q_i) \in E$. But the argument above shows that there are $\tilde{\Phi}_{s_i+\epsilon_i}(q_i) \in F$ with $s_i + \epsilon_i \rightarrow +\infty$ as $i \rightarrow +\infty$. This would produce two points $\tilde{\Phi}_{r_i}(q_i)$ and $\tilde{\Phi}_{s_i+\epsilon_i}(q_i)$ of F in $\tilde{\Phi}_{\mathbf{R}}(q_i)$, contradiction.

We conclude that the stable line leaf $l \subset \partial\Theta(F)$, showing the claim.

Similarly given the t_i defined above, if there is some subsequence $t_{i_n} \rightarrow -\infty$, then we have an unstable line leaf I through p with $I \subset \partial\Theta(F)$. Since $\Theta(F)$ is connected this would force $I \cap \Theta(F) \neq \emptyset$ which is impossible. In addition l is regular on the side containing F . This proves the first statement.

Let now $F \in \tilde{\mathcal{F}}$ and $L \in \tilde{\mathcal{F}}^s$. Suppose that $F \cap L$ has two components u_1, u_2 . Then $\Theta(u_1) \cap \Theta(u_2) = \emptyset$ and there is $x \in L$ with $\Theta(x)$ separating $\Theta(u_1)$ from $\Theta(u_2)$ in $\Theta(L)$ and $\Theta(x) \in \partial\Theta(u_1)$. By the proof above $\Theta(x) \in \partial\Theta(F)$ and there is a line leaf

$$m \subset \partial\Theta(F) \quad \text{with either} \quad m \subset \Theta(\tilde{W}^s(x)) \quad \text{or} \quad m \subset \Theta(\tilde{W}^u(x)).$$




Figure 2: a. Producing boundary in $\Theta(F)$, b. Walls in the back of F .

Since $\Theta(u_1) \subset \Theta(F)$, $\Theta(u_1) \subset \Theta(\widetilde{W}^s(x))$ and m is regular on the side containing $\Theta(F)$, then $m \subset \Theta(\widetilde{W}^s(x))$ cannot happen. Now $\Theta(u_1)$ and $\Theta(u_2)$ must be in the same component of $\mathcal{O} - m$ and $\Theta(x)$ separates $\Theta(u_1)$ from $\Theta(u_2)$ in $\Theta(F)$. Hence there is a component of $\Theta(\widetilde{W}^u(x)) - \Theta(x)$ contained in $\Theta(F)$ and separating $\Theta(u_1)$ from $\Theta(u_2)$ in $\Theta(\widetilde{W}^s(x))$. This contradicts the fact that m is a line leaf of $\Theta(\widetilde{W}^u(x))$ which is regular on the side containing $\Theta(F)$. The second statement follows. This finishes the proof of proposition 2.2. \square

Notice that \mathcal{F} is always transversely orientable because it is transverse to a non vanishing continuous vector field. Let positive flow direction define the positive transversal orientation to \mathcal{F} . This will be fixed throughout the article.

As leaves of $\widetilde{\mathcal{F}}$ separate \widetilde{M} , then given $F \in \widetilde{\mathcal{F}}$ we can define the *positive* or *front* side of F to be the component of $\widetilde{M} - F$ on the positive side of F with respect to the transversal orientation to $\widetilde{\mathcal{F}}$. Similarly define the *negative* or *back* side of F . Notice that if l is a stable boundary leaf of $\Theta(F)$ then $l \times \mathbf{R}$ is in the negative side of F , see fig. 2, b. In addition for any $z \in \widetilde{M}$ with $\Theta(z) \in l$ then $\widetilde{W}^u(z)$ intersects F . Similarly if l is an unstable boundary leaf of $\partial\Theta(F)$, then $l \times \mathbf{R}$ is on the positive side of F . This facts will be used throughout the article.

We stress that the fact that \mathcal{F} is transverse to a pseudo-Anosov flow does not imply that \mathcal{F} is taut, where taut means that every leaf intersects a closed, tranverse loop [Ga1]. For example this can be seen with the Franks-Williams example of an intransitive Anosov flow [Fr-Wi]. Start with a suspension Anosov flow Φ_0 in M_0 and do a DA operation on a closed orbit γ of Φ_0 . This produces a flow called derived from Anosov flow where γ is now a repelling orbit. Remove a solid torus neighborhood of γ obtaining a manifold M_1 with torus boundary T and an incoming semi-flow Φ_1 . Consider a homeomorphic copy M_2 of M_1 with a reversal flow Φ_2 which is outgoing along the boundary. Glue M_1 to M_2 along the boundary to produce closed manifold M with flow Φ . Franks and Williams show that Φ is Anosov and it is not transitive: it has a separating transverse torus.

Now construct the transverse foliation to Φ . Start with the fibration in M_0 which is transverse to Φ_0 . This produces a fibration in M_1 which is transverse to the boundary torus T . Spin the fibers around the boundary torus T and add T as a torus leaf so that the foliation is still transverse to the flow. Then M_1 has a foliation transverse to Φ_1 . Put a similar foliation in M_2 . The resulting foliation \mathcal{F} in M is transverse to the Anosov flow Φ but is not taut: it has a separating torus T . \mathcal{F} is a depth one foliation with a unique compact leaf T and is a fibration over the circle in any component of the complement of this compact leaf [Ga1]. In addition M fibers over the circle with fiber a surface of genus 2 and reducible monodromy.

In the example above notice that Φ is not regulating for \mathcal{F} , because there are closed orbits of Φ contained in M_1 . Their lifts cannot intersect any lift of a leaf of \mathcal{F} in M_2 .

Still there is much one can say if \mathcal{F} is not taut:

Proposition 2.3. *Let Φ be a pseudo-Anosov flow transverse to a foliation \mathcal{F} . If \mathcal{F} is not taut, then Φ is not regulating and \mathcal{F} is not \mathbf{R} -covered.*

Proof. Up to taking a finite cover of M , we may assume that M is orientable and \mathcal{F} is transversely orientable. If \mathcal{F} is not taut, then Goodman [Go] showed that there are finitely many toral leaves T_1, \dots, T_n in \mathcal{F} so that $T_1 \cup \dots \cup T_n$ bounds a submanifold V and the flow is (say) incoming V along T_1, \dots, T_n . Any orbit intersecting T_1 is trapped




Figure 3: a. Rectangles, b. Perfect fits in the universal cover.

inside V in forward time and hence can only limit inside V and any limit is an orbit γ entirely contained in V . Then $\gamma \cap T_i = \emptyset$ for any i , so in the universal cover any lift $\tilde{\gamma}$ will not intersect any lift of T_i . Therefore the flow Φ is not regulating for \mathcal{F} .

Now we show that \mathcal{F} is not \mathbf{R} -covered. By lemma 2.1, \mathcal{F} is Reebless. Suppose first that $n \geq 2$. There is no transversal to \mathcal{F} through T_1 and T_2 , hence no transversal through any lifts of T_1, T_2 to \tilde{M} . This shows that \mathcal{F} is not \mathbf{R} -covered. The remaining case is that $n = 1$ and T_1 bounds on both sides. Let V be the closure of a component of $M - T_1$. If $\pi_1(T_1)$ surjects in $\pi_1(V)$ then since \mathcal{F} is Reebless, $\pi_1(V) \cong \mathbf{Z} \oplus \mathbf{Z}$. This is impossible for a compact manifold V with torus boundary, see chapter 10 of [He]. If $\pi_1(T_1)$ does not surject in $\pi_1(V)$, there is a closed loop $\alpha \in V$ not homotopic into T_1 . Let g be the covering translation of \tilde{M} associated to α . In the universal cover $g(\tilde{T}_1) \neq \tilde{T}_1$. But $g(\tilde{T}_1), \tilde{T}_1$ do not cross a transversal, otherwise there would be a closed transversal to T_1 in M . We conclude that \mathcal{F} is not \mathbf{R} -covered. \square

3 Topological theory of pseudo-Anosov flows

Let Φ be a pseudo-Anosov flow in M^3 closed. We review the results about the topology of $\tilde{\mathcal{F}}^s, \tilde{\mathcal{F}}^u$ which will be used essentially in the next section. We refer to [Fe5, Fe8] for detailed proofs and explanations.

Definition 3.1. Let L be a leaf of $\tilde{\mathcal{F}}^s$ or a line leaf of a leaf of $\tilde{\mathcal{F}}^s$. Then a half leaf of L is a connected component A of $L - \gamma$, where γ is any full orbit in L . The closed half leaf is $\bar{A} = A \cup \gamma$ and its boundary is $\partial A = \gamma$. If ζ is an open, relatively compact, connected subset of $\Theta(L)$ then it defines a flow band L_1 of L by $L_1 = \Theta^{-1}(\zeta)$. Let \bar{L}_1 be the closure of L_1 in \tilde{M} . If ζ is an open segment in $\Theta(L)$, then $\Theta^{-1}(\zeta)$ is called a segment flow band of L .

Two important facts: First if $F \in \tilde{\mathcal{F}}^s$ and $G \in \tilde{\mathcal{F}}^u$ then F and G intersect in at most one orbit, a consequence of index computations for foliations in the plane. A second fact: Suppose that a leaf $F \in \tilde{\mathcal{F}}^s$ intersects two leaves $G, H \in \tilde{\mathcal{F}}^u$ and so does $L \in \tilde{\mathcal{F}}^s$. Then F, L, G, H define a rectangle in \tilde{M} , see fig. 3, a. There are no singularities of Φ in the interior of the rectangle and there is a product structure of $\tilde{\mathcal{F}}^s$ and $\tilde{\mathcal{F}}^u$ in the rectangle.

Definition 3.2. Perfect fits - Two leaves $F \in \tilde{\mathcal{F}}^s$ and $G \in \tilde{\mathcal{F}}^u$, form a perfect fit if $F \cap G = \emptyset$ and there are line leaves F_0, G_0 of F, G respectively and half leaves F_1 of F_0 and G_1 of G_0 and also segment flow bands $L_1 \subset L \in \tilde{\mathcal{F}}^s$ and $H_1 \subset H \in \tilde{\mathcal{F}}^u$, so that F_0 is regular on the side containing L , G_0 is regular on the side containing H and:

$$\begin{aligned} \bar{L}_1 \cap \bar{G}_1 &= \partial L_1 \cap \partial G_1, \quad \bar{L}_1 \cap \bar{H}_1 = \partial L_1 \cap \partial H_1, \quad \bar{H}_1 \cap \bar{F}_1 = \partial H_1 \cap \partial F_1, \\ &\text{with } \bar{L}_1 \cap \bar{G}_1 \neq \emptyset, \quad \bar{L}_1 \cap \bar{H}_1 \neq \emptyset \quad \text{and} \quad \bar{H}_1 \cap \bar{F}_1 \neq \emptyset. \end{aligned}$$

Furthermore

$$\forall S \in \tilde{\mathcal{F}}^u, \quad S \cap L_1 \neq \emptyset \Rightarrow S \cap F_1 \neq \emptyset \quad (1) \text{ and}$$

$$\forall E \in \tilde{\mathcal{F}}^s, \quad E \cap H_1 \neq \emptyset \Rightarrow E \cap G_1 \neq \emptyset \quad (2).$$

We refer to fig. 3, b for perfect fits. Implications (1), (2) imply equivalences (that is $S \cap L_1 \neq \emptyset \Leftrightarrow S \cap F_1 \neq \emptyset$ and the same for (2)). The set $\bar{F}_1 \cup \bar{H}_1 \cup \bar{L}_1 \cup \bar{G}_1$ separates \tilde{M} . Let A be the complementary region which does




Figure 4: a. A lozenge, b. A chain of lozenges.

not contain $F - F_1$ in its closure. Then there are no singularities of $\tilde{\Phi}$ in A . Therefore perfect fits produce “ideal” rectangles, in the sense that even though F and G do not intersect, there is a product structure (of $\tilde{\mathcal{F}}^s$ and $\tilde{\mathcal{F}}^u$) in the interior of A .

Definition 3.3. Given $p \in \tilde{M}$ (or $p \in \mathcal{O}$), and a half leaf H of $\tilde{W}^u(p)$ defined by $\tilde{\Phi}_{\mathbf{R}}(p)$, let

$$\mathcal{J}^u(H) = \{F \in \mathcal{H}(\tilde{\mathcal{F}}^s) \mid F \cap H \neq \emptyset\} \subset \mathcal{H}(\tilde{\mathcal{F}}^s).$$

Notice that $\tilde{W}^s(p) \notin \mathcal{J}^u(H)$. Let also

$$\mathcal{L}^u(H) = \bigcup \{p \in \tilde{M} \mid p \in F \in \mathcal{J}^u(H)\} \subset \tilde{M}.$$

Then $\mathcal{L}^u(H) \subset \tilde{M}$ and $\tilde{W}^s(p) \subset \partial \mathcal{L}^u(H)$. Similarly define $\mathcal{J}^s(L), \mathcal{L}^s(L)$ for a stable half leaf L .

Definition 3.4. Lozenges - Let $p, q \in \tilde{M}$ and half leaves L_p, H_p of $\tilde{W}^s(p), \tilde{W}^u(p)$ defined by $\tilde{\Phi}_{\mathbf{R}}(p)$, half leaves L_q, H_q of $\tilde{W}^s(q), \tilde{W}^u(q)$ defined by $\tilde{\Phi}_{\mathbf{R}}(q)$ so that:

$$\mathcal{L}^u(L_p) \cap \mathcal{L}^s(H_q) = \mathcal{L}^u(L_q) \cap \mathcal{L}^s(H_p) \subset \tilde{M}$$

Then this intersection is called a lozenge \mathcal{A} in \tilde{M} . The corners of the lozenge are $\tilde{\Phi}_{\mathbf{R}}(p)$ and $\tilde{\Phi}_{\mathbf{R}}(q)$ and \mathcal{A} is a subset of \tilde{M} . The sides of \mathcal{A} are L_p, H_p, L_q, H_q . The sides are not contained in the lozenge, but are in the boundary of the lozenge.

Sometimes we also refer to p and q as corners of the lozenge. There are no singularities in the lozenges. This in fact shows that \mathcal{A} is an open region in \tilde{M} . However there may be singular orbits on the sides of the lozenge and the corner orbits also may be singular. The definition of a lozenge implies that L_p, H_q form a perfect fit and so do L_q, H_p . This is an equivalent way to define a lozenge with corners $\tilde{\Phi}_{\mathbf{R}}(p), \tilde{\Phi}_{\mathbf{R}}(q)$. Given any four leaves there is at most one lozenge with sides in them, so we may refer to the full leaves as the sides of the lozenge.

Two lozenges are *adjacent* if they share a corner and there is a stable or unstable leaf intersecting both of them, see fig. 4 b. Therefore they share a side. A *chain of lozenges* is a collection $\{\mathcal{A}_i\}, i \in I$, where I is an interval (finite or not) in \mathbf{Z} ; so that if $i, i+1 \in I$, then \mathcal{A}_i and \mathcal{A}_{i+1} share a corner, see fig. 4 b. Consecutive lozenges may be adjacent or not. The chain is finite if I is finite. We say that two orbits γ, α of $\tilde{\Phi}$ (or the leaves $\tilde{W}^s(\gamma), \tilde{W}^s(\alpha)$) are connected by a chain of lozenges $\{\mathcal{A}_i\}, 1 \leq i \leq n$, if γ is a corner of \mathcal{A}_1 and α is a corner of \mathcal{A}_n .

Definition 3.5. Suppose $\zeta \subset E \in \tilde{\mathcal{F}}^s$ is a (possibly infinite) strong stable segment so that for each $p \in \zeta$ there is a half leaf H_p of $\tilde{W}^u(p)$ defined by $\tilde{\Phi}_{\mathbf{R}}(p)$ so that

$$\forall p, q \in \zeta, \quad \mathcal{J}^u(H_p) = \mathcal{J}^u(H_q). \quad \text{In that case let } \mathcal{P} = \bigcup_{p \in \zeta} H_p.$$

Then $\mathcal{P} \subset \tilde{M}$ is called an unstable product region with base segment ζ . The base segment is not uniquely determined by \mathcal{P} . Similarly define stable product regions.




Figure 5: The correct picture between non separated leaves of $\tilde{\mathcal{F}}^s$.

A fundamental result is the following:

Theorem 3.6. [Fe5, Fe8] *Let Φ be a pseudo-Anosov flow in M^3 closed and let $F_0 \neq F_1 \in \tilde{\mathcal{F}}^s$. Suppose that there is a non trivial covering translation g with $g(F_i) = F_i, i = 0, 1$. Let $\alpha_i, i = 0, 1$ be the periodic orbits of $\tilde{\Phi}$ in F_i so that $g(\alpha_i) = \alpha_i$. Then α_0 and α_1 are connected by a finite chain of lozenges $\{A_i\}, 1 \leq i \leq n$ and g leaves invariant each lozenge A_i as well as their corners.*

A leaf L of $\tilde{\mathcal{F}}^s$ or $\tilde{\mathcal{F}}^u$ is called *periodic* if there is a non trivial covering translation g of \tilde{M} with $g(L) = L$. This is equivalent to $\pi(L)$ having a periodic orbit of Φ . In the same way an orbit γ of $\tilde{\Phi}$ is *periodic* if $\pi(\gamma)$ is a periodic orbit of Φ . The main result concerning non Hausdorff behavior in the leaf spaces of $\tilde{\mathcal{F}}^s, \tilde{\mathcal{F}}^u$ is the following:

Theorem 3.7. [Fe5, Fe8] *Let Φ be a pseudo-Anosov flow in M^3 . Suppose that $F \neq L$ are not separated in the leaf space of $\tilde{\mathcal{F}}^s$. Then F is a periodic leaf and so is L . Let F_0, L_0 be the line leaves of F, L which are not separated from each other. Let V_0 be the sector of F bounded by F_0 and containing L . Let α be the periodic orbit in F_0 and H_0 be the component of $(\tilde{W}^u(\alpha) - \alpha)$ contained in V_0 . Let g be a non trivial covering translation with $g(F_0) = F_0, g(H_0) = H_0$ and g leaves invariant the components of $(F_0 - \alpha)$. Then $g(L_0) = L_0$. This produces closed orbits of Φ which are freely homotopic in M . Theorem 3.6 then implies that F_0 and L_0 are connected by a finite chain of lozenges $\{A_i\}, 1 \leq i \leq n$, all contained in $\mathcal{L}^u(H_0)$ and all intersecting a common stable leaf C . Consecutive lozenges are adjacent. There is an even number of lozenges in the chain, see fig. 5.*

This result does not assume that M is toroidal or atoroidal. In fact almost all Anosov flows satisfy this condition [Fe8].

Finally we have the following result.

Theorem 3.8. [Fe5, Fe8] *Let Φ be a pseudo-Anosov flow in M^3 . If there is a product region in \tilde{M} , then Φ is an \mathbf{R} -covered pseudo-Anosov flow. In particular $\mathcal{F}^s, \mathcal{F}^u$ do not have singularities. Furthermore any leaf of $\tilde{\mathcal{F}}^s$ intersects every leaf of $\tilde{\mathcal{F}}^u$ and vice versa.*

4 Regulating flows

In this section we prove the main result of the article. First we obtain a useful property.

Proposition 4.1. *Let Φ be a pseudo-Anosov flow transverse to an \mathbf{R} -covered foliation \mathcal{F} and not regulating. Then for any leaf G of $\tilde{\mathcal{F}}$ there is an orbit γ of $\tilde{\Phi}$ not intersecting G . Conversely for any orbit α of $\tilde{\Phi}$ there is a leaf E of $\tilde{\mathcal{F}}$ not intersected by α .*

Proof. Since Φ is not regulating for \mathcal{F} , let β be orbit of $\tilde{\Phi}$ and H leaf of $\tilde{\mathcal{F}}$, so that $\beta \cap H = \emptyset$. Without loss of generality assume that β is in the back of H . Since \mathcal{F} is Reebless, \mathbf{R} -covered and transverse to a pseudo-Anosov flow, proposition 2.3 implies that \mathcal{F} is taut. Then we can choose finitely many closed transversals so that their union intersects every leaf of \mathcal{F} . Furthermore concatenating these transversals one finds a single closed transversal δ which intersects every leaf of \mathcal{F} .

Let now G be any leaf of $\tilde{\mathcal{F}}$. Then δ intersects $\pi(H)$ and $\pi(G)$, leaves of \mathcal{F} . Lifting to the universal cover we produce a transversal arc to $\tilde{\mathcal{F}}$ going from H to $g(G)$ in the positive direction. Here $g(G)$ is a covering translate of G . Then $g(G)$ is in the front of H and δ is in the back of H , hence $g(G) \cap \delta = \emptyset$. Consequently $G \cap g^{-1}(\delta) = \emptyset$. This proves the first assertion of the proposition.

As for the second assertion. Since $\Theta(H)$ does not contain $\Theta(\delta)$, then by proposition 2.2, there is a line leaf l in \mathcal{O} , boundary component of $\Theta(H)$ with $l \times \mathbf{R}$ either containing δ or separating δ from H . Since δ is in the back of H , then proposition 2.2 shows that l is a line leaf of $\tilde{\mathcal{F}}_{\mathcal{O}}^s$. Let L be a non singular leaf of $\tilde{\mathcal{F}}^s$, close to $l \times \mathbf{R}$ and not intersecting H . Let W_1 be the component of $\tilde{M} - L$ not containing H .

Claim - $\pi(W_1) = M$.

Otherwise $\pi(W_1)$ is open, \mathcal{F}^s saturated and not M . Let Z be a leaf of \mathcal{F}^s in the boundary of $\pi(W_1)$. Since Z is not compact it limits on a regular point p . Locally near p one finds many distinct local sheets of Z . The definition of W_1 implies that in fact Z intersects $\pi(W_1)$. But then $Z \subset \pi(W_1)$, contradiction. This proves the claim.

Therefore given an arbitrary orbit α of $\tilde{\Phi}$ let h be a covering translation of \tilde{M} with $h(\alpha) \subset W_1$. Then $h(\alpha) \cap H = \emptyset$, so $\alpha \cap h^{-1}(H) = \emptyset$. This finishes the proof of the proposition. \square

We now prove the main theorem of the article:

Theorem 4.2. *Let Φ be a pseudo-Anosov flow transverse to \mathcal{F} , with Φ not an \mathbf{R} -covered Anosov flow. Then Φ is regulating for \mathcal{F} if and only if \mathcal{F} is \mathbf{R} -covered.*

Proof. If Φ is regulating for \mathcal{F} then clearly \mathcal{F} is \mathbf{R} -covered, because any orbit of $\tilde{\Phi}$ provides a homeomorphism from its points to the leaf space of $\tilde{\mathcal{F}}$. Suppose from now on that \mathcal{F} is \mathbf{R} -covered but Φ is not regulating for \mathcal{F} . We will show that Φ must be an \mathbf{R} -covered Anosov flow.

Put an order in the leaf space of $\tilde{\mathcal{F}}$ as follows: $F <_s G$ if F is in the back of G . The goal is to show that there is a unique transversal direction to $\tilde{\mathcal{F}}^s$ or $\tilde{\mathcal{F}}^u$.

1. General construction for periodic orbits of $\tilde{\Phi}$ -

For each periodic orbit of $\tilde{\Phi}$ we produce a chain of leaves of $\tilde{\mathcal{F}}$ escaping in $\tilde{\mathcal{F}}$ and associated to that we produce a chain of leaves of $\tilde{\mathcal{F}}^s$ escaping in $\tilde{\mathcal{F}}^s$. This defines a direction transverse to $\tilde{\mathcal{F}}^s$, which we later prove is unique.

Let then γ be a periodic orbit of $\tilde{\Phi}$, with $g(\gamma) = \gamma$ and $g \in \pi_1(M)$ a generator of the isotropy group of γ . By the previous proposition, γ does not intersect every leaf of $\tilde{\mathcal{F}}$, so let $F' \in \tilde{\mathcal{F}}$ with $F' \cap \gamma = \emptyset$. For simplicity assume that γ is on the back side of F' . Since there are leaves of $\tilde{\mathcal{F}}$ intersecting γ and $\tilde{\mathcal{F}}$ is \mathbf{R} -covered, there is a unique leaf F with $F \cap \gamma = \emptyset$, γ in the back of F and for any $E \in \tilde{\mathcal{F}}$ sufficiently close to F and in the back of F then $E \cap \gamma \neq \emptyset$. In other words F is the least element in $\tilde{\mathcal{F}}$ satisfying $F \cap \gamma = \emptyset$ and γ is in the back of F . For the existence and uniqueness of F , it is essential the fact that \mathcal{F} is \mathbf{R} -covered as this may not be true otherwise. Taking g^{-1} if necessary we may assume that $g(F) \leq_s F$. Since $g(\gamma) = \gamma$ and $g(F) \cap \gamma = \emptyset$, the definition of F implies that $g(F) = F$.

Since $F \cap \gamma = \emptyset$ and γ is in the back of F , there is a stable line leaf l of $\tilde{\mathcal{F}}_{\mathcal{O}}^s$ with $l \subset \partial\Theta(F)$ and either $\gamma \subset l \times \mathbf{R}$ or $l \times \mathbf{R}$ separating γ from F . By the same reasoning as above $g(l \times \mathbf{R}) = l \times \mathbf{R}$. Let $L = l \times \mathbf{R}$ and V be the component of $\tilde{M} - L$ not containing F .

The following technical fact is needed later:

Claim 1.1 - There is F_1 in $\tilde{\mathcal{F}}$ with γ in the back of F_1 and $\tilde{W}^u(\gamma) \cap F_1 = \emptyset$.

In order to prove that we go in the opposite direction to F , that is in the region V .

Suppose first that L intersects all leaves of $\tilde{\mathcal{F}}^u$ which intersect V . In particular the leaf space of $\tilde{\mathcal{F}}^u$ restricted to V is homeomorphic to \mathbf{R} and therefore V does not contain singularities of $\tilde{\Phi}$. Since $\pi(V) = M$ this implies that there are no singularities of Φ , that is Φ is an Anosov flow. Suppose there are $A, B \in \tilde{\mathcal{F}}^u$ which are not separated from each other. Because $\pi(V) = M$, there is $h \in \pi_1(M)$ so that $h(B) \cap L \neq \emptyset$ and $h(A) \subset V$. But then $h(A) \cap L = \emptyset$ contradicting the above property of L . Hence the leaf space of $\tilde{\mathcal{F}}^s$ is Hausdorff and Φ is an \mathbf{R} -covered Anosov flow. This finishes the proof in this case.

The remaining case is that there is Z leaf of $\tilde{\mathcal{F}}^u$ with $Z \subset V$ and $Z \cap L = \emptyset$. We may assume that Z is non singular. Let $h_1 \in \pi_1(M)$ with $h_1(\gamma)$ in the component of $\tilde{M} - Z$ not intersecting L . Then $h_1(\tilde{W}^u(\gamma)) \subset V$ and $h_1(\tilde{W}^u(\gamma))$ will not intersect any leaf $F' \in \tilde{\mathcal{F}}$ with $F' \geq_s F$. So $\tilde{W}^u(\gamma)$ will not intersect any leaf $\geq_s h_1^{-1}(F)$. This settles claim 1.1.

Now we construct the chains in $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}^s$. As done before let F_1 be the smallest leaf of $\tilde{\mathcal{F}}$ in the front of F and not intersecting $\tilde{W}^u(\gamma)$. Since $g(\tilde{W}^u(\gamma)) = \tilde{W}^u(\gamma)$, then as above $g(F_1) = F_1$. Also γ is in the back of F_1 . Now there is l_1 a stable line leaf in $\partial\Theta(F_1)$ with either γ in $l_1 \times \mathbf{R}$ or γ separated from F_1 by $l_1 \times \mathbf{R}$. If $\gamma \subset \partial\Theta(F_1)$ then since l_1 is a stable line leaf it follows that $\tilde{W}^u(\gamma_1) \cap F_1 \neq \emptyset$. Then $\tilde{W}^u(\gamma) \cap F_1 \neq \emptyset$, contradiction. So the second option occurs. Then as seen before l_1 is a stable line leaf. Let $L_1 = l_1 \times \mathbf{R}$. Then $g(L_1) = L_1$, so there is an orbit γ_1 in L_1 with $g(\gamma_1) = \gamma_1$.




Figure 6: *The sequence of leaves invariant under g .*

We can iterate this process: As before produce F_2 smallest in $\tilde{\mathcal{F}}$ with γ_1 in the back of F_2 and $\tilde{W}^u(\gamma_1) \cap F_2 = \emptyset$. Then $g(F_2) = F_2$ and produce l_2 stable line leaf $l_2 \subset \partial\Theta(F_2)$ and $L_2 = l_2 \times \mathbf{R}$ with $g(L_2) = L_2$. In this way we produce γ_i orbits of $\tilde{\Phi}$, F_i leaves of $\tilde{\mathcal{F}}$, L_i leaves of $\tilde{\mathcal{F}}^s$ with

$$g(\gamma_i) = \gamma_i, \quad g(F_i) = F_i, \quad g(L_i) = L_i, \quad F_i <_s F_{i+1} \quad \text{and} \quad g(\tilde{W}^u(\gamma_i)) \cap F_{i+1} = \emptyset,$$

for all $i \in \mathbf{N}$, see fig. 6. In addition for all $i \in \mathbf{N}$, L_{i+1} separates L_i from L_{i+2} , that is, the L_i are nested in \tilde{M} .

Then the F_i escapes in \tilde{M} when $i \rightarrow \infty$. Otherwise $F_i \rightarrow F^*$ unique leaf of $\tilde{\mathcal{F}}$ since \mathcal{F} is \mathbf{R} -covered. As the family $l_i = \Theta(L_i) \subset \mathcal{O}$, $i \in \mathbf{N}$ is nested and trapped by $\Theta(F)$, then the l_i also converge. There may be more than one limit as the leaf space of $\tilde{\mathcal{F}}_{\mathcal{O}}^s$ may fail to be Hausdorff, but nevertheless there is a unique limit l^* line leaf of $\tilde{\mathcal{F}}_{\mathcal{O}}^s$ which separates $\Theta(F)$ from all $\Theta(L_i)$. Let $L^* = l^* \times \mathbf{R}$. Then $g(L_i) = L_i$ and $g(F) = F$ implies that $g(L^*) = L^*$ so there is a periodic orbit $\gamma^* \subset L^*$ with $g(\gamma^*) = \gamma^*$. Then $g(\tilde{W}^u(\gamma^*)) \cap L_i \neq \emptyset$, for big enough i , producing two orbits invariant by g in $\tilde{W}^u(\gamma^*)$, namely $\tilde{W}^u(\gamma^*) \cap L_i$ and γ^* . This is a contradiction. Hence the F_i escapes in \tilde{M} .

We stress that the construction above works for any periodic orbit γ of $\tilde{\Phi}$. The point here is the following: The general construction suggests there is a standard “direction” in the universal cover “transverse” to $\tilde{\mathcal{F}}^s$ which is created by the increasing F_i . This is the direction of the nested L_i . The following property shows that this “direction” given by the L_i is essentially independent of the periodic orbit γ . This will eventually force the leaf space of $\tilde{\mathcal{F}}^s$ (or $\tilde{\mathcal{F}}^u$) to be non singular and Hausdorff proving the theorem.

2. Nesting property -

If α is another periodic orbit of $\tilde{\Phi}$ associated to the covering translation f of \tilde{M} , the general construction above produces sequences α_i of periodic orbits of $\tilde{\Phi}$, G_i leaves of $\tilde{\mathcal{F}}$ and E_i line leaves of $\tilde{\mathcal{F}}^s$ which are all invariant under f for any $i \in \mathbf{N}$.

Claim 2.1 - The nested sequence $\{E_i\}$, $i \in \mathbf{N}$ is eventually nested with the sequence $\{L_j\}$, $j \in \mathbf{N}$.

This will prove there is a unique “up” direction transverse to $\tilde{\mathcal{F}}^s$.

To prove the claim, first notice that $G_i \in \tilde{\mathcal{F}}$ also has to escape in \tilde{M} when $i \rightarrow \infty$. Hence there is $i_0 \in \mathbf{N}$ so that $G_{i_0} \geq_s F_{j_0}$ for some $j_0 \in \mathbf{N}$. Given $i > i_0$ there is a unique $j(i) \in \mathbf{N}$ ($j(i) \geq j_0$) so that either $G_i = F_{j(i)}$ or $F_{j(i)} \leq_s G_i <_s F_{j(i)+1}$. This shows that the sequences $\{F_k\}$, $k \in \mathbf{N}$ and $\{G_k\}$, $k \in \mathbf{N}$ are eventually nested. We want to show the same thing for the $\{L_k\}$, $\{E_k\}$, $k \in \mathbf{N}$.

Fix i and let $j = j(i)$. Consider L_j, L_{j+1} and the 3 components of $\tilde{M} - (L_j \cup L_{j+1})$:

- W_1 is the component not intersecting F_j or F_{j+1} , essentially “below” F_j ,
- W_2 is the component intersecting F_j but not F_{j+1} . This is the component between L_j and L_{j+1} , equivalently the one which has both L_j and L_{j+1} in its boundary,
- W_3 the component intersecting F_{j+1} , see fig. 7, a.

For simplicity we consider the situation where both L_j and L_{j+1} are non singular. In general $\tilde{M} - (L_j \cup L_{j+1})$ may have more components, but the same arguments apply.

We claim that E_i has to be either L_j or to separate L_j from L_{j+1} , so it must be in the closure of W_2 in \tilde{M} . If for instance $E_i \subset W_1$, then as $\Theta(G_i)$ has a boundary component in $\Theta(E_i)$, then G_i has points in W_1 too. This forces $G_i <_s F_j$, contradiction. If on the other hand $E_i \subset W_3$ then $G_i >_s F_{j+1}$. We conclude that E_i is in the closure of W_2 . Assume that $E_i \neq L_j, L_{j+1}$. If E_i does not separate L_j from L_{j+1} then E_k , $k > i$ will be contained




Figure 7: a. The nesting between E_i and L_j , b. Producing a product region in \widetilde{M} .

in the component of $\widetilde{M} - E_i$ not containing L_{j+1} , see fig. 7, a. This is a contradiction because $G_k > F_{j+1}$. We conclude that E_i separates L_j from L_{j+1} . This proves claim 2.1 This is the nesting we are looking for.

In addition suppose the $\{L_i\}, i \in \mathbf{N}$ are line leaves of $\widetilde{\mathcal{F}}^s$ of any such sequence given by the general construction. Then $\{L_i\}, i \in \mathbf{N}$ escapes in \widetilde{M} as $i \rightarrow \infty$. Otherwise $L_i \rightarrow L^*$ line leaf of $\widetilde{\mathcal{F}}^s$ and maybe other limits as well since the leaf space of $\widetilde{\mathcal{F}}^s$ may fail to be Hausdorff. Let V_2 be the component of $\widetilde{M} - L^*$ not containing the L_i . Then $\pi(V_2) = M$ so let h be a covering translation so that $h(L^*) \subset V_2$. For j big enough $h(L_j) \subset V_2$ and cannot be nested with the sequence $L_i, i \in \mathbf{N}$. This fails the nesting property and is therefore not possible. Therefore we can now conclude that the sequence $\{L_i\}, i \in \mathbf{N}$ escapes in \widetilde{M} .

The general construction and the nesting property will imply that starting in any periodic orbit in \widetilde{M} there is a chain of lozenges going to the “canonical” direction of the $L_i, i \in \mathbf{N}$. By carefully choosing where to put the initial periodic orbits we will show this is incompatible with the existence of singularities of $\widetilde{\Phi}$ or non Hausdorff behavior in the leaf space of $\widetilde{\mathcal{F}}^s$ or $\widetilde{\mathcal{F}}^u$.

CASE I - Φ has singularities.

We will show this does not happen. By way of contradiction, let β be a singular orbit of $\widetilde{\Phi}$. Consider one chain $\gamma_i, L_i, F_i, i \in \mathbf{N}$ given by the general construction, all invariant under $g \in \pi_1(M)$. Since the sequence $\{L_i\}, i \in \mathbf{N}$ escapes in \widetilde{M} and are nested, it follows that the L_i are eventually in a fixed sector V_0 of $\widetilde{W}^s(\beta)$.

Suppose first that all leaves $L_i, i \in \mathbf{N}$ intersect $\widetilde{W}^u(\beta)$. Let $h \in \pi_1(M)$. The sequences of leaves $h(\gamma_i), h(L_i), h(F_i), i \in \mathbf{N}$ are those obtained by the general construction if one starts with the orbit $h(\gamma)$. By the nesting property above, the $\{h(L_j)\}, j \in \mathbf{N}$ are eventually nested with the $\{L_i\}, i \in \mathbf{N}$. Since $h(\widetilde{W}^u(\beta))$ intersects all of $h(L_i)$, it follows that $h(\widetilde{W}^u(\beta))$ will intersect all $L_i, i \geq i_0$ for some i_0 big enough. Choose h so that $h(\widetilde{W}^u(\beta)) \neq \widetilde{W}^u(\beta)$. For any $j > i_0$, the leaves $\widetilde{W}^u(\beta), h(\widetilde{W}^u(\beta)), L_{i_0}, L_j$ form a rectangle in \widetilde{M} , hence there is a product structure in the interior of the rectangle. Let V_1 be the component of $\widetilde{M} - L_{i_0}$ containing L_i for $i > i_0$. Since L_j escapes in \widetilde{M} as $j \rightarrow \infty$, it follows that any leaf unstable leaf of $\widetilde{\mathcal{F}}^u$ between $\widetilde{W}^u(\beta)$ and $h(\widetilde{W}^u(\beta))$ intersects the same set of stable leaves restricted to V_1 . This creates a stable product region contained in V_1 and with base segment in L_{i_0} see fig. 7, b. By theorem 3.8, the foliations $\widetilde{\mathcal{F}}^s, \widetilde{\mathcal{F}}^u$ have leaf space homeomorphic to \mathbf{R} . This contradicts the existence of singularities.

Therefore we may assume that $L_i \cap \widetilde{W}^u(\beta) = \emptyset$ for i big enough and therefore the L_i are eventually in a single component C of $Y = \widetilde{M} - (\widetilde{W}^u(\beta) \cup \widetilde{W}^s(\beta))$. Since there are at least 3 prongs in $\widetilde{W}^s(\beta), \widetilde{W}^u(\beta)$, there are at least 6 components of Y . Let D be a component of Y which is separated from C by at least 3 separatrices on each side. For simplicity we analyse the case that $\widetilde{W}^s(\beta), \widetilde{W}^u(\beta)$ have 3 prongs, see fig. 8, a.

The proof now will go as follows: start with a periodic orbit in D and use the general sequence. This eventually has to go into C , which will be a contradiction.

First consider $\pi(D) \subset M$. If this is not M then it is a region whose boundary may have a corner, coming from




Figure 8: *a. Separating components of Y and first option, b. The union $Q_1 \cup Q_2$ traps lozenges away from C , c. The union $R_1 \cup R_2$ traps lozenges away from C .*

β and is otherwise saturated by leaves of \mathcal{F}^s or \mathcal{F}^u . In a similar way to a previous argument, this is impossible, therefore $\pi(D) = M$. Hence there are periodic orbits of $\tilde{\Phi}$ in D . Let α be one such periodic orbit and f the covering translation of \tilde{M} associated to it. The general construction produces sequences α_i periodic orbits of $\tilde{\Phi}$, E_i line leaves of $\tilde{\mathcal{F}}^s$ and G_i leaves of $\tilde{\mathcal{F}}$, $i \in \mathbf{N}$, all invariant under $f \in \pi_1(M)$ associated to α . By the nesting property the sequence $E_i, i \in \mathbf{N}$ is eventually nested with $L_i, i \in \mathbf{N}$. Therefore both sequences are eventually contained in C . Let i_0 with $L_{i_0} \subset C$. Then $f(\alpha) = \alpha$ and $f(\alpha_{i_0}) = \alpha_{i_0}$. By theorem 3.6 there is a chain of lozenges $\mathcal{B}_k, 1 \leq k \leq n$ from α to α_{i_0} . Let $\delta_k, 0 \leq k \leq n$ be the periodic orbits which are the corners of these lozenges, so $\delta_0 = \alpha, \delta_n = \alpha_{i_0}$. The goal is to show that any such sequence of lozenges has to have a corner in β which will turn out to be impossible.

Let D_1, D_2, D_3, D_4 be the other components of Y as described in fig. 8, a. Let also S_1, S_2, S_3 be the components of $\tilde{W}^s(\beta) - \beta$ as described in fig. 8, a and let U_1, U_2, U_3 be the components of $\tilde{W}^u(\beta) - \beta$. These are the separatrices of Y . The chain of lozenges has to go from D to C so it has to intersect the boundary of D . For simplicity suppose that δ_0 is the last corner in the chain which is contained in D . Therefore \mathcal{B}_1 has to intersect the boundary of D . Suppose first that it crosses the stable boundary of D namely S_1 . The term cross here means that the lozenge \mathcal{B}_1 intersects S_1 in the interior. Therefore an unstable side Q_1 of \mathcal{B}_1 crosses the interior of S_1 into the adjacent component D_1 , see fig. 8, a. In order for the lozenge to close up, the other corner of \mathcal{B}_1 has to be in D_1 , see fig. 8, a. Consider the second lozenge \mathcal{B}_2 . If it is adjacent to \mathcal{B}_1 along an unstable side Q_2 , this forces the corner δ_2 to be contained in D , see fig. 8, a. This contradicts the choice of δ_0 as the only corner in D . Suppose \mathcal{B}_2 is adjacent to \mathcal{B}_1 along a stable side, see fig. 8, b. Then there is an unstable side Q_2 of \mathcal{B}_2 and an unstable side Q_3 of \mathcal{B}_1 , so that Q_2 and Q_3 belong to the same unstable leaf Q , see fig. 8, b. The leaf Q separates the lozenges $\mathcal{B}_1, \mathcal{B}_2$ from C and all further lozenges in the chain are contained in the same component of $M - Q$. Therefore the chain can never reach C , contradiction. The same occurs if \mathcal{B}_2 does not immediately cross to the next component D_2 . The final possibility is that the other corner of \mathcal{B}_2 is in D_2 , that is, \mathcal{B}_2 crosses the unstable standard leaf $U_2 \subset \tilde{W}^u(\beta)$, see fig. 8, c. Then there is a stable side R_1 of \mathcal{B}_1 and a stable side R_2 of \mathcal{B}_2 which are in the same stable leaf R of $\tilde{\mathcal{F}}^s$, see fig. 8, c. The leaf R separates \mathcal{B}_2 from C and will also separate all further lozenges in the chain from the set C . Again this is a contradiction.

We conclude that the chain cannot cross S_1 . In the same way the chain cannot cross U_1 .

The point here is that the chain of lozenges can only cross at most two separatrices of Y but not 3. Since D is separated from C by at least 3 separatrices in each side, there is no way to go from D to C crossing the interior of the separatrices.

The remaining option is that one of the corners in the chain is actually β and \mathcal{B}_1 has sides in S_1 and U_1 rather than crossing these standard leaves. Then \mathcal{B}_2 could jump to the other side and be contained in C , without the chain having to cross the interior of the separatrices. A priori there is no contradiction. But since this has to happen for any periodic orbit in D then any periodic orbit in D has to be connected to β by a chain of lozenges. If the set of periodic orbits in D projects to an infinite collection of closed orbits in M , then they have to limit somewhere in M and so in \tilde{M} also. Therefore there will be two periodic orbits ζ_1, ζ_2 in D so that $\tilde{W}^u(\zeta_1) \cap \tilde{W}^s(\zeta_2) \neq \emptyset$. Let g_0 be the generator of the isotropy group of β . As both ζ_1 and ζ_2 are connected to β by a chain of lozenges, then $g_0(\zeta_1) = \zeta_1, g_0(\zeta_2) = \zeta_2$. This would produce two periodic orbits in $\tilde{W}^u(\zeta_1)$, contradiction. Consequently the collection of periodic orbits in D projects to a finite collection of orbits in M . Since $\pi(D) = M$ this would imply there are only finitely many closed orbits of Φ in M . This is a contradiction [Mo1, Man].

We conclude that case I cannot happen. Therefore Φ does not have singularities.

CASE II - Suppose $\tilde{\mathcal{F}}^s$ does not have Hausdorff leaf space.

Now we know there are no singular orbits in Φ . The general thrust of the proof will be similar to Case I. By hypothesis there are $A_1, A_2 \in \tilde{\mathcal{F}}^s$ which are not separated in their leaf space. By theorem 3.7 A_1, A_2 are connected by a chain of adjacent lozenges all intersecting a common stable leaf. There are an even number of lozenges. For simplicity we may assume there are two lozenges $\mathcal{C}_1, \mathcal{C}_2$ in the chain. A stable leaf intersects \mathcal{C}_1 if and only if it intersects \mathcal{C}_2 . Let ξ_1, ξ_2 be the periodic orbits in A_1, A_2 respectively. There are some special leaves for us: A_1, A_2 are the line leaves of $\tilde{\mathcal{F}}^s$ not separated from each other. A_3 is the line leaf in the boundary of the union of the two lozenges $\mathcal{C}_1 \cup \mathcal{C}_2$ - that is, contains the union of the stable sides of $\mathcal{C}_1, \mathcal{C}_2$ which are not in A_1, A_2 . Finally B_1 (B_2) is the component of $\tilde{W}^u(\xi_1) - \xi_1$ ($\tilde{W}^u(\xi_2) - \xi_2$) which is a side of one of the lozenges \mathcal{C}_1 or \mathcal{C}_2 , see fig. 9, a. This defines 5 regions of \tilde{M} which will be important for us:

- Region (1) - the component of $\tilde{M} - A_1$ not containing A_2 ,
- Region (2) - the component of $\tilde{M} - A_2$ not containing A_1 ,
- Region (3) - the component of $\tilde{M} - A_3$ not containing A_1 ,
- Region (4) - the component of $\tilde{M} - (A_1 \cup B_1)$ not containing A_2 ,




Figure 9: a. Non separated leaves in $\tilde{\mathcal{F}}^s$ and regions of \tilde{M} associated to them. Trapping back and forth of corners of lozenges producing contradiction, b. The union $R_1 \cup R_2$ separates B_2 from A_2 .

– Region (5) - the component of $\tilde{M} - (A_2 \cup B_2)$ not containing A_1 .

Notice that \tilde{M} is the union of these 5 regions, plus the two lozenges and their sides, plus A_1, A_2 , see fig. 9, a.

Let γ be a periodic orbit of $\tilde{\Phi}$ and the sequences γ_i orbits of $\tilde{\Phi}$, L_i line leaves of $\tilde{\mathcal{F}}^s$ and $F_i \in \tilde{\mathcal{F}}$ as given by the general construction. As seen in the beginning of the proof of Part I, the L_i cannot keep intersecting any of the leaves B_1 or B_2 , or also the leaves A_1, A_2, A_3 since these last ones are stable leaves. Therefore the $L_i, i \in \mathbf{N}$ are eventually contained in one of the 5 regions defined above.

We analyse each possibility in turn.

Case II.1 - Suppose the L_i are eventually contained in region (2).

Start with a periodic orbit α in region (1). The general construction produces orbits α_i of $\tilde{\Phi}$, line leaves E_i of $\tilde{\mathcal{F}}^s$ and leaves $G_i \in \tilde{\mathcal{F}}, \forall i \in \mathbf{N}$. These are all invariant under $f \in \pi_1(M)$ associated to α . By the nesting property the E_i are eventually nested with the L_i and so eventually in region (2). As before this produces a chain of lozenges $B_k, 1 \leq k \leq n$, from $\delta_0 = \alpha$ to $\delta_n = \alpha_{i_0}$, with $\alpha_{i_0} \subset$ region (2). Let $\delta_j, 0 \leq j \leq n$ be the corner orbits in this chain. Assume δ_0 is the last corner in region (1). Then B_1 has to intersect A_1 . Assume first the interior of B_1 intersects A_1 . If the second lozenge is adjacent to B_1 along the common unstable side Q_1 , then Q_1 intersects A_1 and this forces δ_2 to be contained in region (1), see fig. 9, a. This contradicts the choice of δ_0 . The other option is that there are stable sides R_1 of B_1 and R_2 of B_2 contained in the stable leaf $\tilde{W}^s(\delta_1)$ and so that $R_1 \cup R_2$ separates B_2 from A_1 and hence from A_2 , see fig. 9, b. Then $R_1 \cup R_2$ separates all further lozenges from A_2 and A_2 cannot be reached by the chain, contradiction. The last option is that the chain has a corner in the periodic orbit ξ_1 in A_1 . But then all periodic orbits in region (1) would have to be connected to this orbit by a chain of lozenges. As seen in the proof of case I, this is not possible either. We conclude that this case cannot happen.

If the sequence $L_i, i \in \mathbf{N}$ is eventually in region (1), then we switch the roles of regions (1) and (2) and apply the same proof.

Case II.2 - The $L_i, i \in \mathbf{N}$ are eventually in region (3).

In this case we let α be a periodic orbit contained in region (4). There is a chain of lozenges $B_k, 1 \leq k \leq n$, with corners $\delta_k, 0 \leq k \leq n$ from α to $\alpha_{i_0} \subset$ region (3). As before assume δ_0 is the last corner in region (4). Assume first the interior of B_1 intersects the unstable leaf $\tilde{W}^u(\xi_1)$ - notice $B_1 \subset \tilde{W}^u(\xi_1)$. Then there is a stable side R_0 of B_1 which intersects $B_1 \subset \tilde{W}^u(\xi_1)$ because $\delta_0 \subset$ region (4). Since the lozenges C_1, C_2 are adjacent, this forces R_0 to intersect C_2 also and enter region (5). Hence δ_1 is contained in region (5). If B_2 is adjacent to B_1 along a stable side R_1 , then this forces B_2 to cross back to region (4) and $\delta_2 \subset$ region (4), see fig. 10, a. This contradicts the choice of δ_0 . Otherwise there is an unstable side Q_1 of B_1 and an unstable side Q_2 of B_2 so that $Q_1 \cup Q_2 \subset \tilde{W}^u(\delta_1)$ and $Q_1 \cup Q_2$ separates B_2 from region (3), see fig. 10, b. In fact $Q_1 \cup Q_2$ will separate all further lozenges from region (3), so α_{i_0} can never be achieved, contradiction. The remaining option is that B_1 has a side in $\tilde{W}^u(\xi_1)$. But then $\delta_1 = \xi_1$. But this is also impossible, as seen before. This proves that case II.2 cannot happen either.

Conversely if the $L_i, i \in \mathbf{N}$ are eventually in region (4), then we would start with $\alpha \subset$ region (3). The above proof shows there is no chain of lozenges from region (4) to region (3), consequently no chain of lozenges from




Figure 10: *a. Trapping back and forth of lozenges, b. Separating leaf which traps lozenges.*

region (3) to region (4). Therefore this cannot happen either. Finally the symmetry between regions (4) and (5) shows that the sequence $L_i, i \in \mathbf{N}$ cannot be eventually in region (5) either.

Since none of the possibilities can occur we conclude that $\tilde{\mathcal{F}}^s$ has Hausdorff leaf space. It now follows that Φ is a (topological) **R**-covered Anosov flow. This finishes the proof of theorem 4.2. \square

5 **R**-covered Anosov flows

The main theorem leaves open the question as to whether **R**-covered pseudo-Anosov flows transverse to **R**-covered foliations can be non regulating. At first it may seem this is not possible, but in fact there are many examples.

We first describe the topological types of **R**-covered pseudo-Anosov flows. If Φ is an **R**-covered pseudo-Anosov flow, then up to isotopy there are only two topological models for the joint structure of the stable and unstable foliations in the universal cover [Ba2, Fe3], which we now describe.

The set \mathcal{O} is homeomorphic to $H = (0, 1) \times \mathbf{R}$, which is more convenient for our description of **R**-covered pseudo-Anosov flows. In the *product* type the foliation $\tilde{\mathcal{F}}^s$ is a foliation by horizontal segments in H and $\tilde{\mathcal{F}}^u$ is a foliation by vertical lines. Notice that any leaf F of $\tilde{\mathcal{F}}^s$ intersects every leaf of $\tilde{\mathcal{F}}^u$ and vice versa.

In the *skewed* model $\tilde{\mathcal{F}}^s$ is again a foliation by horizontal segments in H and now $\tilde{\mathcal{F}}^u$ is a foliation by parallel segments in H making an angle $\theta \neq \pi/2$ with the horizontal. Hence any leaf of $\tilde{\mathcal{F}}^s$ does not intersect all leaves of $\tilde{\mathcal{F}}^u$ and vice versa.

For the remainder of this section we consider smooth Anosov flows. In this case there are 1-dimensional strong stable and unstable foliations $\mathcal{F}^{ss}, \mathcal{F}^{uu}$ which are transverse to each other and generated by subbundles $E^s, \overline{E^u}$ of the tangent bundle. For details see [An, An-Si].

We first study suspension Anosov flows. Let \mathcal{F} be the unstable foliation of a suspension Anosov flow. Then \mathcal{F} is **R**-covered. Notice any stable leaf of $\tilde{\mathcal{F}}^s$ intersects every unstable leaf of $\tilde{\mathcal{F}}^u$ in a single orbit of $\tilde{\Phi}$ and vice versa. We will produce a new Anosov flow, transverse to \mathcal{F} and which is not regulating for \mathcal{F} .

Assume that \mathcal{F} is transversely oriented or that \mathcal{F}^{ss} is oriented. For simplicity assume that M has the solv metric so that $T\Phi$ is perpendicular to both the stable E^s and unstable E^u bundles of Φ . Let $-\pi/2 < \theta < \pi/2$. Define a new flow Φ_θ as follows. In each point of M consider the stable leaf R of \mathcal{F}^s through it. Staying in R , tilt the vector $T\Phi$ by the angle θ , see fig. 12, a. This can be done coherently because \mathcal{F}^{ss} is oriented. This produces a new vector field Φ_θ . For small $|\theta|$, the vector fields Φ_θ and Φ are C^1 -close so by structural stability of Φ [An, An-Si] it follows that Φ_θ is Anosov and topologically conjugate to Φ .

But in fact this is true for any $-\pi/2 < \theta < \pi/2$: Since orbits of Φ_θ are tangent to \mathcal{F}^s it follows that the new orbits are contained in the old stable leaves. These flow lines make constant angle θ with the old flow lines. The stable leaves of $\tilde{\mathcal{F}}^s$ are isometric to the hyperbolic plane \mathbf{H}^2 . Fix a stable leaf $L \in \tilde{\mathcal{F}}^s$. It is easier to see in the upper half plane model for \mathbf{H}^2 where flow lines of $\tilde{\Phi}$ are vertical rays and strong stable leaves of $\tilde{\Phi}$ are horizontal lines (or horocycles), see fig. 12, a. The vertical rays are the intersection of unstable leaves of $\tilde{\mathcal{F}}^u$ with L or the intersections of leaves of $\tilde{\mathcal{F}}$ with L . A curve making a constant angle with the verticals is a ray (Euclidean ray) making angle θ with vertical direction. This curve is a bounded distance from a vertical line. Still flow lines of $\tilde{\Phi}_\theta$ converge exponentially in forward time. The reader may check that the horizontal vectors are uniformly exponentially contracted in the forward direction. The old strong unstable vectors are still uniformly exponentially




Figure 11: *a. Product type, b. Skewed type. The diagrams are bounded in the horizontal direction and unbounded in the vertical direction.*




Figure 12: *a. The tilt of an Anosov flow, b. The picture in a stable leaf.*

expanded in negative time. We conclude that the flow Φ_θ is Anosov with same strong stable foliations as those of Φ , same stable foliation, but different unstable foliation.

Notice that for $\Theta = -\pi/2, \pi/2$ the flow lines will be tangent to the fibers of a fibration of M over the circle and the flow Φ_θ obviously will not be Anosov. For $-\pi < \theta < -\pi/2$ or $\pi/2 < \theta < \pi$, the flow is conjugate to the inverse of the old one.

Now given an orbit γ of $\tilde{\Phi}_\theta$ contained in L , γ is a ray making angle θ with the verticals and γ will only intersect a sub collection of the vertical lines in L . Therefore γ will not intersect all leaves of $\tilde{\mathcal{F}}$. We conclude that $\tilde{\Phi}_\theta$ is transverse to \mathcal{F} but is not regulating for \mathcal{F} and still \mathcal{F} is **R**-covered. This provides counterexamples to the main theorem if one allows Φ to be an **R**-covered pseudo-Anosov flow.

Once this example is well understood, it is easy to see that any perturbation of Φ keeping it tangent to leaves of \mathcal{F}^s , transverse to $\mathcal{F}^u = \mathcal{F}$ and perturbed with angle $< \pi/2$, will produce an Anosov flow transverse to \mathcal{F} and which is not regulating for \mathcal{F} . We can also do the same for small transverse perturbations of Φ outside of being tangent to \mathcal{F}^s .

As for the skewed case: Assume again that \mathcal{F}^u is transversely orientable. A small perturbation of Φ tangent to \mathcal{F}^s and transverse to $\mathcal{F}^u = \mathcal{F}$, will produce a new flow Φ' topologically conjugate to Φ , hence Anosov. Let L be a leaf of $\tilde{\mathcal{F}}^s$ (same for $\tilde{\Phi}$ and $\tilde{\Phi}'$). Orbits of Φ' in L can only intersect the leaves of $\tilde{\mathcal{F}} = \tilde{\mathcal{F}}^u$ (unstable from the original flow $\tilde{\Phi}$) intersected by L . But these are not all of the leaves of $\tilde{\mathcal{F}}$ because of the skewed picture. This shows that Φ' is not regulating for \mathcal{F} . The easier picture to see in this case, is that of a geodesic flow in the unit tangent bundle T_1S , S a closed surface of constant negative curvature.

6 Non regulating pseudo-Anosov flows and non \mathbf{R} -covered foliations

In this section we use the main theorem to analyse a large class of known examples of foliations transverse to pseudo-Anosov flows and so that the foliations are not \mathbf{R} -covered. There are two types of constructions: branched covers and cut and paste operations.

I. Branched cover construction -

As in the last section consider \mathcal{F} to be the unstable foliation of a suspension Anosov flow Φ . Assume that \mathcal{F}^{ss} is orientable. Fix $\theta \in (-\pi/2, \pi/2)$. As in the previous section construct the Anosov flow Φ_θ which is transverse to \mathcal{F} but not regulating for \mathcal{F} .

Fix a periodic leaf R of \mathcal{F}^s with periodic orbit γ_0 . Then Φ_θ also has a periodic orbit $\gamma \subset R$ which is transverse to the flow lines of Φ when restricted to R . The simple closed curve γ is transverse to \mathcal{F} . Also γ is transverse to the fibration of M over the circle associated to the suspension flow Φ .

The complement $M - \gamma$ is irreducible, atoroidal and has an incompressible surface, hence it has a complete hyperbolic metric of finite volume [Th2, Th4]. By Thurston's Dehn surgery theorem, for n big enough, the $(0, n)$ Dehn filling along γ has a hyperbolic structure [Th2]. Here $(0, 1)$ is the meridian, that is, the curve bounding a fiber in $M - \gamma$. Dehn filling produces a hyperbolic orbifold M_n with singular locus γ and index n along γ . By Selberg's theorem about finite index subgroups without torsion [Se], there is a finite index orbifold cover of M_n which is a manifold N . In other words N is a branched cover of M with branch locus γ and index n along γ . For background on branched covers consult [Rol].

Since the branch set is a closed orbit γ of Φ_θ , the branched cover construction produces a flow Φ_1 in N which is a branched cover of the flow Φ_θ . Locally near γ the construction unwraps the manifold M_n n times around γ , therefore the new flow Φ_1 is a pseudo-Anosov flow in N . The only singular orbits are those coming from γ , where there is a n to 1 branched cover of a neighborhood of γ .

The curve γ is transverse to \mathcal{F} in M , hence leaves of \mathcal{F} meet a neighborhood of γ in meridian disks. Under the branched cover construction $N \rightarrow M_n$, \mathcal{F} lifts to a foliation \mathcal{F}_1 in N . Locally near lifts of γ , leaves of \mathcal{F}_1 are n to 1 branched covers of disks in leaves of \mathcal{F} . The pseudo-Anosov flow Φ_1 in N is transverse to the foliation \mathcal{F}_1 . The foliation \mathcal{F}_1 is the one we are interested in analysing.

We claim that Φ_1 is not regulating for \mathcal{F}_1 . To see this consider first the universal cover \widetilde{M} of M , with lifts $\widetilde{\mathcal{F}}, \widetilde{\Phi}$. Let Z be the leaf of \mathcal{F} containing γ_0 , that is the unstable leaf (of Φ !) containing γ_0 . Let $\tilde{\gamma}$ be a lift of γ to \widetilde{M} and $F \in \widetilde{\mathcal{F}}$ the corresponding lift of Z . By construction of γ and Z , it follows that $\tilde{\gamma} \cap F = \emptyset$. The universal cover \widetilde{N} is a branched cover of \widetilde{M} , branched along all lifts of γ to \widetilde{M} . It follows that any lift of $\tilde{\gamma}$ to \widetilde{N} will not intersect any lift of F to \widetilde{N} . Hence some orbits of Φ_1 do not intersect every leaf of $\widetilde{\mathcal{F}_1}$. It follows that Φ_1 is not regulating for \mathcal{F}_1 .

Since Φ_1 is obviously a singular pseudo-Anosov flow it cannot be an \mathbf{R} -covered pseudo-Anosov flow. If \mathcal{F}_1 were \mathbf{R} -covered, the flow Φ_1 would have to be regulating, by the main theorem. We conclude that \mathcal{F}_1 is not \mathbf{R} -covered.

The orbit γ of Φ_θ is transverse to the original fibration of M over the circle. Under the branched cover construction, the fibration in M lifts to a fibration in N and hence Φ_1 is a suspension pseudo-Anosov flow.

In summary the main ingredient for the construction to work is a branch cover along a closed orbit of the given flow which is not regulating for the original foliation. It follows that this applies to a large class of examples.

II. Cut and paste construction -

The second class of examples is obtained by cut and paste operations on a fibration. Let M be a manifold fibering over the circle with pseudo-Anosov monodromy η , that is $M = S \times [0, 1]/(x, 1) \cong (\eta(x), 0)$, where S is a hyperbolic surface and η is a pseudo-Anosov homeomorphism of S [Th3, FLP]. Suppose that all singularities of η have an even number of prongs. This can be achieved for instance by taking branched covers of Anosov diffeomorphisms of the torus T^2 , see expose 13 in [FLP]. Then choose η so that there is a simple closed curve α in S so that $\alpha \cap \eta(\alpha) = \emptyset$. This can also be achieved by taking finite covers and the fact that $\pi_1(S)$ is LERF [Sc] (see explicit construction in section 2 of [CLR1]). Let λ be the expansion/contraction constant associated to the homeomorphism η , that is, under η unstable lengths get multiplied by λ and stable lengths get divided by λ .

Now consider the suspension flow Φ of η which is a suspension pseudo-Anosov flow. This flow will be fixed throughout the discussion. Let \mathcal{F}_0 be the original fibration of M .

In [CLR1] Cooper et al did cut and paste to the fiber S along an annulus transverse to the fibration and with base in $\alpha \subset S$ to obtain closed immersed surfaces transverse to Φ . Many of these surfaces were not virtual fibers producing fascinating examples. Here we do cut and paste to produce a non \mathbf{R} -covered foliation.

Let A be the annulus $\alpha \times [0, 1]$ in $S \times [0, 1]$. Since $\alpha \cap \eta(\alpha) = \emptyset$, then A embeds in M . Extend A a little bit beyond both boundaries to get an embedded annulus in M transverse to \mathcal{F}_0 with boundaries in leaves of \mathcal{F}_0 covering a little more than one period of the fibration. Let this be the annulus A thought to be in M . Then \mathcal{F}_0




Figure 13: *a. Cutting and pasting in fibrations, b. Shearing along an annulus.*

induces a foliation by circles on A - parametrize this $\alpha \times [-a, 1 + a]$ (where $a > 0$). Cut open M along A and glue back by $\alpha \times \{t\} \rightarrow \alpha \times \{h(t)\}$, where h is a homomorphism of $[-a, 1 + a]$ so that $h(x) > x$ for any $x \in (-a, 1 + a)$. There is $b > 0$ so that for any $x \in [0, 1]$, then $h(x) > x + b$. The cut and paste of \mathcal{F}_0 produces a new foliation \mathcal{F} still transverse to the suspension flow Φ .

This construction is well known to Thurston, Gabai and others. Here we analyse the leaf space of the ensuing foliations.

We think of Φ as a true suspension so that it takes time 1 to go between any two lifts of S to \tilde{M} . We will measure height in \tilde{M} using the time parameter. Let \tilde{S}_0 be the lift of S at height 0.

Think of S as $S \times \{0\}$ embedded in M . Let \mathcal{F}_0^{uu} be the singular strong unstable foliation of η in S . Notice \mathcal{F}_0^{uu} is orientable because of the even parity of prongs. Therefore following leaves of \mathcal{F}_0^{uu} in the positive direction, they will always intersect α in the same direction, which we may assume is the direction which increases the height under the glueing by h . In addition since leaves of \mathcal{F}_0^{uu} are dense in S any segment in a leaf of \mathcal{F}_0^{uu} intersects α after at most length $c_0 > 0$. It easily follows that there is $c > 0$ so that any segment of a leaf of \mathcal{F}^{uu} (of Φ !) of length $> c$, has to intersect a lift of $\alpha \times [0, 1]$.

Let F be a leaf of $\tilde{\mathcal{F}}$. Since Φ is transverse to F the unstable foliation $\tilde{\mathcal{F}}^u$ induces a singular one dimensional foliation $\tilde{\mathcal{F}}_F^u$ in F . Let l be a leaf of $\tilde{\mathcal{F}}_F^u$, contained in $L \in \tilde{\mathcal{F}}^u$. We will keep track of the progress of this leaf in \tilde{M} . For simplicity suppose we start at a point in l of height 0 and follow l in the positive direction. Assume L is non singular. Each time l crosses a lift of $\alpha \times [0, 1]$, then l goes up by at least height b and this happens at least every length c in l . Projecting back to \tilde{S}_0 along flow lines of $\tilde{\Phi}$ the distances along l are contracted by at least λ^b as compared to what happened before the crossing [Ca-Th]. Therefore as seen from the projection in \tilde{S} , then after length at most $\lambda^{-b}c$, the leaf l will hit a lift of $\alpha \times [0, 1]$ again and l goes up again by at least height b . When projecting to \tilde{S} distance are now contracted by at least λ^{2b} . By induction after length at most kc along l , which corresponds to the length

$$c + \lambda^{-b}c + \lambda^{-2b}c + \dots + \lambda^{-b(k-1)}c = c \frac{1 - \lambda^{-bk}}{1 - \lambda^{-b}}$$

when projected to \tilde{S} , then l is at a height at least kb . Let l' be $L \cap \tilde{S}_0$. By letting $k \rightarrow \infty$ in the argument above we see that the height of l blows up to ∞ before the projection of l to l' achieves a length of

$$c \frac{1}{1 - \lambda^{-b}}$$

from the initial point in l' . It follows that F cannot intersect the orbit of $\tilde{\Phi}$ corresponding to the intersection of \tilde{S} with the limit of the projection of l to l' . Otherwise by uniqueness of intersections of F with orbits of $\tilde{\Phi}$, the height of l would be bounded near this limit orbit, contradiction to it increasing without bound.

We conclude that F does not intersect this orbit of $\tilde{\Phi}$ and hence Φ is not regulating for \mathcal{F} . Since Φ is a singular pseudo-Anosov flow the main theorem implies that \mathcal{F} is not \mathbf{R} -covered. This finishes the analysis of the second class of examples.




Figure 14: *The skewed picture produces a slithering over the circle.*

7 Foliations transverse to \mathbf{R} -covered Anosov flows

If Φ is a pseudo-Anosov flow transverse to an \mathbf{R} -covered foliation \mathcal{F} then as seen in section 5, Φ may fail to be regulating if Φ is an \mathbf{R} -covered pseudo-Anosov flow. We will show in this section that this failure is very rare. The goal is to prove a topological rigidity result: that \mathcal{F} is essentially topologically conjugate to one of the invariant foliations of Φ .

First we need some facts about the structure of \mathbf{R} -covered pseudo-Anosov flows. Consider Φ an \mathbf{R} -covered pseudo-Anosov flow of product type. If g is a covering translation of \widetilde{M} , then g leaves invariant at most one leaf of $\widetilde{\mathcal{F}}^s$ (or $\widetilde{\mathcal{F}}^u$) [Fe2]. Equivalently there are no non trivial free homotopies between closed orbits of Φ .

In this case $\pi_1(M)$ is solvable and there has been important work of Plante [Pl2] completely classifying foliations in such manifolds. However Plante assumes that \mathcal{F} is at least C^2 and he makes use of Kopell’s lemma [Ko]. We do not assume that \mathcal{F} is C^2 , in fact in this particular case it is very easy to produce \mathcal{F} transverse to Φ which cannot be C^2 .

Recall that \mathcal{H} is the leaf space of $\widetilde{\mathcal{F}}$ and similarly $\mathcal{H}^s, \mathcal{H}^u$.

The structure of a skewed \mathbf{R} -covered pseudo-Anosov flow is much more complex and rich. Let Φ have skewed type. For simplicity suppose that $\mathcal{F}^s, \mathcal{F}^u$ are transversely orientable. Fix a transverse orientation to \mathcal{F}^s and put an order $<_s$ in \mathcal{H}^s coherent with this transverse orientation. Given $\widetilde{F} \in \widetilde{\mathcal{F}}^s$ there is a unique unstable leaf U which makes a perfect fit with F and is contained in the component of $\widetilde{M} - F$ defining the positive orientation from F [Fe3]. Intuitively the leaf U is “asymptotic” to F , see fig. 14. Let this U be denoted by $\tau_+(F)$. Then τ_+ defines a homeomorphism from \mathcal{H}^s to \mathcal{H}^u . Similarly define $\tau_-(F)$ which makes a perfect fit with F and is on the negative side of F , see fig. 14. Then τ_- is also a homeomorphism from \mathcal{H}^s to \mathcal{H}^u . We choose a transversal orientation $<_u$ to \mathcal{F}^u so that $\tau_-(F) <_u \tau_+(F)$.

Let $\zeta : \mathcal{H}^s \rightarrow \mathcal{H}^s$ be the homeomorphism $(\tau_-)^{-1} \circ \tau_+$. Then ζ is fixed point free [Fe3, Fe6]. Intuitively it goes one step up in the infinite ladder of asymptotic leaves. Then \mathcal{H}^s / ζ is homeomorphic to a circle and there is a map $\delta : \widetilde{M} \rightarrow \mathbf{S}^1$ induced by the map $\mathcal{H}^s \rightarrow \mathcal{H}^s / \zeta$. The map δ is group equivariant: there is a group homomorphism $\rho : \pi_1(M) \rightarrow \text{Homeo}(\mathbf{S}^1)$, so that

$$\forall g \in \pi_1(M), \forall x \in \widetilde{M}, \quad \delta(g(x)) = \rho(g)(\delta(x)).$$

or $\delta \circ g = \rho(g) \circ \delta$ [Fe3, Fe6]. Therefore the foliation $\widetilde{\mathcal{F}}^s$ (and consequently \mathcal{F}^s) is induced by δ . The map δ is an example of what Thurston calls a *slithering* over the circle [Th5]. We refer the reader to [Th5] for the beautiful theory of slitherings. They are very common in 3-manifolds. We will use a few simple properties of foliations defined by slitherings. One property which follows easily from the slithering map is that given any $F, L \in \widetilde{\mathcal{F}}^s$ they are a bounded (Hausdorff) distance apart: there is $c_0 > 0$ so that for any $x \in F, y \in L$ then $d(x, L) < c_0$ and $d(y, F) < c_0$. Here the number $d(x, L)$ is the infimum of $d(x, z)$ for all $z \in L$. Clearly c_0 depends on the pair F, L . Let $d_H(F, L)$ be the smallest c_0 which works. This is the *Hausdorff* distance between leaves in \widetilde{M} .

Notice that distance in \mathcal{H}^s is not well defined but slithering units are: we say that $a, b \in \mathcal{H}^s$ are n slithering units apart if $a = \zeta^n(b)$ or $b = \zeta^n(a)$. Then given any $a, b \in \mathcal{H}^s$ with $a <_s b$ let

$$d_s(a, b) = n, \text{ if } \zeta^n(a) \leq_s b <_s \zeta^{n+1}(a).$$

Extend it to $a \geq_s b$ by making d_s symmetric. This is a roughly defined distance in \mathcal{H}^s . One fundamental fact is that d_s is invariant under covering translations: if $g \in \pi_1(M)$ then for any $a, b \in \mathcal{H}^s$, $d_s(a, b) = d_s(g(a), g(b))$. This




Figure 15: *a. The case where boundary of $\partial\Theta(F_0)$ has only one component produces impossible intersection, b. An unstable boundary also produces impossible intersection.*

follows from the invariance of the infinite staircase picture. Another important fact proved in [Th5] is that there are constants c_1, c_2 so that for any $E, L \in \tilde{\mathcal{F}}^s$ then

$$c_1(d_s(E, L) - 1) \leq d_H(E, L) \leq c_2(d_s(E, L) + 1) \quad (1).$$

This says that the slithering distance between leaves roughly measures the Hausdorff distance between leaves. In addition the distance between points in leaves is also bounded below: there is $c_3 > 0$ so that for any $E, L \in \tilde{\mathcal{F}}^s$ then

$$\forall p \in E, q \in L, \quad d(p, q) > c_3(d_s(E, L) - 1). \quad (2)$$

Whenever the distance d_s is used, it is implicit that Φ is a skewed \mathbf{R} -covered pseudo-Anosov flow and d_s refers to the distance defined here.

Many 3-manifolds containing skewed Anosov flows are Seifert fibered. Ghys [Gh] showed the manifold is up to finite covers T_1S where S is a hyperbolic surface. Foliations in Seifert fibered manifolds have also been extensively studied by Thurston [Th1], Levitt [Le] and finally Brittenham [Br]. This could help us a bit in the Seifert fibered case. However skewed \mathbf{R} -covered pseudo-Anosov flows can occur in much more general manifolds, for instance the manifolds may be hyperbolic [Fe3].

Unless otherwise stated, for the remainder of this section Φ will denote an \mathbf{R} -covered pseudo-Anosov flow and \mathcal{F} a transverse foliation. We first show that \mathcal{F} has to be \mathbf{R} -covered.

Proposition 7.1. *If Φ is an \mathbf{R} -covered pseudo-Anosov flow transverse to a foliation \mathcal{F} then \mathcal{F} is an \mathbf{R} -covered foliation.*

Proof. Let the positive transversal direction to \mathcal{F} be defined by positive flow direction of Φ . Up to a double cover if necessary, assume that \mathcal{F}^s is transversely oriented. We assume that the order $<_s$ in \mathcal{H}^s is coherent with the transverse orientation to \mathcal{F}^s . Notice that it is preserved by covering translations.

Suppose the proposition is not true. Clearly Φ is not regulating for \mathcal{F} . Let $F_0, F_1 \in \tilde{\mathcal{F}}$ not separated from each other, say in their negative sides. As they are not separated, they cannot intersect a common orbit of $\tilde{\Phi}$, so $\Theta(F_0) \cap \Theta(F_1) = \emptyset$. Hence there is either a stable or unstable boundary leaf of $\Theta(F_0)$ separating it from $\Theta(F_1)$. If it is unstable, then the corresponding unstable leaf of $\tilde{\mathcal{F}}^u$ is on the positive side of F_0 . Therefore F_1 is also in the positive side of F_0 , contradiction to assumption. Let therefore L_0 be a leaf of $\tilde{\mathcal{F}}^s$ with $\Theta(L_0) \subset \partial\Theta(F_0)$ and L_0 separating F_0 from F_1 . In the same way there is $L_1 \in \tilde{\mathcal{F}}^s$ with $\Theta(L_1) \subset \partial\Theta(F_1)$ and L_1 separating F_1 from F_0 . Notice that L_0 and L_1 may be the same leaf. Let $l_i = \Theta(L_i)$. Assume that F_1 intersects stable leaves which are $<_s L_0$, see fig. 15 a. Given these conditions we make the following claim.

Claim – For any leaf $F \in \tilde{\mathcal{F}}$, then $\partial\Theta(F)$ is a union of two stable leaves of $\tilde{\mathcal{F}}^s$.

We first prove the claim for F_0 defined above. If the boundary of $\Theta(F_0)$ consists only of l_0 , then F_0 intersects all $L \in \tilde{\mathcal{F}}^s$ with $L >_s L_0$. Take a covering translate $g(L_0)$ of L_0 so that $g(L_0) \cap F_1 \neq \emptyset$. Hence $g(L_0) <_s L_1$. We now consider the picture near $g(L_0)$. Let $p \in \Theta(g(L_0))$ and $p_i \in \Theta(g(F_0))$ with $p_i \rightarrow p$. Let $y_i \in \Theta(F_1)$ with $\Theta(y_i) = p_i$. Since $p \in \Theta(F_1)$ then $y_i \rightarrow y$ with $\Theta(y) = p$. Let $x \in g(F_0)$ with $\Theta(x_i) = p_i$. As $p \in \partial\Theta(g(F_0))$, then $x_i = \tilde{\Phi}_{t_i}(y_i)$ with $|t_i| \rightarrow \infty$. Since $\Theta(g(L_0))$ is a stable boundary leaf of $\Theta(g(F_0))$ it follows that $t_i \rightarrow +\infty$. This implies that $g(F_0)$ is in the front of F_1 . Applying the same argument to points in L_1 we obtain that $g(F_0)$ is in the back of F_1 . But that would imply that $g(F_0)$ and F_1 have to intersect, which is a contradiction, see fig. 15, a.




Figure 16: *Producing unstable boundary.*

Hence $\Theta(F_0)$ has more boundary. If it is unstable boundary, then points in F_0 go down when points in $\Theta(F_0)$ approach this boundary. Take $g(L_0) \cap F_1 \neq \emptyset$. As seen in the argument of the previous paragraph this implies the contradiction $g(F_0) \cap F_1 \neq \emptyset$, see fig. 15, b. We conclude that there should be more stable boundary to $\Theta(F_0)$. Finally given any 3 stable leaves of $\tilde{\mathcal{F}}^s$, one of them separates the other two, hence it follows that $\partial\Theta(F_0)$ can have only two components. This proves the claim for F_0 .

Let l'_0 be the other component of $\partial\Theta(F_0)$ and let $L'_0 \in \tilde{\mathcal{F}}^s$ with $\Theta(L'_0) = l'_0$. Now given any $F \in \tilde{\mathcal{F}}$, there is $h(F)$ which is in front of F_0 . But then $h(F)$ cannot intersect L_0 or L'_0 . Therefore $h(F)$ has to escape in the positive direction before it gets to L_0 or L'_0 . It follows that $\partial\Theta(h(F))$ also has two components and the same holds for F . This finishes the proof of the claim.

It is this claim that will eventually lead to a contradiction.

If all leaves of \mathcal{F} are simply connected, then \mathcal{F} is a foliation by planes and this implies that M is the 3-torus T^3 [Ros]. But T^3 does not support an \mathbf{R} -covered pseudo-Anosov flow, because it has abelian fundamental group and 3-manifolds supporting \mathbf{R} -covered pseudo-Anosov flows have exponential fundamental group [Pl-Th].

Let then F_2 with $\pi(F_2)$ not simply connected. Let $g \in \pi_1(M)$, not identity, with $g(F_2) = F_2$. Let $\partial\Theta(F_2) = l_2 \cup l'_2$ and $L_2, L'_2 \in \tilde{\mathcal{F}}^s$ with $\Theta(L_2) = l_2$, $\Theta(L'_2) = l'_2$. Since g preserves orientation of \mathcal{H}^s , the fact that $g(F_2) = F_2$ implies that $g(L_2) = L_2$ and $g(L'_2) = L'_2$.

If Φ has product type this immediately leads to a contradiction, because g would have two invariant leaves in $\tilde{\mathcal{F}}^s$.

In the skewed case this does not lead to a contradiction yet. Suppose that $d_s(L_2, L'_2) = n$. Consider $h \in \pi_1(M)$, with $h(L_2) >_s L'_2$ and $d_s(L_2, h(L_2)) > n + 3$. Fix an orbit γ in L_2 . Consider $E \in \tilde{\mathcal{F}}$ with $E \cap L_2 \neq \emptyset$. Then $E \cap \gamma \neq \emptyset$. Suppose first that no such E intersects $h(L_2)$. Let E_i be a sequence of leaves of $\tilde{\mathcal{F}}^s$ so that $E_i \cap \gamma$ is monotonic in γ and escapes in the negative flow direction of γ . Let S_i be the stable leaf with $S_i >_s L_2$ and $\Theta(S_i) \subset \partial\Theta(E_i)$, see fig. 16. Because $E_i \cap \gamma$ is monotonic in γ then S_i are (weakly) monotonic and increasing in \mathcal{H}^s . But since no E_i intersects $h(L_2)$ then $S_i \leq_s h(L_2)$. Therefore $S_i \rightarrow S$ in \mathcal{H}^s as $i \rightarrow \infty$, see fig. 16.

Choose $R \cap S \neq \emptyset$. Then R is in the back of any E_i so $R \cap L_2 = \emptyset$. Also $R \cap S_i \neq \emptyset$ for i big enough and therefore R cannot escape in the positive direction before getting to L_2 . One concludes that R has to escape down, producing unstable boundary in $\Theta(R)$, see fig. 16. This contradicts the claim above.

We conclude that there is E with $E \cap L_2 \neq \emptyset$ and $E \cap h(L_2) \neq \emptyset$. Let $A_0, A'_0 \in \tilde{\mathcal{F}}^s$ with $\partial\Theta(E) = \Theta(A_0) \cup \Theta(A'_0)$. Then $A_0 <_s L_2$ and $A'_0 >_s h(L_2)$. Therefore $d_s(A_0, A'_0) \geq n + 3$.

Choose $f \in \pi_1(M)$ with $f(E)$ in front of F . This implies that

$$L_2 \leq_s f(A_0) <_s f(A'_0) \leq_s L'_2.$$

Then

$$d_s(f(A_0), f(A'_0)) \leq d_s(L_2, L'_2) = n \leq d_s(A_0, A'_0) - 3.$$

This contradicts the fact that the covering translation f preserves d_s . This contradiction shows there is also a problem in the skewed case. We conclude that the claim is impossible, that is, for any $F \in \tilde{\mathcal{F}}^s$, the set $\Theta(F)$ can have at most one stable boundary component. Finally, the failure of the claim implies that \mathcal{F} is \mathbf{R} -covered as desired. This finishes the proof of proposition 7.1. \square

Given that \mathcal{F} is \mathbf{R} -covered we now start to analyse the relationships between $\mathcal{F}, \mathcal{F}^s, \mathcal{F}^u$. We first lift to a double cover if necessary so that \mathcal{F}^s is transversely orientable. As shown by Barbot [Ba2], this implies in the skewed case that \mathcal{F}^u is also transversely oriented. The standing hypothesis for the remainder of this section (unless otherwise stated) is that there is a fixed leaf $F^* \in \tilde{\mathcal{F}}$ with $\Theta(F^*)$ having a stable boundary component.

Lemma 7.2. *For any $F \in \tilde{\mathcal{F}}$ the set $\Theta(F)$ has a unique stable boundary component.*

Proof. The leaf F^* has a stable boundary component l^* . Since Φ is transitive, it follows from symbolic dynamics [Bow1, Bow2] that there are dense orbits of Φ in M . Let γ be an orbit of Φ which is dense in M and let $\tilde{\gamma}$ a lift of γ which intersects F^* . There is a translate $f_0(F)$ of F so that $\tilde{\gamma}$ intersects $f_0(F)$ after intersecting F^* . Then $\Theta(F^*), \Theta(f_0(F))$ intersect in a point $p = \Theta(\gamma) \in \mathcal{O}$. Let α be a path in $\Theta(F^*) \cup l^*$ from p to $p' \in l^*$, intersecting l^* only in p' . This path cannot be totally lifted to F^* because it escapes in the positive direction as the projection to \mathcal{O} approaches p' . Since $f_0(F)$ is above F^* , the path cannot be totally lifted to $f_0(F)$ either and has to escape in the positive flow direction before the projection in \mathcal{O} gets to p' . This produces stable boundary in $\Theta(f_0(F))$. Translating back by f_0^{-1} implies that $\Theta(F)$ also has stable boundary.

Finally the proof of the previous proposition shows that there is at most one boundary stable leaf. \square

These results show that given $F \in \tilde{\mathcal{F}}$ there is a unique $L \in \tilde{\mathcal{F}}^s$ with $\Theta(L) \subset \partial\Theta(F)$. Let this L be $\eta_s(F)$. This defines a function $\eta_s : \mathcal{H} \rightarrow \mathcal{H}^s$, $\eta_s(F) = L$. The function η_s commutes with covering translations:

$$\forall g \in \pi_1(M), \quad \eta_s \circ g = g \circ \eta_s.$$

We now fix a transversal orientation $<_s$ to \mathcal{F}^s so that under this orientation F^* is contained in the front of $\eta_s(F^*)$. As explained in the beginning of the section this defines a transversal orientation $<_u$ to \mathcal{F}^u so that $\tau_+ \circ (\tau_-)^{-1}$ is increasing in \mathcal{H}^u .

Recall that $\eta_s(F^*)$ is in the back of F^* with respect to positive flow direction of $\tilde{\Phi}$, hence $\eta_s(F^*)$ is in the back of F^* with respect to the transverse orientation of $\tilde{\mathcal{F}}$. Given a translate $g(F^*)$ of F^* then $\eta_s(g(F^*)) = g(L^*)$ is in the back of $g(F^*)$ (as seen from the transverse orientation to $\tilde{\mathcal{F}}$) and $g(F^*)$ is in the front of $g(L^*)$ (as seen from the transverse orientation to $\tilde{\mathcal{F}}^s$). Therefore because the translates of L^* are dense in \tilde{M} , this property follows for all $F \in \tilde{\mathcal{F}}$ and $L = \eta_s(F) \in \tilde{\mathcal{F}}^s$ – this is because \mathcal{F} is \mathbf{R} -covered. Hence if $F_1 < F_2$ in \mathcal{H} , then F_2 is in the front of F_1 and so F_1 separates $\eta_s(F_1)$ from F_2 . It follows that $\eta_s(F_2)$ is also in the front of $\eta_s(F_1)$ (from the point of view of $\tilde{\mathcal{F}}^s$). This implies that $\eta_s : \mathcal{H} \rightarrow \mathcal{H}^s$ is a weakly monotone increasing map.

If the original \mathcal{F}^s is not transversely orientable, there is $g \in \pi_1(M)$ reversing any orientation to \mathcal{H}^s . Then $g(F^*)$ would be in the back of $\eta_s(g(F^*))$, contradicting the discussion above. It follows that the original \mathcal{F}^s is transversely orientable and it is not necessary to lift to a double cover. In addition if Φ is skewed, then \mathcal{F}^u is also transversely oriented.

We stress that the fact that both \mathcal{H} and \mathcal{H}^s are homeomorphic to the real numbers, is essential throughout the proofs in this section. The leaf F^* of $\tilde{\mathcal{F}}$ is fixed throughout the discussion. Let $L^* = \eta_s(F^*)$.

Lemma 7.3. *The map $\eta_s : \mathcal{H} \rightarrow \mathcal{H}^s$ is continuous.*

Proof. Consider $F_i \in \mathcal{H}$ with $F_i \rightarrow F$ in \mathcal{H} . We may assume that F_i is monotonic in \mathcal{H} .

Case 1 – F_i is decreasing.

Then $L_i = \eta_s(F_i)$ is weakly monotone decreasing. Clearly $L = \eta_s(F) \leq_s \eta_s(F_i)$ for all i , hence L_i is bounded below and $L_i \rightarrow L^-$ in \mathcal{H}^s . If $L <_s L^-$, then either F intersects L^- or L^- separates L_i from F for all i , see fig. 17, a. Hence for a fixed j big enough, either F_j intersects L^- or F_j jumps to the other side of L^- . Both options are disallowed by $L_i \geq_s L^-$ for all i . Hence $L^- = L$ and $\eta_s(F_i) \rightarrow \eta_s(F)$.

Case 2 – F_i is increasing in \mathcal{H} .

Suppose that $L_i = \eta_s(F_i) \rightarrow L^-$ with $L^- <_s L = \eta_s(F)$, see fig. 17, b. Then choose a covering translate $g(L_1)$ with

$$L^- <_s g(L_1) <_s L.$$

The map η_s is weakly monotonic, hence $g(F_1) < F$ in \mathcal{H} . But also as $\eta_s(g(F_1)) = g(L_1) >_s L^-$, then $g(F_1) > F_i$ in \mathcal{H} for any i . This contradicts the fact that $F_i \rightarrow F$, see fig. 17, b. This shows that $L^- = L$ and so $\eta_s(F_i) \rightarrow \eta_s(F)$.

It now follows that η_s is continuous as desired. \square

Corollary 7.4. *For any stable leaf $L \in \tilde{\mathcal{F}}^s$ there is $F \in \tilde{\mathcal{F}}$ with $L = \eta_s(F)$. In addition $\eta_s^{-1}(L)$ is a compact interval in \mathcal{H} .*




Figure 17: a. The case F_i is decreasing, b. The case F_i is increasing.

Proof. To prove the first statement choose $g_0, g_1 \in \pi_1(M)$ with $g_0(L^*) <_s L <_s g_1(L^*)$. Then $g_0(L^*) = \eta_s(g_0(F^*))$ and $g_1(L^*) = \eta_s(g_1(F^*))$ are in the image of η_s . As η_s is continuous, its image is connected, which implies that L is also in the image of η_s . The second statement follows from η_s being continuous, surjective, monotonic and \mathcal{H} homeomorphic to \mathbf{R} . \square

We now show that leaves of $\tilde{\mathcal{F}}$ are boundedly near leaves of $\tilde{\mathcal{F}}^s$.

Proposition 7.5. *Let $F \in \tilde{\mathcal{F}}^s$ and $L = \eta_s(F)$. Then the Hausdorff distance between F and L is finite and bounded by a global constant r_0 .*

Proof. Suppose not. Then there are $F_i \in \tilde{\mathcal{F}}$, $L_i = \eta_s(F_i)$ which are not a bounded distance from each other.

First option - Up to subsequence there are $q_i \in L_i$ with $d(q_i, F_i) \rightarrow +\infty$.

Up to covering translations we may assume that $q_i \rightarrow q_0$, so $L_i \rightarrow L_0 \in \tilde{\mathcal{F}}^s$ - the limit is unique because \mathcal{F}^s is \mathbf{R} -covered. Therefore the L_i are in compact interval I of \mathcal{H}^s . It follows that F_i are in $\eta_s^{-1}(I) \subset \mathcal{H}$, which is a compact interval of \mathcal{H} by lemma 7.3 and corollary 7.4. Up to taking a subsequence we may assume that $F_i \rightarrow F$ in \mathcal{H} . Take $z \in F$ and choose $z_i \in F_i$ with $z_i \rightarrow z$. Then

$$d(q_i, F_i) \leq d(q_i, z_i) \rightarrow d(q, z).$$

So $d(q_i, F_i)$ is bounded, contradiction.

Second option - Up to subsequence there are $p_i \in F_i$ with $d(p_i, L_i) \rightarrow +\infty$.

Again up to covering translations, assume that $p_i \rightarrow p$ so $F_i \rightarrow F$ for some $F \in \mathcal{H}$. By continuity of η_s , $L_i = \eta_s(F_i) \rightarrow \eta_s(F) = L$. The argument in the first option shows that $d(p_i, L_i)$ is bounded, again contradiction to assumption. This finishes the proof of the proposition. \square

The goal now is to show that \mathcal{F} is topologically equivalent to \mathcal{F}^s . However this is not true in general for the following easy reason: given a foliation \mathcal{F} transverse to Φ one can blow up a leaf (or a collection) of leaves of \mathcal{F} into foliated I -bundles [Ga-Oe] and still get a foliation transverse to Φ . To get the conjugacy, one first needs to blow down these gaps or foliated I -bundles. At this point we need to differentiate a bit between the two cases:

Lemma 7.6. *Suppose that Φ is a product \mathbf{R} -covered pseudo-Anosov flow. Then for any $F \in \tilde{\mathcal{F}}$, $\partial\Theta(F) = l$, with l a stable leaf in \mathcal{O} (there is no unstable boundary in $\Theta(F)$). If $F, F' \in \tilde{\mathcal{F}}$ satisfy $\eta_s(F) = \eta_s(F')$, then $\Theta(F) = \Theta(F')$.*

Proof. Suppose that $\Theta(F)$ has an unstable boundary component $u \in \tilde{\mathcal{F}}^u$. But Φ is product type, so $u \cap l \neq \emptyset$, contradiction to two boundary components of $\Theta(F)$ not intersecting. As seen before there can only one stable boundary component. This shows that $\partial\Theta(F)$ is a single stable leaf $l = \Theta(\eta_s(F))$. Hence if $\eta_s(F) = \eta_s(F')$, then $\partial\Theta(F) = \partial\Theta(F')$ and both F, F' are contained in the front of $\eta_s(F)$, so $\Theta(F) = \Theta(F')$. This finishes the proof. \square



Figure 18: a. Boundary in the product case, b. Boundary in the skewed case.

In the skewed case we have a different situation.

Proposition 7.7. *Suppose that Φ is skewed. Then for any $F \in \tilde{\mathcal{F}}$, there are unique $L \in \tilde{\mathcal{F}}^s, U \in \tilde{\mathcal{F}}^u$ with $\partial\Theta(F) = \Theta(L) \cup \Theta(U)$. In addition U and L make a perfect fit and $U = \tau_+(L)$. In particular given $F, F' \in \tilde{\mathcal{F}}$ with $\eta_s(F) = \eta_s(F')$ then $\Theta(F) = \Theta(F')$.*

Proof. Let $F \in \tilde{\mathcal{F}}$. By lemma 7.2, $\partial\Theta(F)$ can have only one stable boundary component, which is $\Theta(L)$ where $L = \eta_s(F)$. If $\Theta(F)$ does not have unstable boundary component, then $\Theta(F)$ is a half plane in \mathcal{O} bounded by $\Theta(L)$. Choose $L_i \rightarrow +\infty$ in \mathcal{H}^s , which implies that $\Theta(L_i) \subset \Theta(F)$. Hence $d_s(L, L_i) \rightarrow \infty$. Choose $p_i \in L_i \cap F$. It follows from the distance equation (2) (from the beginning of the section) applied to $p_i \in L_i$ and $L \in \tilde{\mathcal{F}}^s$, that $d(p_i, L) \rightarrow +\infty$. But $p_i \in F$ and proposition 7.5 showed that $d_H(F, L) < r_0$ so there are $z_i \in L$ with $d(p_i, z_i) < r_0$. This contradicts $d(p_i, L) \rightarrow \infty$.

We conclude that $\Theta(F)$ must have unstable boundary. In addition it can have only one unstable boundary component – the argument is the same as in the proof of proposition 7.1. Let this unstable boundary be denoted by $\eta_u(F)$. In the same way as in lemma 7.3, corollary 7.4 and proposition 7.5 it follows that η_u is continuous, weakly monotone increasing, surjective and for any $F \in \tilde{\mathcal{F}}$, then F and $\eta_u(F)$ are a bounded Hausdorff distance from each other. We stress that η_u is only defined in the skewed case (given the assumption that there is $F^* \in \tilde{\mathcal{F}}$ with $\Theta(F)$ having stable boundary).

Given $L \in \tilde{\mathcal{F}}^s$ let $J = [F_0, F_1] = \eta_s^{-1}(L)$ be a non empty, compact interval in \mathcal{H} . Assume $F_0 < F_1$ in \mathcal{H} . Given $g \in \pi_1(M)$, suppose that $g(J) \cap J \neq \emptyset$. Then some leaf $F' \in g(J)$ has $\eta_s(F') = L$. But all leaves in $g(J)$ have the same image under η_s . It follows that $g(J) \subset J$. In the same way $J \subset g(J)$. So the translates of J are either equal to J or disjoint from it. It follows that the stabilizers of L, F_0 and F_1 in $\pi_1(M)$ are exactly the same. Notice that this also works in the product case.

We now prove that if $\eta_s(F) = L$, then $\eta_u(F) = \tau_+(L)$.

We first assume that L is periodic, that is, there is a non trivial $g \in \pi_1(M)$ with $g(L) = L$. Let $J = [F_0, F_1] = \eta_s^{-1}(L)$. Then $g(F_0) = F_0$. Let $U = \tau_+(L)$, which makes a perfect fit with L and is contained in the same component of $\tilde{M} - L$ which contains F_0 . Then $g(U) = U$. Let γ be the periodic orbit of $\tilde{\Phi}$ in U so $g(\gamma) = \gamma$. If $F_0 \cap U \neq \emptyset$ then F_0 intersects all orbits in U – because the only stable boundary to $\Theta(F_0)$ is $\Theta(L)$. Hence F_0 intersects γ . Recall that the intersection of F_0 with γ can be at most one point and let this be p . Then

$$g(p) = g(F_0 \cap \gamma) = g(F_0) \cap g(\gamma) = F_0 \cap \gamma = p,$$

which obviously contradict the fact that g acts freely in \tilde{M} . It follows that $F_0 \cap U = \emptyset$. On the other hand

$$g(\eta_u(F_0)) = \eta_u(g(F_0)) = \eta_u(F_0).$$

But U is the smallest (in $\tilde{\mathcal{F}}^u$) leaf contained in $\tilde{M} - L$ and intersecting stable leaves $>_s L$, which is invariant by g . Hence either $U = \eta_u(F_0)$ or U separates $\eta_u(F_0)$ from L . The second option would imply $F_0 \cap U \neq \emptyset$, contradiction. We conclude that $U = \eta_u(F_0)$ and so $\eta_u(F_0) = \tau_+(L)$. In the same way one proves that $\eta_u(F_1) = U$ and so for any $F \in [F_0, F_1]$, then $\eta_u(F) = \eta_+(L)$.

If L is now any leaf in $\tilde{\mathcal{F}}^s$, choose $L_i \in \tilde{\mathcal{F}}^s$ periodic with $L_i \rightarrow L$. Choose $F_i \in \eta_s^{-1}(L_i)$. Then as seen before, we may assume up to subsequence that $F_i \rightarrow F$. As proved in the previous paragraph $\eta_u(F_i) = U_i = \tau_+(L_i)$. As τ_+ is continuous, then $U_i \rightarrow U = \tau_+(L)$. But η_u is also continuous, hence $\eta_u(F_i) = U_i \rightarrow U = \eta_u(F)$. It follows that $\eta_u(F) = \tau_+(L) = \tau_+(\eta_s(F))$ for an arbitrary $F \in \tilde{\mathcal{F}}$.

Finally this shows that given any $F, F' \in \tilde{\mathcal{F}}$, if $\eta_s(F) = \eta_s(F')$ then $\eta_u(F) = \eta_u(F')$ also. It follows that $\Theta(F) = \Theta(F')$, that is, F and F' intersect exactly the same orbits of $\tilde{\Phi}$. This finishes the proof of the proposition. \square

Given any foliation \mathcal{F}^* transverse to Φ and $F \in \widetilde{\mathcal{F}}^*$, let $\partial^s \Theta(F)$ be the union of the stable boundary components of $\Theta(F)$ and similarly define $\partial^u \Theta(F)$. In our situation of \mathcal{F} transverse to Φ , $\partial^s \Theta(F) = \Theta(\eta_s(F))$, $\partial^u \Theta(F) = \Theta(\eta_u(F))$ (the second term only occurs in the skewed case).

Proposition 7.8. *There is a foliation \mathcal{F}' , which is obtained from \mathcal{F} by blowing down at most countably many foliated I -bundles of \mathcal{F} to single leaves and so that \mathcal{F}' satisfies: \mathcal{F}' is transverse to Φ and for any $L \in \widetilde{\mathcal{F}}^s$ there is a unique $F \in \widetilde{\mathcal{F}}'$ with $\Theta(L) = \partial^s \Theta(F)$.*

Proof. Suppose there is $L \in \widetilde{\mathcal{F}}^s$ and there are $F, F' \in \widetilde{\mathcal{F}}$ with $\Theta(L) = \partial^s \Theta(F) = \eta_s(F) = \partial^s \Theta(F') = \eta_s(F')$. Let $J = \eta_s^{-1}(L) = [F_0, F_1]$, which is a compact, non degenerate interval in $\mathcal{H}(\widetilde{\mathcal{F}}^s)$. Assume $F_0 <_s F_1$. By proposition 7.5, the Hausdorff distances from L to F_0 and from L to F_1 are bounded by r_0 . It follows that the Hausdorff distance from F_0 to F_1 is bounded by $2r_0$.

By lemma 7.6 and proposition 7.7, then for any $F, F' \in J$, $\Theta(F) = \Theta(F')$, that is F and F' intersect exactly the same orbits of $\widetilde{\Phi}$. There is a homeomorphism from F_0 to F_1 obtained by following orbits of $\widetilde{\Phi}$. Also for any $g \in \pi_1(M)$, $g(J) = J$ or $g(\widetilde{J}) \cap J = \emptyset$, so the stabilizers of F_0 and F_1 are equal to a subgroup \mathbf{A} of $\pi_1(M)$. Let V be the closed subset of M bounded by $F_0 \cup F_1$. The conditions above imply that V/\mathbf{A} embeds in M and is homeomorphic to $\pi(F_0) \times [0, 1]$ with $\pi(F_0)$ and $\pi(F_1)$ corresponding to levels 0 and 1 respectively and following orbit segments of Φ produces a homeomorphism from $\pi(F_0)$ to $\pi(F_1)$. Therefore V/\mathbf{A} is a foliated product so that transverse segments are given by orbits of Φ . Collapse $\pi(F_0)$ to $\pi(F_1)$ by identifying those segments to points. This is the inverse operation of blowing up a single leaf. There are at most countably many such regions, corresponding exactly to the non degenerate intervals of \mathcal{H} where η_s is constant. Do the blow down operation for the at most countable foliated products as above to produce a foliation \mathcal{F}' . Then \mathcal{F}' is still transverse to the flow Φ . In addition for any $F, F' \in \widetilde{\mathcal{F}}'$, if $\partial^s(F) = \partial^s(F')$ then $F = F'$ so \mathcal{F}' satisfies the requirements of the proposition. \square

The goal now is to show that \mathcal{F}' is topologically conjugate to \mathcal{F}^s . First we need a couple of facts. Given a leaf F of a foliation \mathcal{F} let d_F be the path metric in F - $d_F(x, y)$ is the infimum of the lengths of paths from x to y contained in F . The metric d will always denote the original metric in M or \widetilde{M} .

Lemma 7.9. *([Fe2]) Let \mathcal{F} be an \mathbf{R} -covered foliation in M^3 closed. Then for any $u > 0$ there is $v(u) > 0$ satisfying: for any leaf G of $\widetilde{\mathcal{F}}$ and any two points $x, y \in G$:*

$$d(x, y) < u \quad \Rightarrow \quad d_G(x, y) < v(u) \quad (3)$$

The point here is that $v(u)$ depends only on \mathcal{F} and u and not on G, x or y .

Definition 7.10. *Let $\varphi : (M_1, d_1) \rightarrow (M_2, d_2)$ be a map between metric spaces. Then φ is a quasi-isometric embedding [Th2, Gr] if there is $k > 0$ so that for any $x, y \in M_1$ then*

$$\frac{1}{k} d_1(x, y) - k < d_2(\varphi(x), \varphi(y)) - k < k d_1(x, y) + k. \quad (4)$$

If in addition there is k' so that for any

$$\forall z \in M_2, \quad \exists x \in M_1, \quad \text{with} \quad d_2(z, \varphi(x)) < k' \quad (5),$$

then φ is called a quasi-isometry between M_1 and M_2 . If the constant is important then we say that φ is a k quasi-isometry.

In the case of quasi-isometry it easily follows that there is $\varphi' : M_2 \rightarrow M_1$ which is also a quasi-isometry and which is an inverse of φ in the quasi-isometry category: $\varphi' \circ \varphi$ is a bounded distance from the identity map of M_1 and similarly for $\varphi \circ \varphi'$. Notice that is NOT at all required that the quasi-isometry φ is continuous – one is only interested in what happens for large distances. For background on quasi-isometries we refer the reader to [Gr, Gh-Ha, CDP].

Let \mathcal{H}' be the leaf space of $\widetilde{\mathcal{F}}'$ which is obviously homeomorphic to \mathbf{R} . Clearly for any $F' \in \widetilde{\mathcal{F}}'$ there is $F \in \widetilde{\mathcal{F}}$ with $\Theta(F) = \Theta(F')$. Hence $\partial^s(F') = \partial^s(F) = \eta_s(F)$. Define $\eta(F') = \eta_s(F)$. By proposition 7.8, the map η is well defined and η is a homeomorphism from \mathcal{H}' to \mathcal{H}^s .

Since \mathcal{F}' is obtained from \mathcal{F} by collapsing at most countably many foliated products, there is a collapsing map $\beta : M \rightarrow M$ sending leaves of \mathcal{F} to leaves of \mathcal{F}' and collapsing the required foliated products. The map β is continuous and is a homotopy equivalence which is homotopic to the identity. It lifts to a map $\widetilde{\beta} : \widetilde{M} \rightarrow \widetilde{M}$, which leaves orbits of $\widetilde{\Phi}$ invariant and moves points of \widetilde{M} a bounded distance. Hence the Hausdorff distance $d_H(\widetilde{\beta}(F), F)$

is finite and bounded for any $F \in \tilde{\mathcal{F}}$. Let $F' = \beta(F)$. Since $d_H(F, \eta_s(F)) < r_0$ and $\eta(F') = \eta_s(F)$, it follows that $d_H(F', \eta(F'))$ is globally bounded by a constant r_1 for any $F' \in \mathcal{F}'$.

We begin by relating leaves of $\tilde{\mathcal{F}}'$ and $\tilde{\mathcal{F}}^s$ metrically.

Lemma 7.11. *For any $F \in \tilde{\mathcal{F}}'$ and $\eta(F) \in \tilde{\mathcal{F}}^s$ there is a roughly natural map $\varphi : \eta(F) \rightarrow F$ which is a uniform quasi-isometry.*

Proof. The term roughly natural will become clear in the proof. Both \mathcal{F} and \mathcal{F}' are \mathbf{R} -covered. Let v be the function given by lemma 7.9 for \mathcal{F}^s and v' the respective function for \mathcal{F}' . Given $F \in \tilde{\mathcal{F}}'$ let $L = \eta(F)$. Then $d_H(F, L) < r_1$. So given $p \in L$ choose a point $q \in F$ with $d(p, q) < r_1$. Let this be $\varphi(p)$. Obviously $\varphi(p)$ is not uniquely defined, but we show it is roughly uniquely defined in F as follows: if $q' \in F$ with $d(q', p) < r_1$ then $d(q, q') < 2r_1$. Let $a_0 = v'(2r_1)$. By lemma 7.9 $d_F(q, q') < v'(2r_1) = a_0$, which is a fixed constant. So the diameter of the set of possible choices for $\varphi(p)$ is globally bounded and $\varphi(p)$ is well defined in F up to a distance a_0 . Notice that clearly there is no guarantee that φ is even continuous.

Given $x, y \in L$ let $n = \lfloor d_L(x, y) \rfloor$ where $\lfloor \cdot \rfloor$ denotes the function greatest integer less than or equal the given number. Choose a minimal length geodesic γ in L from x to y and choose points p_0, p_1, \dots, p_{n+1} in γ so that $p_0 = x$, $p_{n+1} = y$, $d_L(p_i, p_{i+1}) = 1$ if $0 \leq i < n$ and $d_L(p_n, p_{n+1}) < 1$. Let $q_i = \varphi(p_i)$. Then $d(q_i, p_i) < r_1$, hence $d(q_i, q_{i+1}) < 2r_1 + 1$. Let $a_1 = v'(2r_1 + 1)$. By lemma 7.9 $d_F(q_i, q_{i+1}) < a_1$. Therefore

$$d_F(\varphi(x), \varphi(y)) = d_F(q_0, q_{i+1}) < (n+1)a_1 = na_1 + a_1 \leq a_1 d_L(x, y) + a_1. \quad (6)$$

This proves one side of the needed quasi-isometric inequality (4) for the map φ . To prove the other direction we will use a dual argument.

Given $w \in F$ choose $z \in \eta(F) = L$ with $d(z, w) < r_1$. Let this z be $\varphi'(w)$. As above the function φ' is (roughly) natural – its possible values are in a set of diameter $< 2r_1$ in \tilde{M} and hence of diameter $< v(2r_1)$ in L .

Let $a_2 = v(2r_1 + 1)$. An argument entirely analogous as above shows that for any $z, w \in F$ then

$$d_L(\varphi'(z), \varphi'(w)) \leq a_2 d_F(z, w) + a_2. \quad (7)$$

Notice that for any $z \in F$ then $\varphi(z) \in L$ and $d(z, \varphi'(z)) < r_1$. Let $y = \varphi'(z)$. Then $\varphi(y) \in F$ and $d(z, \varphi(y)) < 2r_1$. So $d_F(z, \varphi(y)) < v'(2r_1) = a_0$. Since z is arbitrary in F and F arbitrary in \mathcal{H} , this proves the second condition (5) of the quasi-isometry definition for φ with $k' = a_0$. In addition for any $z \in F$, $d_F(z, \varphi \circ \varphi'(z)) < v'(2r_1) = a_0$. Let $a_3 = v(2r_1)$. In a completely analogous fashion, for any $x \in L$ then

$$d_L(x, \varphi' \circ \varphi(x)) < v(2r_1) = a_3. \quad (8)$$

Finally we show the other side of the required inequality (4). Let $x, y \in L$ arbitrary. Then (7) applied to $z = \varphi(x), w = \varphi(y)$ shows that

$$d_L(\varphi' \circ \varphi(x), \varphi' \circ \varphi(y)) \leq a_2 d_F(\varphi(x), \varphi(y)) + a_2. \quad (9)$$

But $d_L(\varphi' \circ \varphi(x), x) < a_3$ and similarly for y , hence

$$d_L(x, y) < a_2 d_F(\varphi(x), \varphi(y)) + (a_2 + 2a_3). \quad (10)$$

or

$$\frac{1}{a_2} d_L(x, y) - \left(1 + \frac{2a_3}{a_2}\right) < d_F(\varphi(x), \varphi(y)). \quad (11)$$

Choose $k = \max(a_1, a_2, 1 + \frac{2a_3}{a_2})$. Then equation (4) holds for any $x, y \in L$. Therefore φ is a k quasi-isometry from L to F . This finishes the proof of the proposition. \square

We now produce a flow Φ' tangent to \mathcal{F}' which “shadows” Φ in a particular way: for each orbit of $\tilde{\Phi}$ there is an orbit of $\tilde{\Phi}'$ which is a bounded Hausdorff distance from it. Recall that a *quasigeodesic* in a metric space is a quasi-isometric embedding of \mathbf{R} or \mathbf{Z} (with their natural metrics).

Proposition 7.12. (*shadowing flow*) *There is a flow Φ' in M tangent to \mathcal{F} which shadows Φ : for any orbit γ of $\tilde{\Phi}$ there is unique orbit γ' of $\tilde{\Phi}'$ associated to it and γ' is is a bounded distance from γ .*




Figure 19: *The shadowed flow in $\tilde{\mathcal{F}}'$.*

Proof. It is well known that the leaves of $\tilde{\mathcal{F}}^s$ are Gromov negatively curved in the large in their path metrics [Su, P11, Gr]. Therefore each leaf L of $\tilde{\mathcal{F}}^s$ has an ideal boundary $\partial_\infty L$ which is then homeomorphic to a circle [Fe2] and $L \cup \partial_\infty L$ is homeomorphic to a closed disk [Fe2]. Given $F \in \tilde{\mathcal{F}}'$ then F is k quasi-isometric to $\eta(F) = L$ – it follows that F is also Gromov negatively curved in the large [Gr, Gh-Ha, CDP].

Since all leaves of $\tilde{\mathcal{F}}'$ are Gromov negatively curved, Candel [Can] showed that there is a metric d' in M making leaves of \mathcal{F}' hyperbolic. The metric d' is obviously smooth along the leaves of \mathcal{F}' but may be only C^0 transversely. Here we use both metrics d and d' . Let d'_F be the induced path metric in each leaf of \mathcal{F}' – it is a hyperbolic metric. Then the metric d_F and d'_F are uniformly quasi-isometric, that is, the identity map $(F, d_F) \rightarrow (F, d'_F)$ is a uniform quasi-isometry. Uniform means that the quasi-isometric constant does not depend on F and this follows from compactness of M .

Let $L = \eta(F)$. The flow lines of $\tilde{\Phi}$ in L are uniform quasigeodesics in L [Fe3] and therefore limit to two distinct ideal points in $\partial_\infty L$ [Fe3]. In fact there is a common forward ideal point $b(L)$ of all flow lines (they are forward asymptotic) and all other points of $\partial_\infty L$ are negative ideal points of flow lines of $\tilde{\Phi}$ in L [Fe3].

An important fact about Gromov negatively curved spaces is that since $\varphi : L \rightarrow F$ is a quasi-isometry, it extends to a homeomorphism $\tilde{\varphi} : L \cup \partial_\infty L \rightarrow F \cup \partial_\infty F$ [Gr, Gh-Ha, CDP]. This defines a point $b(F) \in \partial_\infty F$ which is the image of $b(L)$ under this map. The point $b(F)$ is uniquely defined in $\partial_\infty F$. We now construct a flow $\tilde{\Phi}'$ in \tilde{M} as follows: its orbits are contained in leaves of $\tilde{\mathcal{F}}'$. For each $F \in \tilde{\mathcal{F}}'$ the orbits are geodesics in the hyperbolic metric d'_F of F . They all have one ideal point in $b(F)$ and move toward this ideal point with speed 1.

We first need to show that this defines a continuous flow in \tilde{M} . In each leaf of $\tilde{\mathcal{F}}'$ it is obviously smooth but we need to consider what happens transversally. Given $p \in \tilde{M}$ let $D\tilde{\Phi}'(p)$ be the tangent vector field to the flow lines. We aim to prove that $D\tilde{\Phi}'$ is continuous with p .

Let \mathcal{O}' be the orbit space of $\tilde{\Phi}'$. Before proving continuity we produce a map from orbits of $\tilde{\Phi}$ to orbits of $\tilde{\Phi}'$ or equivalently a map from \mathcal{O} to \mathcal{O}' . Here we identify orbits of $\tilde{\Phi}$ to points of \mathcal{O} and similarly for \mathcal{O}' . Given γ and orbit of $\tilde{\Phi}$, it is contained in $L \in \tilde{\mathcal{F}}^s$. Let $F = \eta^{-1}(L) \in \tilde{\mathcal{F}}'$. Notice that F is unique. Then γ is a uniform quasigeodesic in L [Fe3] and as $\varphi : L \rightarrow F$ is a quasi-isometry then $\varphi(\gamma)$ is a quasigeodesic in F . Since φ is not continuous, $\varphi(\gamma)$ may not be a continuous curve. Another point of view is to choose a sequence of points in γ so that consecutive points are 1 unit apart in γ . The image is a quasi-isometric embedding from \mathbf{Z} in F and its image is a bounded distance from a unique geodesic in F [Gr, Gh-Ha, CDP]. One of the ideal points is obviously $b(F)$ and the other depends on γ . There is a unique geodesic in F which has these two ideal points and let this be $\chi(\gamma)$. The geodesic $\chi(\gamma)$ is an orbit of $\tilde{\Phi}'$ and this defines a map $\chi : \mathcal{O} \rightarrow \mathcal{O}'$. Then $\chi(\gamma)$ is a bounded Hausdorff distance from $\varphi(\gamma)$ and hence from γ also. In fact $\chi(\gamma)$ is the unique geodesic in F which is bounded distance from γ in F and hence in \tilde{M} . The bound is global for all orbits of $\tilde{\Phi}$. The map χ sends the set of orbits of $\tilde{\Phi}$ in L bijectively to the set of orbits of $\tilde{\Phi}'$ in F . Finally $\chi : \mathcal{O} \rightarrow \mathcal{O}'$ is a bijection.

We now prove continuity of $D\tilde{\Phi}'$. Let $p_i \rightarrow p$ in \tilde{M} . Let α_i be the orbits of $\tilde{\Phi}'$ through p_i . We can choose a subsequence so that $D\tilde{\Phi}'(p_i)$ converges to a tangent vector w_0 at p . Let $p_i \in F_i \in \tilde{\mathcal{F}}'$ and $p \in F \in \tilde{\mathcal{F}}'$. Let α be the geodesic of F through p in the direction of w_0 . Then every point of α is a limit of points in α_i . Let $L_i = \eta(F_i)$ and $\gamma_i \subset L_i$ orbits of $\tilde{\Phi}$ which are a uniform bounded distance from the orbits α_i . By the construction above the

γ_i are unique. Also $L_i \rightarrow L$ with $L = \eta(F)$. The γ_i all have points q_i a bounded distance from p_i so up to a further subsequence we may assume that $q_i \rightarrow q$ and γ_i converge to an orbit γ of $\tilde{\Phi}$ in F . For any fixed size r_2 , the segments of size r_2 of γ_i centered at q_i converge uniformly to the segment of size r_2 of γ . But these segments are a bounded distance from segments of size $z(r_2)$ of α_i centered at p_i , where $z(r_2) \rightarrow \infty$ when $r_2 \rightarrow \infty$. But segments of fixed size in α_i converge uniformly to segments in α , so for any fixed size segment in α it is a bounded distance from a segment in γ . In addition the bound is independent of the segment or its size. It follows that the whole orbit α is a bounded distance from γ . But the only geodesic of F which is a bounded distance from the orbit γ of $\tilde{\Phi}$ is $\chi(\gamma)$. Hence $\alpha = \chi(\gamma)$ and the tangent to α at p is $D\tilde{\Phi}'(p)$ which is w_0 . This proves continuity of $D\tilde{\Phi}'$ and shows that $\tilde{\Phi}'$ is a continuous flow in \tilde{M} .

Furthermore the function $b(F)$ is clearly equivariant under covering translations: if $g \in \pi_1(M)$ and $F \in \tilde{\mathcal{F}}'$ then $b(g(F)) = g(b(F))$ – where the second g denotes the extension of the isometry $g : F \rightarrow g(F)$ to the ideal boundaries of F and $g(F)$. It follows that $\tilde{\Phi}'$ induces a flow Φ' in M . Notice that χ is a homeomorphism from \mathcal{O} to \mathcal{O}' .

Finally for any $F \in \tilde{\mathcal{F}}'$, $\eta(F) = L \in \tilde{\mathcal{F}}^s$, the maps $\partial_\infty L \rightarrow \partial_\infty F$ are also equivariant, hence the map $\chi : \mathcal{O} \rightarrow \mathcal{O}'$ is also group equivariant. \square

We now prove topological equivalence of \mathcal{F}' and \mathcal{F}^s .

Theorem 7.13. *The flows Φ and Φ' are topologically conjugate. The foliation \mathcal{F}' is topologically equivalent to the stable foliation \mathcal{F}^s of Φ .*

Proof. By the previous proposition there is a homeomorphism $\chi : \mathcal{O} \rightarrow \mathcal{O}'$. This homeomorphism is group equivariant. There is a general theory which deals with such situations [Hae]: the theory of classifying spaces for flows produces a map $h : M \rightarrow M$ which is a homotopy equivalence and sends orbits of Φ to orbits of Φ' and is transversely injective. The problem is that h may not be injective along the orbits. Ghys [Gh] encountered this situation when studying Anosov flows in Seifert fibered spaces and did an averaging trick to produce an injective map, hence topological equivalence. Barbot did the same for general \mathbf{R} -covered Anosov flows [Ba2]. We cannot apply these results directly because we do not know that the flow Φ' is Anosov, let alone that it is a smooth flow. On the other hand there is much more structure in our situation which allows us to get the topological equivalence more directly.

In our situation the homeomorphism $\chi : \mathcal{O} \rightarrow \mathcal{O}'$ and the quasi-isometries $\varphi : L \rightarrow \eta^{-1}(L)$ give a way to relate the orbits and points in the orbits. Let \mathcal{U} be a very fine, locally finite foliation box cover (of \mathcal{F}^s) of M . Choose a partition of unit subordinated to this cover. For each element of the cover, lift it to \tilde{M} and define a map from this set to a foliated box of \mathcal{F}' by sending each plaque of $\tilde{\mathcal{F}}^s$ to a plaque of $\tilde{\mathcal{F}}'$, each orbit segment of $\tilde{\Phi}$ to an orbit segment of $\tilde{\Phi}'$ so that this is a homeomorphism between the two sets and monotonic increasing along orbits. The final requirement is that any point and its image are $< r_3$ distant apart, where r_3 is a globally fixed constant. This is doable because $d_H(\gamma, \chi(\gamma))$ is bounded over all orbits of $\tilde{\Phi}$. Iterate by covering translations of \tilde{M} . Do this for all elements of the foliated box cover of M . Given $p \in \tilde{M}$ with $p \in \gamma$ orbit of $\tilde{\Phi}$, then the above construction defines finitely many images of p in $\chi(\gamma)$. Each orbit of $\tilde{\Phi}'$ has a well defined affine structure: use the partition of unity as affine coordinates of the images of p in the orbit $\chi(\gamma)$ of $\tilde{\Phi}'$ to define a map $\tilde{h} : \tilde{M} \rightarrow \tilde{M}$ which sends orbits of $\tilde{\Phi}$ to orbits of $\tilde{\Phi}'$ and induces $\chi : \mathcal{O} \rightarrow \mathcal{O}'$. This map is continuous and equivariant, producing a map $h : M \rightarrow M$ which is a homotopy equivalence, sends orbits of Φ to orbits of Φ' and is transversely injective.

Furthermore the construction implies that for any $p \in \tilde{M}$, then $d(p, \tilde{h}(p)) < r_4$ for a globally fixed constant r_4 . Therefore even though \tilde{h} may not be monotonic in a fixed orbit of $\tilde{\Phi}$, it eventually progresses in the forward direction. This is called by Thurston as *quasi-monotonic* [Th5] and it is the key property which allows for the construction of a monotonic map [Ba2] (see details of the proof of theorem 3.4 of [Ba2]). The new map is obtained by averaging h on orbit segments of long but fixed size t . This provides a new map $h' : M \rightarrow M$, which is now injective along orbits and is therefore a topological equivalence from Φ to Φ' .

This immediately shows that \mathcal{F}' is topologically equivalent to \mathcal{F}^s , finishing the proof of the theorem. \square

We collect these results together:

Corollary 7.14. *Let Φ be an \mathbf{R} -covered pseudo-Anosov flow transverse to a foliation \mathcal{F} and suppose that Φ is not regulating for \mathcal{F} . Then \mathcal{F} is topologically equivalent to a blow up of either (1) the stable foliation \mathcal{F}^s or (2) the unstable foliation \mathcal{F}^u of Φ .*

Proof. Suppose first that some leaf $F \in \widetilde{\mathcal{F}}$ has $\partial^s(\Theta(F)) \neq \emptyset$. Then the results of this section show that conclusion (1) occurs. If on the other hand $\partial^u(\Theta(F)) \neq \emptyset$, the same arguments show that (2) occurs. This finishes the proof. Notice that in the skewed case \mathcal{F}^s and \mathcal{F}^u are topologically equivalent [Ba2]. \square

Finally we also mention a topological rigidity for the flows: Let \mathcal{F} be an \mathbf{R} -covered foliation in M^3 closed. Then, up to topological conjugacy, there is at most one pseudo-Anosov flow Φ which is transverse to \mathcal{F} , but not regulating. The proof is very similar to the arguments given in this section and is omitted for brevity.

References

- [An] D. V. Anosov, *Geodesic flows on closed Riemannian manifolds with negative curvature*, Proc. Steklov Inst. Math. **90** (1969).
- [An-Si] D. V. Anosov and Y. Sinai, *Some smoothly ergodic systems*, Russian Math. Surveys **22** (1967) 5 103-167.
- [Ba1] T. Barbot, *Géométrie transverse des flots*, Thesis, École Normale Sup. Lyon, 1992.
- [Ba2] T. Barbot, *Caractérisation des flots d'Anosov en dimension 3 par leurs feuilletages faibles*, Ergod. Th. Dynam. Sys. **15** (1995) 247-270.
- [Ba3] T. Barbot, *Flots d'Anosov sur les variétés graphées au sens de Waldhausen*, Ann. Inst. Fourier Grenoble **46** (1996) 1451-1517.
- [Ba4] T. Barbot, *Generalizations of Bonatti-Langevin example of Anosov flow and their classification up to topological equivalence*, Comm. Anal. Geom. **6** (1998) 749-798.
- [Bl-Ca] S. Bleiler and A. Casson, *Automorphisms of surfaces after Nielsen and Thurston*, Cambridge Univ. Press, 1988.
- [Bow1] R. Bowen, *Periodic orbits for hyperbolic flows*, Amer. Jour. Math. **94** (1970) 1-30.
- [Bow2] R. Bowen, *Symbolic dynamics for hyperbolic flows*, Amer. Jour. Math. **95** (1970) 429-459.
- [Br] M. Brittenham, *Essential laminations in Seifert fibered spaces*, Topology **32** (1993) 61-85.
- [Ca1] D. Calegari, *\mathbf{R} -covered foliations of hyperbolic 3-manifolds*, Geom. Topol **3** (1999) 137-153.
- [Ca2] D. Calegari, *The geometry of \mathbf{R} -covered foliations*, Geom. Topol. **4** (2000) 457-515.
- [Ca3] D. Calegari, *Foliations with one sided branching*, to appear.
- [Ca4] D. Calegari, *Promoting essential laminations, Part I*, preprint, 2001.
- [Can] A. Candel, *Uniformization of surface laminations*, Ann. Sci. Ecole Norm. Sup. (4) **26** (1993) 489-516.
- [CDP] M. Coornaert, T. Delzant and A. Papadopoulos, *Géométrie et théorie des groupes, Les groupes hyperboliques de Gromov*, Lecture Notes in Math. **1441** Springer (1991).
- [Ca-Th] J. Cannon and W. Thurston, *Group invariant Peano curves*, to appear.
- [CLR1] D. Cooper, D. Long and A. Reid, *Bundles and finite foliations*, Inven. Math. **118** (1994) 255-283.
- [CLR2] D. Cooper, D. Long and A. Reid, *Finite foliations and similarity interval exchange maps*, Topology, **36** (1997) 209-227.
- [FLP] A. Fathi, Laudenbach, Poenaru, *Travaux de Thurston sur les surfaces*, Séminaire Orsay, Astérisque **66-67** Société Mathématique de France, 1979.
- [Fe1] S. Fenley, *Asymptotic properties of depth one foliations in hyperbolic 3-manifolds*, Jour. Diff. Geom., **36** (1992) 269-313.
- [Fe2] S. Fenley, *Quasi-isometric foliations*, Topology **31** (1992) 667-676.
- [Fe3] S. Fenley, *Anosov flows in 3-manifolds*, Ann. of Math. **139** (1994) 79-115.
- [Fe4] S. Fenley, *Quasigeodesic Anosov flows and homotopic properties of closed orbits*, Jour. Diff. Geo. **41** (1995) 479-514.
- [Fe5] S. Fenley, *The structure of branching in Anosov flows of 3-manifolds*, Comm. Math. Helv. **73** (1998) 259-297.
- [Fe6] S. Fenley, *Continuous extension of Anosov foliations in 3-manifolds with negatively curved fundamental group*, Pac. Jour. Math. **186** (1998) 201-216.
- [Fe7] S. Fenley, *Surfaces transverse to pseudo-Anosov flows and virtual fibers in 3-manifolds*, Topology **38** (1999) 823-859.
- [Fe8] S. Fenley, *Foliations with good geometry*, Journal of the A.M.S. **12** (1999) 619-676.
- [Fe9] S. Fenley, *Pseudo-Anosov flows and incompressible tori*, submitted for publication.
- [Fe10] S. Fenley, *Foliations and the topology of 3-manifolds: \mathbf{R} -covered foliations and transverse pseudo-Anosov flows*, to appear.
- [Fe11] S. Fenley, *Topological rigidity of pseudo-Anosov flows transverse to foliations*, in preparation.
- [Fe-Mo] S. Fenley and L. Mosher, *Quasigeodesic flows in hyperbolic 3-manifolds*, Topology **40** (2001) 503-537.
- [Fr-Wi] J. Franks and R. Williams, *Anomalous Anosov flows*, in Global theory of Dyn. Systems, Lecture Notes in Math. **819** Springer (1980).
- [Ga1] D. Gabai, *Foliations and the topology of 3-manifolds*, J. Diff. Geo. **18** (1983) 445-503.
- [Ga2] D. Gabai, *Foliations and the topology of 3-manifolds III*, J. Diff. Geo. **26** (1987) 479-536.
- [Ga-Oe] D. Gabai and U. Oertel, *Essential laminations and 3-manifolds*, Ann. of Math. **130** (1989) 41-73.
- [Gh] E. Ghys, *Flots d'Anosov sur les 3-variétés fibrées en cercles*, Ergod. Th and Dynam. Sys. **4** (1984) 67-80.
- [Gh-Ha] E. Ghys and P. de la Harpe, editors, *Sur les groupes hyperboliques d'après Mikhael Gromov*, Progress in Math. **83** Birkhäuser, (1996).
- [Go] S. Goodman, *Closed leaves in foliations of codimension one*, Comm. Math. Helv. **50** (1975) 383-388.
- [Go-Sh] S. Goodman and S. Shields, *A condition for the stability of \mathbf{R} -covered for foliations of 3-manifolds*, to appear in Trans. A.M.S. (2000).

- [Gr] M. Gromov, *Hyperbolic groups*, in Essays on group theory, Springer, 1987, 75-263.
- [Hae] A. Haefliger, *Groupoïdes d'holonomie et classifiants*, Asterisque **116** (1984) 70-97.
- [Han] M. Handel, *Global shadowing of pseudo-Anosov homeomorphisms*, Ergod. Th. Dynam. Sys. **5** (1985) 373-377.
- [He] J. Hempel, *3-manifolds*, Ann. of Math. Studies 86, Princeton University Press, 1976.
- [Ko] N. Kopell, *Commuting diffeomorphisms*, Proceedings of Symposia in Pure Mathematics, A.M.S. **14** (1970) 165-184.
- [Le] G. Levitt, *Feuilletages des variétés de dimension trois qui son fibrés en cercles*, Comm. Math. Helv. **53** (1978) 572-594.
- [Man] B. Mangum, *Incompressible surfaces and pseudo-Anosov flows*, Topol. Appl. **87** (1998) 29-51.
- [Mo1] L. Mosher, *Dynamical systems and the homology norm of 3-manifolds, I*, Duke Jour. Math. **65** (1992) 449-500.
- [Mo2] L. Mosher, *Laminations and flows transverse to finite depth foliations*, manuscript available in the web from <http://newark.rutgers.edu:80/mosher/>, Part I: Branched surfaces and dynamics, Part II in preparation.
- [No] S. P. Novikov, *Topology of foliations*, Trans. Moscow Math. Soc. **14** (1963) 268-305.
- [Pl1] J. Plante, *Foliations with measure preserving holonomy*, Ann. of Math. **107** (1975) 327-361.
- [Pl2] J. Plante, *Foliations of 3-manifolds with solvable fundamental group*, Inven. Math. **51** (1979) 219-230.
- [Pl3] J. Plante, *Solvable groups acting on the line*, Trans. A.M.S. **278** (1983) 401-414.
- [Pl4] J. Plante, *Anosov flows, transversely affine and a conjecture of Verjovsky*, Jour. London Math. Soc. **83** (1981) 359-362.
- [Pl-Th] J. Plante and W. Thurston, *Anosov flows and the fundamental group*, Topology **11** (1972) 147-150.
- [Rob1] R. Roberts, *Taut foliations in punctured surface bundles, I*, to appear.
- [Rob2] R. Roberts, *Taut foliations in punctured surface bundles, II*, to appear.
- [Rol] D. Rolfsen, *Knots and links*, Publish or perish, 1976.
- [Ros] H. Rosenberg, *Foliations by planes*, Topology **7** (1968) 131-138.
- [Sc] P. Scott, *Subgroups of surface groups are almost geometric*, Jour. Lond. Math. Soc. **17** (1978) 555-565..
- [Se] A. Selberg, *On discontinuous groups in higher-dimensional symmetric spaces*, in Contributions to Function Theory, Tata Inst. of Fund. Research, Bombay (1960) 147-164.
- [St] K. Strebel, *Quadratic differentials*, Springer-Verlag, 1984.
- [Su] D. Sullivan, *Cycles for the dynamical study of foliated manifolds and complex manifolds*, Invent. Math. **36** (1976) 225-255.
- [Th1] W. Thurston, *Foliations of 3-manifolds that are circle bundles*, PhD thesis, University of California Berkeley, 1972.
- [Th2] W. Thurston, *The geometry and topology of 3-manifolds*, Princeton University Lecture Notes, 1982.
- [Th3] W. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. A.M.S **19** (1988) 417-431.
- [Th4] W. Thurston, *Hyperbolic structures on 3-manifolds II, Surface groups and 3-manifolds that fiber over the circle*, preprint.
- [Th5] W. Thurston, *Three manifolds, foliations and circles I*, preprint, 1997.
- [Th6] W. Thurston, *Three manifolds, foliations and circles II, The transverse asymptotic geometry of foliations*, preprint, 1998.
- [Th7] W. Thurston, *Private communication*.
- [Ve] A. Verjovsky, *Codimension one Anosov flows*, Bol. Soc. Mat. Mex. **19** (1977) 49-77.

Florida State University
Tallahassee, FL 32306-4510