



Demonstrates $3 \cdot 2 = 2 \cdot 3$ by sending apple (x, y) to (y, x) .
The same idea is used in Exercise 6.

Intro Advanced Math. Answers to questions 1–8

1. For each, simplify the cardinality to one of: $0, 1, 2, \dots, \aleph_0, c, 2^c, 2^{2^c}, \dots$

For (a)–(h) you do not need to show your work, but for (i),(j) you need to justify your answer by showing all steps.

- (a) \mathbb{N} : \aleph_0
 - (b) $\emptyset \times \mathbb{R}$: $0 \cdot c = 0$
 - (c) \mathbb{Q} : \aleph_0
 - (d) $\mathbb{R} \times P(\mathbb{Q})$: $c \cdot 2^{\aleph_0} = c \cdot c = c$
 - (e) $\mathbb{Q} - \mathbb{Z}$: \aleph_0
 - (f) $P(\mathbb{N})$: $2^{\aleph_0} = c$
 - (g) $P(\mathbb{R})$: 2^c
 - (h) $\{2, 2\}$: 1
 - (i) $\mathbb{R}^{\mathbb{N}}$: $c^{\aleph_0} = (2^{\aleph_0})^{\aleph_0} = 2^{\aleph_0 \aleph_0} = 2^{\aleph_0} = c$
 - (j) $\mathbb{R}^{\mathbb{R}}$: $c^c = (2^{\aleph_0})^c = 2^{\aleph_0 c} = 2^c$
2. Based on the answer in your previous question, does there exist:
(it suffices to write yes/no):
- (a) an injective function from $\mathbb{R}^{\mathbb{R}}$ to $P(\mathbb{R})$? Yes
 - (b) an injective function from \mathbb{Q} to \mathbb{N} ? Yes
 - (c) an injective function from $P(\mathbb{N})$ to \mathbb{N} ? No
3. Prove, using only the definition, that the intervals $(0, 1)$ and $(0, 2)$ have the same cardinality.
- The function $f(x) = 2x$ is a bijection from $(0, 1)$ to $(0, 2)$.
[must know item 1!]
4. Let A, B be sets and let $C = A \cup B$. Suppose that $A \cap B = \emptyset$ and:
- (1) there is **no** bijection from A to C
 - (2) there is **no** bijection from B to C

Prove that A and B are finite sets.

Let a, b, c_0 be the cardinalities of A, B, C . Then $c_0 = a + b$ (item 21).

Suppose a or b is infinite. Then $c_0 = \max(a, b)$ (item 22).

So c_0 is a or b . But $c_0 \neq a$ by (1) and $c_0 \neq b$ by (2), contradiction.

5. Let A be any set. Prove that there is no bijection from \mathbb{N} to $P(A)$.

Suppose there is a bijection. Then $o(P(A)) = \aleph_0$ [memorize items 1-6].

Item 7 says $o(A) < o(P(A))$, so $o(A) < \aleph_0$.

Then A is finite (items 11, 10, 6, 5). Then $P(A)$ is finite, contradiction.

6. TURN IN: We know that if d, e are natural numbers then $d \cdot e = e \cdot d$. But do you remember how to prove that? Lets prove this not only for natural numbers, but for all cardinal numbers! I will type the first line in the proof, and you finish it:

Proof: Let D, E be sets for which $d = o(D)$ and $e = o(E)$.

- (a) Give the definition of $D \times E$.

This is the set of all pairs (x, y) with $x \in D$ and $y \in E$.

The shortest way to denote that is: $\{(x, y) : x \in D, y \in E\}$.

- (b) Give a bijection from $D \times E$ to $E \times D$.

Let $f : D \times E \rightarrow E \times D$ be the function that sends (x, y) to (y, x) .

[Explanation: Any element of $D \times E$ can be written as (x, y) with $x \in D$ and $y \in E$. But then the pair (y, x) will be an element of $E \times D$. Clearly this is a bijection.

Notation: If we write $f : (x, y) \mapsto (y, x)$ then this indicates that (x, y) is an element of the domain of f and that (y, x) is an element of the codomain (range) of f . In other words, this notation means that f sends this element (x, y) in $D \times E$ to the element (y, x) in $E \times D$.]

- (c) Why does this bijection prove $d \cdot e = e \cdot d$?

Because $d \cdot e$ is by definition $o(D \times E)$. But giving a bijection from $D \times E$ to $E \times D$ proves that $o(D \times E) = o(E \times D)$ but the latter is by definition equal to $e \cdot d$.

7. TURN IN: Find all sets A for which the following is true: Every element of A is equal to 1.

Answer: $A = \emptyset$ and $A = \{1\}$.

8. TURN IN:

Item 22 says that if d, e are cardinals, and if at least one of them is infinite, then $d + e = \max(d, e)$. It is quite hard to prove this in general. Lets prove it in a special case, when $d = e = \aleph_0$, as follows:

Let $\mathbb{N}^* = \{1, 2, 3, 4, \dots\}$, $E = \{2, 4, 6, 8, \dots\}$, $D = \{1, 3, 5, 7, \dots\}$.

So $E = \{\text{all even positive integers}\}$, and $D = \{\text{all odd positive integers}\}$.

- (a) Give a bijection $f : \mathbb{N}^* \rightarrow E$ (write down: $f(n) = \dots$)

This function: $f(n) = 2n$ is a bijection (there are other correct answers, but this one is the most obvious one).

(b) Give a bijection $g : \mathbb{N}^* \rightarrow D$.

This function: $g(n) = 2n - 1$ is a bijection.

(c) Explain why parts (a),(b) prove that $\aleph_0 + \aleph_0 = \aleph_0$.

If D, E are disjoint sets, each with cardinality \aleph_0 , then $\aleph_0 + \aleph_0$ is by definition the cardinality of the union $D \cup E$. So $\aleph_0 + \aleph_0 = o(D \cup E) = o(\mathbb{N}^*) = \aleph_0$.

[Note: this is exactly like the second part of Hotel Infinity.]

List of facts on cardinal numbers, shortened version.

Note: During the actual test, basic definitions that everyone must know (such as items 1–7) may be deleted!

1. $o(A) = o(B)$ means $\exists f : A \rightarrow B$ with f bijection.
2. $o(A) \leq o(B)$ means $\exists f : A \rightarrow B$ with f one-to-one.
3. \aleph_0 is short notation for $o(\mathbb{N}^*)$.
4. c is short notation for $o(\mathbb{R})$.
5. The set A is *countably infinite* when: $o(A) = \aleph_0$.
By item 1 this means: $\exists f : \mathbb{N}^* \rightarrow A$ with f bijection. Note, in that case $A = f(\mathbb{N}^*) = f(\{1, 2, \dots\}) = \{f(1), f(2), \dots\}$ and this means that all elements of A fit into one sequence $f(1), f(2), \dots$
6. Notation: $x < y$ is short for: $x \leq y \wedge x \neq y$.
7. $o(A) < o(P(A))$.
8. Item 7 implies that not all infinite sets have the same cardinality!
The cardinal number $o(\mathbb{N}^*) = \aleph_0$, is NOT the largest possible cardinality despite the fact that it is infinite! After all, $P(\mathbb{N}^*)$ has larger cardinality by item 7. And $P(P(\mathbb{N}^*))$ has larger cardinality still!
9. If $f : A \rightarrow B$ is onto then $o(B) \leq o(A)$.
10. A is *countable* when either: A is countably infinite (defined in item 5) or A is finite.
11. A is countable when $o(A) \leq \aleph_0$.
12. A subset of a countable set is again countable.
13. If $A \subseteq B$ then $o(A) \leq o(B)$.
14. The ordering \leq on cardinal numbers is a *partial ordering*.
In particular: whenever $d \leq e$ and $e \leq d$ we may conclude $d = e$.
You might remember that the proof was not easy!

15. The ordering \leq on cardinal numbers is a *total ordering*. So given any two cardinals d, e we have $d \leq e$ or $d \geq e$. This means that one of these things must be true: $d < e$ or $d = e$ or $d > e$.
16. Set A is uncountable when $o(A) \not\leq \aleph_0$. Using item 15 we can reformulate this by saying: A is uncountable when $o(A) > \aleph_0$.
17. Any infinite set contains a countably infinite subset. (note: That an uncountable set has a countably infinite subset follows from item 16).
18. \mathbb{Z} and \mathbb{Q} are countable.
19. If you have countably many sets, and if each of these sets is countable, then their union is also countable.
20. \mathbb{R} is uncountable. $c = o(\mathbb{R}) = o(P(\mathbb{N}^*))$.
21. If $d = o(D)$ and $e = o(E)$ then $d + e$ is the cardinality of $D \cup E$ if we assume that $D \cap E = \emptyset$. Likewise, $d \cdot e$ is the cardinality of $D \times E$. d^e is the cardinality of D^E where $D^E = \{\text{all functions from } E \text{ to } D\}$.
22. If d, e are cardinal numbers, and if at least one of them is infinite, then $d + e = \max(d, e)$.
If $d \neq 0$ and $e \neq 0$ and at least one of them is infinite, then $d \cdot e$ equals $\max(d, e)$ as well. So for non-zero cardinals with at least one infinite, the operations $+, \cdot, \max$ are the same!
23. There is a bijection between $P(A)$ and $\{0, 1\}^A$, and hence $o(P(A)) = o(\{0, 1\}^A) = o(\{0, 1\})^{o(A)} = 2^{o(A)}$.
24. $c = o(\mathbb{R}) = o(P(\mathbb{N}^*)) = o(\{0, 1\}^{\mathbb{N}^*}) = 2^{o(\mathbb{N}^*)} = 2^{\aleph_0}$.
25. $(d_1 d_2)^e = d_1^e d_2^e$, $d^{e_1 + e_2} = d^{e_1} d^{e_2}$, $(d^e)^f = d^{ef}$
26. If you have d sets, and each of these sets has cardinality e , and if A is the union of all those sets, then $o(A) \leq de$ (if the d sets are disjoint, then you may replace the \leq by $=$). Now if d or e is infinite, and both are non-zero, then we can also replace de by $\max(d, e)$, see item 22.
27. So far we have encountered these increasing cardinals:

$$0, 1, 2, 3, \dots, \aleph_0, \quad c = 2^{\aleph_0}, \quad 2^c, \quad 2^{2^c}, \dots$$

and we can wonder if there are any cardinals in between. Specifically, the *continuum hypothesis* asks if there is a cardinal d with $\aleph_0 < d < c$.

From the axioms of set theory (= the only statements mathematicians accept without a proof) it is impossible to prove or disprove this.