

MLC Module 1 Section 8 Exercises

- Given a double decrement table with $q_x^{(1)} = .04$ and $p_x^{(\tau)} = .94$, determine $q_x^{(2)}$.
- For a double decrement table, given $p_x'^{(1)} = 0.95$ and $q_x'^{(2)} = 0.1$, determine $q_x^{(\tau)}$.
- For a double decrement table, given $p_x'^{(1)} = 0.9$, $q_x^{(1)} = 0.09$, and $p_x^{(\tau)} = 0.72$, determine
 - $q_x^{(2)}$
 - $q_x'^{(2)}$
- You are given the double decrement table:

x	$l_x^{(\tau)}$	$q_x^{(1)}$	$q_x^{(2)}$	$q_x'^{(1)}$	$q_x'^{(2)}$
30	1000	0.09			0.20
31	712		0.20	0.15	0.20
32		0.15		0.16	
33	305.0208				

Determine

- ${}_2q_{31}^{(2)}$
 - ${}_{1|2}q_{30}^{(1)}$
- For a triple decrement table, given $\mu_x^{(1)}(t) = .1$, $\mu_x^{(2)}(t) = .2$, and $\mu_x^{(3)}(t) = .3$. determine
 - ${}_5q_x^{(\tau)}$
 - ${}_{5|10}q_x^{(2)}$
 - For a triple decrement table, given $\mu_x^{(1)}(t) = .01$, $\mu_x^{(2)}(t) = .02$, and $\mu_x^{(3)}(t) = .03$. determine
 - ${}_{10}q_x^{(\tau)}$
 - ${}_{10}q_x^{(2)}$
 - ${}_{10|10}q_x^{(\tau)}$
 - ${}_{10|10}q_x^{(1)}$
 - the expected time until departure, $e_x^{\circ(\tau)}$

7. For a double decrement table given: $\mu_x^{(1)}(t) = .01 + .01t$, $\mu_x^{(2)}(t) = .02 + .02t$, and $\mu_x^{(3)}(t) = .03 + .03t$, determine

- (a) ${}_5q_x^{(\tau)}$
 (b) ${}_5q_x^{(3)}$
 (c) the conditional probability that departure was by decrement 2, given that departure occurred at age $x+5$
 (d) the conditional probability that departure was by decrement 1, given that departure occurred before age $x+5$

8. You are given the double decrement table:

x	$l_x^{(\tau)}$	$d_x^{(1)}$	$d_x^{(2)}$
50		75	
51	900		
52			25
53			

You are also given:

- (i) $q_{50}^{(\tau)} = .1$
 (ii) ${}_2p_{50}^{(\tau)} = .825$
 (iii) there are twice as many departures from decrement 1 at age 51 as there are from decrement 2 at age 51
 (iv) ${}_2|q_{50}^{(1)} = .025$

Determine

- (a) $q_{50}^{(2)}$
 (b) ${}_2q_{51}^{(\tau)}$
 (c) ${}_1|q_{51}^{(1)}$
 (d) ${}_1|_2q_{50}^{(2)}$
9. Given $l_x^{(\tau)} = 1000$ and a triple decrement table with $\mu_x^{(j)} = 0.1 + 0.2(j - 1)$ for $j = 1, 2$, and 3, determine the expected number of departures between ages x and $x+1$ by decrement 2.

10. Given a double decrement model with $\mu_x^{(1)} = 0.02$ and $\mu_x^{(2)} = 0.03$, determine

- (a) ${}_2p_x^{(\tau)}$
- (b) ${}_2q_x^{(1)}$
- (c) ${}_2q_x^{(2)}$
- (d) ${}_2q_x'^{(1)}$
- (e) ${}_2q_x'^{(2)}$

11. Given a double decrement table where decrement 1 is DML(80) in the associated single decrement table and decrement 2 has $\mu_x^{(2)} = 0.1$, determine

- (a) ${}_{10}q_{50}^{(1)}$
- (b) $\mu_{50}^{(\tau)}(10)$

12. Given a double decrement table where decrement 1 is DML(100) in the associated single decrement table and decrement 2 has $\mu_x^{(2)} = 0.05$, determine

- (a) ${}_{10}q_{30}^{(1)}$
- (b) ${}_{10}q_{30}^{(2)}$

13. Given a double decrement model with $p_x'^{(1)} = 0.9$ and $p_x'^{(2)} = 0.8$, determine $q_x^{(1)}$ and $q_x^{(2)}$ using

- (a) MUDD
- (b) SUDD

14. Given a double decrement model with $q_x^{(1)} = 0.1$ and $q_x^{(2)} = 0.2$, determine $q_x'^{(1)}$ and $q_x'^{(2)}$ using

- (a) MUDD
- (b) SUDD

15. For a double decrement table where each decrement is UDD in the double decrement table, given $q_x'^{(1)} = 0.1$ and $q_x'^{(2)} = 0.2$, determine

- (a) $0.3q_x^{(2)}$
- (b) $0.5|0.3q_x^{(2)}$
- (c) $0.3q_{x+0.5}^{(2)}$

16. Given a double decrement model with $p'_{40}{}^{(1)} = p'_{41}{}^{(1)} = 0.9$ and $p'_{40}{}^{(2)} = p'_{41}{}^{(2)} = 0.8$, determine ${}_{1.5}q_{40}^{(1)}$ using the SUDD assumption.
17. Given a double decrement table where decrement 1 is BOY and decrement 2 is UDD in the associated single decrement table, and given $q_x^{(1)} = 0.1$ and $q_x^{(2)} = 0.2$, determine
- $q_x^{(1)}$
 - $q_x^{(2)}$
18. For a triple decrement table where decrement 1 and decrement 2 are each UDD in their associated single decrement tables, and decrement 3 is EOY, given $q_x^{(j)} = 0.2j$ for $j = 1, 2$, and 3 , determine
- $q_x^{(1)}$
 - $q_x^{(2)}$
 - $q_x^{(3)}$
19. For a double decrement table where decrement 1 is MOY and decrement 2 is UDD in the associated single decrement table, given $q_x^{(1)} = 0.1$ and $q_x^{(2)} = 0.3$ determine
- $q_x^{(1)}$
 - $q_x^{(2)}$
20. For a double decrement table where decrement 1 is SUDD and 25% of decrement 2 occurs at time 0.3 with the rest occurring at time 0.7, given $q_x^{(1)} = 0.2$ and $q_x^{(2)} = 0.4$ determine
- $q_x^{(1)}$
 - $q_x^{(2)}$