

PROPERTY R_∞ FOR SOME SPHERICAL AND AFFINE ARTIN–TITS GROUPS

MATTHIEU CALVEZ AND IGNAT SOROKO

ABSTRACT. Let $n \geq 2$. In this note we give a short uniform proof of property R_∞ for the Artin–Tits groups of spherical types $A_n, B_n, D_4, I_2(m)$ ($m \geq 3$), their pure subgroups, and for the Artin–Tits groups of affine types \tilde{A}_{n-1} and \tilde{C}_n . In particular, we provide an alternative proof of a recent result of Dekimpe, Gonçalves and Ocampo, who established property R_∞ for pure Artin braid groups.

1. INTRODUCTION

Let G be a group and φ an automorphism of G . An equivalence relation on G is defined by saying that $g, h \in G$ are φ -twisted conjugate if and only if there exists $x \in G$ such that $h = xg\varphi(x)^{-1}$. The number of its equivalence classes is the *Reidemeister number* of φ , denoted by $R(\varphi)$. We say that G has property R_∞ if $R(\varphi) = \infty$ for all $\varphi \in \text{Aut}(G)$.

The notion of twisted conjugacy classes arises in Nielsen fixed point theory, where under certain natural conditions the Reidemeister number serves as an upper bound for a homotopy invariant called the Nielsen number. Also the twisted conjugacy classes appear naturally in Selberg theory and in some topics of algebraic geometry. See the introduction sections in [TW11], [Rom11] and references therein.

Property R_∞ has been proved for a number of groups (see [FT15] or [FN16] for a detailed list) among which are Artin braid groups [FGD10], some classes of large Artin–Tits groups [Juh10], and, very recently, pure Artin braid groups [DGO21] and some right-angled Artin–Tits groups [DS21].

Recall that a *Coxeter matrix* over a finite set S is a symmetric matrix $(m_{st})_{s,t \in S}$ with entries in $\{1, 2, \dots, \infty\}$, such that $m_{ss} = 1$ for all $s \in S$ and $m_{st} = m_{ts} \geq 2$ if $s \neq t$. A Coxeter matrix can be encoded by the corresponding *Coxeter graph* X having S as the set of vertices. Two distinct vertices $s, t \in S$ are connected with an edge in X if $m_{st} \geq 3$, and this edge is labeled with m_{st} if $m_{st} \geq 4$. The *Artin–Tits group of type X* is the group $A(X)$ given by the presentation:

$$A(X) = \langle S \mid \text{prod}(s, t, m_{st}) = \text{prod}(t, s, m_{ts}), \text{ for all } s \neq t, m_{st} \neq \infty \rangle,$$

where $\text{prod}(s, t, m_{st})$ is the word $stst\dots$ of length $m_{st} \geq 2$. The *Coxeter group $W(X)$ of type X* is the quotient of $A(X)$ by all relations of the form $s^2 = 1$, $s \in S$. The most famous example is the Artin braid group on n strands, or Artin–Tits group $A(A_{n-1})$, whose corresponding Coxeter group is the symmetric group on n elements.

The groups $A(X)$ and $W(X)$ said to be *irreducible* if X is connected. An Artin–Tits group $A(X)$ is of *spherical type* if the corresponding Coxeter group $W(X)$ is finite, and of *affine type* if $W(X)$ acts geometrically (i.e. properly discontinuously and cocompactly by isometries) on a euclidean space.

We denote by $P(X)$ the kernel of the natural epimorphism $A(X) \rightarrow W(X)$. It is called the *pure Artin–Tits group of type X* . If X is of spherical type, the group $P(X)$ has finite index in $A(X)$. The group $P(A_{n-1})$ is also called the *pure Artin braid group on n strands*.

Date: June 19, 2022.

2020 Mathematics Subject Classification. 20F36, 20F65, 57K20, 20E36, 20E45, 57M07 (primary).

Key words and phrases. Artin–Tits groups, twisted conjugacy, spherical type, affine type.




FIGURE 1. The defining Coxeter graphs for the Artin–Tits groups under consideration.

In this note we focus on irreducible Artin–Tits groups associated to the Coxeter graphs depicted in Figure 1 and on the pure Artin–Tits groups corresponding to those of them which are of spherical type. For a group G , we denote by $\overline{G} = G/Z(G)$ its quotient by the center. We prove the following theorem.

Theorem 1. *Let $n \geq 2$.*

- (1) *Let $X \in \{A_n, B_n, D_4, I_2(m) (m \geq 3, m \neq \infty)\}$. Then the Artin–Tits group $A(X)$, the pure Artin–Tits group $P(X)$, and their central quotients $\overline{A(X)}$ and $\overline{P(X)}$ have property R_∞ .*
- (2) *Let $X \in \{\tilde{A}_{n-1}, \tilde{C}_n\}$. Then the Artin–Tits group $A(X)$ has property R_∞ .*

As is well known (see for example [Rom11]), free abelian groups do not have property R_∞ ; therefore $A(A_1) \simeq P(A_1) \simeq \mathbb{Z}$ and $A(I_2(2)) \simeq P(I_2(2)) \simeq \mathbb{Z}^2$ do not have property R_∞ .

Our proof is uniform and is an application of the strategy implemented in [FGD10] for $A(A_n)$ to all other groups of Theorem 1 (except $A(I_2(m)), P(I_2(m))$, and $A(\tilde{A}_1)$, which are virtually free). Let G be one of these groups and $\Gamma = G/Z(G)$. We use the fact that the groups Γ and $\text{Aut}(\Gamma)$ are finite index subgroups of the extended mapping class group of a suitable punctured surface to deduce that the “mapping torus” of any automorphism φ of Γ acts non-elementarily on the curve complex of that surface. This, combined with a result of Delzant (Lemma 5) and Proposition 4, allows us to distinguish infinitely many twisted conjugacy classes of φ .

In particular, we obtain a short independent proof of a recent theorem of Dekimpe–Gonçalves–Ocampo [DGO21] who established property R_∞ for the pure Artin braid groups $P(A_n)$. For all other groups in Theorem 1 (except $A(A_n)$ and $P(A_n)$) our result is new, to the best of our knowledge.

Acknowledgments. The authors are grateful to Karel Dekimpe for his encouragement to write this note and to Peter Wong for a useful discussion. The authors would also like to thank the anonymous referee for his or her valuable suggestions which improved the quality of the paper. The first author was supported by the EPSRC New Investigator Award EP/S010963/1 and FONDECYT 1180335. The second author acknowledges support from the AMS–Simons Travel Grant.

2. PRELIMINARY RESULTS

In this section we collect preliminary results which we will use to prove Theorem 1 in Section 3.

Recall that the center of an irreducible Artin–Tits group $A(X)$ (and its corresponding pure Artin–Tits group $P(X)$) of spherical type is cyclic, see [BS72, Satz 7.2] (and [CP22, Corollary 7], respectively). On the other hand, the center of an irreducible Artin–Tits group of affine type is trivial, see [McCS20, Proposition 11.9]. It is known (and is easy to prove) that the central quotient $\overline{A(X)}$ has trivial center itself, which is also true for $\overline{P(X)}$ (see [CP22, Corollary 7]).

The following Proposition allows us to reduce the proof of Theorem 1 about $A(X)$ and $P(X)$ to proving a similar statement for their central quotients $\overline{A(X)}$ and $\overline{P(X)}$, respectively.

Proposition 2 (See e.g. [FGD10, (2.2)]). *Let G be a group, φ be an automorphism of G , and H be a normal φ -invariant subgroup of G . Denote by $\bar{\varphi}$ the automorphism induced by φ on G/H . Then the condition $R(\bar{\varphi}) = \infty$ implies $R(\varphi) = \infty$. In particular, if $G/Z(G)$ has property R_∞ , then so does G . \square*

Recall that for a group G , $\text{Inn}(G)$ denotes the subgroup of $\text{Aut}(G)$ consisting of all inner automorphisms. For $g \in G$, we denote by ι_g the inner automorphism $x \mapsto gxg^{-1}$. The map $\iota : g \mapsto \iota_g$ identifies $\text{Inn}(G)$ with $G/Z(G)$ and $\text{Inn}(G)$ is a normal subgroup of $\text{Aut}(G)$. The *outer automorphism group* of G is the quotient $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$.

Now, let us review an interesting relationship between the twisted conjugacy classes of automorphisms of a centerless group G and the actual conjugacy classes in an extension of G . Let φ be an automorphism of a centerless group G , and let $m \in \{1, 2, 3, \dots, \infty\}$ be the order of φ in $\text{Out}(G)$; if $m < \infty$, let $p \in G$ be the unique element such that $\varphi^m = \iota_p$. Consider the group

$$G_\varphi = \begin{cases} G * \langle t \rangle / \langle tgt^{-1} = \varphi(g) \text{ for all } g \in G \rangle, & \text{if } m = \infty, \\ G * \langle t \rangle / \langle tgt^{-1} = \varphi(g) \text{ for all } g \in G, t^m = p \rangle, & \text{if } m < \infty. \end{cases}$$

Note that if $m = \infty$, G_φ is the semidirect product $G \rtimes_\varphi \mathbb{Z}$, and if $m = 1$, G_φ is isomorphic to G .

Lemma 3. *Let G be a centerless group, $\varphi \in \text{Aut}(G)$, and let G_φ be defined as above. Then G is a normal subgroup of G_φ , the quotient G/G_φ is cyclic, and the assignments $g \mapsto \iota_g$ and $t \mapsto \varphi$ define an injective homomorphism from G_φ to $\text{Aut}(G)$.*

Proof. We prove the second statement first. Indeed, the given assignments $g \mapsto \iota_g$ and $t \mapsto \varphi$ define a homomorphism since relations in G_φ hold in $\text{Aut}(G)$: the conjugation by φ in $\text{Aut}(G)$ sends the inner automorphism ι_g for $g \in G$ to $\varphi \iota_g \varphi^{-1} = \iota_{\varphi(g)}$. Every element of G_φ can be written as $t^k g$, for some $g \in G$ and $k \in \mathbb{Z}$ if $m = \infty$ and $k \in \{0, \dots, m-1\}$ if $m < \infty$. Then the element $t^k g$ is sent to $\varphi^k \iota_g$, and if $\varphi^k \iota_g = \text{id}_G$, we deduce first $\varphi^k = \iota_g^{-1} = \iota_{g^{-1}}$, which forces $k = 0$ by definition of m . Then since G is assumed to be centerless, $\iota_g = \text{id}_G$ implies that $g = 1$ and hence the homomorphism in question is injective.

If $m = \infty$, then G_φ is the semidirect product $G \rtimes_\varphi \langle t \rangle$, hence G is a subgroup of G_φ . If $m < \infty$, we note that the injective homomorphism $\iota : G \rightarrow \text{Aut}(G)$, $g \mapsto \iota_g$, factors through the composition $G \rightarrow G_\varphi \rightarrow \text{Aut}(G)$, and hence the map $G \rightarrow G_\varphi$ is itself injective. In either case, G is a normal subgroup and G_φ/G is generated by the coset tG . \square

Proposition 4 (cf. [FGD10, Lemma 6.2]). *Let G be a centerless group, $\varphi \in \text{Aut}(G)$, and let G_φ be defined as above. Two elements $g, h \in G$ are φ -twisted conjugate if and only if the elements gt and ht of G_φ are conjugate in G_φ . In particular, $R(\varphi) = \infty$ if and only if the coset Gt in G_φ contains infinitely many conjugacy classes.*

Proof. Suppose that $g, h \in G$ are φ -twisted conjugate. Then there exists some $x \in G$ such that $h = xg\varphi(x)^{-1}$. In G_φ , we then have $h = xgtx^{-1}t^{-1}$, which is to say that $ht = x(gt)x^{-1}$. Conversely, suppose that gt and ht are conjugate in G_φ . Then there is some element $t^k x$, with $x \in G$ and $k \in \mathbb{Z}$ if $m = \infty$, and $k \in \{0, \dots, m-1\}$ if $m < \infty$, such that $ht = t^k x \cdot gt \cdot x^{-1} t^{-k}$, or $ht^{k+1} x = t^k xgt$. Using the relation $t^k \gamma = \varphi^k(\gamma)t^k$ for $\gamma \in G$, we obtain $h\varphi^{k+1}(x)t^{k+1} = \varphi^k(xg)t^{k+1}$. Canceling t^{k+1} on the right and setting $y = \varphi^k(x)$, we obtain $h\varphi(y) = y\varphi^k(g)$, or $h = y\varphi^k(g)\varphi(y)^{-1}$. This is to say that h and $\varphi^k(g)$ are φ -twisted conjugate. To conclude, it suffices to know that $\varphi^k(g)$ is φ -twisted conjugate to g : in general, for every automorphism ψ , any $\gamma \in G$ is ψ -twisted conjugate to its image $\psi(\gamma)$ according to the equality $\gamma = \gamma\psi(\gamma)\psi(\gamma)^{-1}$. \square

In particular, we recover from Proposition 4 the well-known fact that for any inner automorphism φ , the number of φ -twisted conjugacy classes $R(\varphi)$ is equal to the number of conjugacy classes in G (see e.g. [Tro19, Lemma 1.3] for $\varphi = \text{id}$).

Let G be a group acting on a Gromov hyperbolic space S . For any $s \in S$ consider the set $\Lambda(G)$ of accumulation points of the orbit Gs on the boundary ∂S . The action of G on S is said to be *non-elementary*, if $\Lambda(G)$ contains at least three points, see [Osi16, Section 3].

The main tool for our proof is the following result due to Delzant:

Lemma 5 ([FGD10, Lemma 6.3],[LL00, Lemma 3.4]). *Let Γ be a group acting non-elementarily on a Gromov hyperbolic space, and let K be a normal subgroup of Γ such that the quotient Γ/K is abelian. Then any coset of K contains infinitely many conjugacy classes.* \square

This lemma will be used as follows. Let G be a centerless group, $\varphi \in \text{Aut}(G)$ and G_φ as above, then if G_φ acts non-elementarily on a Gromov hyperbolic space, Lemma 5 (with $\Gamma = G_\varphi$ and $K = G$) and Proposition 4 show that $R(\varphi) = \infty$. We will also make use of the following result of Fel'shtyn:

Theorem 6 ([Fel04, Theorem 3]). *Every non-elementary hyperbolic group has property R_∞ .* \square

3. PROOF OF THEOREM 1

Proposition 7. *Let $n \geq 3$ and let $X \in \{A_n, B_n, D_4, \tilde{A}_{n-1}, \tilde{C}_{n-1}\}$. Let G be either the Artin–Tits group $A(X)$, or the pure Artin–Tits group $P(A_n)$, $P(B_n)$, $P(D_4)$, and denote $\Gamma = G/Z(G)$. Then there exists a surface Σ_Γ with punctures such that both Γ and $\text{Aut}(\Gamma)$ can be embedded as nested subgroups of finite index in the extended mapping class group of Σ_Γ :*

$$\Gamma \leq \text{Aut}(\Gamma) \leq \text{Mod}^\pm(\Sigma_\Gamma).$$

Here Γ is identified with the subgroup $\text{Inn}(\Gamma)$ of inner automorphisms. If g and p denote the genus and the number of punctures of Σ_Γ , respectively, then they satisfy the inequality: $3g + p - 4 > 0$.

Proof. It is shown in [CC05, Section 2] that for $G = A(X)$ with $X \in \{A_n, B_n, \tilde{A}_{n-1}, \tilde{C}_{n-1}\}$, the group $\Gamma = G/Z(G)$ is a subgroup of finite index in the extended mapping class group of a sphere with $n + 2$ punctures $\Sigma_\Gamma = S_{n+2}$. Namely, $\overline{A(A_n)}$, $\overline{A(B_n)}$, and $\overline{A(\tilde{C}_{n-1})}$ are the subgroups of all orientation preserving mapping classes of S_{n+2} that fix 1, 2, and 3 punctures of S_{n+2} , respectively. Similarly, $\overline{A(\tilde{A}_{n-1})}$ is a subgroup of index n in $\overline{A(B_n)}$. For $G = P(A_n)$, it is briefly mentioned in [CC05, p. 324] and discussed in detail in [FM12, Section 9.3] that $\Gamma = G/Z(G)$ is the group of all orientation preserving mapping classes of S_{n+2} that fix all punctures pointwise, i.e. it is the pure orientation preserving mapping class group of the sphere with $n + 2$ punctures, which has index $2(n + 2)!$ in $\text{Mod}^\pm(S_{n+2})$. In general, if $G = P(X)$ is the pure Artin–Tits group for a connected Coxeter graph X of spherical type, it was proven in [CP22, Corollary 7] that $G/Z(G)$ is isomorphic to a subgroup of finite index in $\overline{A(X)}$. In particular, for $G = P(B_n)$, the group $\Gamma = G/Z(G)$ is isomorphic to a subgroup of finite index in $\overline{A(B_n)}$, and hence to a subgroup of finite index in $\text{Mod}^\pm(S_{n+2})$. Note that for $\Sigma_\Gamma = S_{n+2}$ we have $g = 0$, $p = n + 2$, so the required inequality becomes: $3g + p - 4 = n - 2 > 0$, which is true if $n \geq 3$.

Now we invoke the Corollary to the Ivanov–Korkmaz theorem [CC05, Corollary 4], and conclude that in all the above cases the group $\text{Aut}(\Gamma)$ is isomorphic to the normalizer of Γ in $\text{Mod}^\pm(S_{n+2})$. Since $\text{Aut}(\Gamma)$ contains Γ as a subgroup (identified with the subgroup $\text{Inn}(\Gamma)$ of all inner automorphisms), and Γ has finite index in $\text{Mod}^\pm(S_{n+2})$, we conclude that $\text{Aut}(\Gamma)$ has itself finite index in $\text{Mod}^\pm(S_{n+2})$.

If $G = A(D_4)$, let $\Gamma = \overline{A(D_4)}$ and let Σ_Γ be the torus with three punctures. It was shown in [Sor20, Theorem 1] that Γ is isomorphic to the pure orientation preserving mapping class group of Σ_Γ and in [Sor21, Corollary 6] that $\text{Aut}(\Gamma)$ is isomorphic to the extended mapping class group $\text{Mod}^\pm(\Sigma_\Gamma)$, which contains Γ as a subgroup of index 12. For $G = P(D_4)$, we refer to [CP22, Corollary 7] again to conclude that $\Gamma = G/Z(G)$ is isomorphic to a subgroup of finite index in $\overline{A(D_4)}$, and hence to a subgroup of finite index in $\text{Mod}^\pm(\Sigma_\Gamma)$, with Σ_Γ being the torus with three punctures. Clearly, the required inequality is satisfied for Σ_Γ in this case as well: $3g + p - 4 = 3 + 3 - 4 > 0$. \square

Now we are ready to prove Theorem 1.

Proof of Theorem 1. Let G be any of the groups from the theorem; we are going to prove that its central quotient $G/Z(G)$ has property R_∞ . Then Proposition 2 will yield the desired conclusion for G .

First, let $X = I_2(m)$, $m \geq 3$, $m \neq \infty$. It is known that $\overline{A(X)}$ is isomorphic to $C_m \star C_2$ if m is odd, and to $C_{m/2} \star \mathbb{Z}$ if m is even, where C_n denotes the cyclic group of order n , see e.g. [CP02, Section 5]. In any case, $\overline{A(X)}$ contains a finite index nonabelian free subgroup and hence is non-elementary hyperbolic. By [CP22, Corollary 7], the central quotient $\overline{P(X)}$ is isomorphic to a subgroup of finite index in $\overline{A(X)}$, hence also contains a nonabelian free subgroup of finite index, and thus is also non-elementary hyperbolic. (One can actually prove that $\overline{P(I_2(m))} \cong F_{m-1}$, the free group of rank $m-1$, see [CP22, Proof of Lemma 6.2].) Theorem 6 now shows that $\overline{A(X)}$ and $\overline{P(X)}$ have property R_∞ , and hence, by Proposition 2, $A(X)$ and $P(X)$ have property R_∞ as well.

In view of the isomorphisms $A(A_2) \simeq A(I_2(3))$ and $A(B_2) \simeq A(I_2(4))$, the above reasoning covers the groups $A(A_2)$, $A(B_2)$, $P(A_2)$, $P(B_2)$; the result for $A(\tilde{A}_1) \simeq F_2$ also follows from Theorem 6.

Suppose now that $n \geq 3$ and $G = A(X)$ for $X \in \{A_n, B_n, D_4, \tilde{A}_{n-1}, \tilde{C}_{n-1}\}$, or $G = P(A_n)$, $P(B_n)$, or $P(D_4)$. Denote $\Gamma = G/Z(G)$ (recall that Γ is isomorphic to G for $G = A(\tilde{A}_{n-1})$ or $A(\tilde{C}_{n-1})$). Let $\varphi \in \text{Aut}(\Gamma)$ and consider the group Γ_φ as defined in Section 2. Then by Lemma 3 and Proposition 7, there exist a surface with punctures Σ_Γ and a tower of inclusions of finite index subgroups of the extended mapping class group of Σ_Γ ,

$$\Gamma \leq \Gamma_\varphi \leq \text{Aut}(\Gamma) \leq \text{Mod}^\pm(\Sigma_\Gamma).$$

Now recall that $\text{Mod}^\pm(\Sigma_\Gamma)$, and hence in particular its subgroup Γ_φ , acts by isometries on the curve complex $\mathcal{C}(\Sigma_\Gamma)$ of the surface Σ_Γ . This curve complex is Gromov hyperbolic by the result of Masur–Minsky [MM99, Theorem 1.1] when $3g + p - 4 > 0$, where g denotes the genus of the surface Σ_Γ and p is the number of punctures. We claim that the action of Γ_φ on $\mathcal{C}(\Sigma_\Gamma)$ is non-elementary. Indeed, it is known (see for instance [Bow08] and [Osi16, Theorem 1.1]) that $\text{Mod}^\pm(\Sigma_\Gamma)$ contains infinitely many independent elements acting loxodromically on $\mathcal{C}(\Sigma_\Gamma)$. (Recall that an element acts loxodromically if its orbit has two accumulation points on the boundary and two such elements are independent if the sets of accumulation points of their orbits are disjoint.) Note that if an element acts loxodromically, so do all of its powers. Since Γ_φ has finite index in $\text{Mod}^\pm(\Sigma_\Gamma)$, this implies that Γ_φ itself contains infinitely many independent elements acting loxodromically on $\mathcal{C}(\Sigma_\Gamma)$. Therefore, the action of Γ_φ on $\mathcal{C}(\Sigma_\Gamma)$ is non-elementary and, by Proposition 4 and Lemma 5, $R(\varphi) = \infty$. As φ was arbitrary, this implies that Γ has property R_∞ , and from Proposition 2 we deduce that G has property R_∞ . This finishes the proof of Theorem 1. \square

Remark 8. Given an Artin–Tits group $A(X)$, there is another (in most cases, conjecturally) hyperbolic space with an action of the central quotient $\overline{A(X)}$. This is the *graph of irreducible parabolic subgroups* $\mathcal{C}_{\text{parab}}(A(X))$ introduced in [CGGW19] and also studied in [MW21, CC21]. Hyperbolicity of $\mathcal{C}_{\text{parab}}(A(X))$ was established for X being A_n ($n \geq 3$), B_n ($n \geq 3$), \tilde{A}_n ($n \geq 2$), and \tilde{C}_n ($n \geq 2$), see [CC21]. Also, it can be shown using results from [Sor21] that $\mathcal{C}_{\text{parab}}(A(D_4))$ is isomorphic to the curve graph of the three times punctured torus, and hence it is also hyperbolic by Masur–Minsky’s theorem. Given a non-inner automorphism φ of $\overline{A(X)}$, we found that in some cases, the action of $\overline{A(X)}$ extends to an action of $\overline{A(X)}_\varphi$ on $\mathcal{C}_{\text{parab}}(A(X))$; this is for instance the case when φ is a parabolic-preserving automorphism. However, we also found that for some non-parabolic-preserving automorphisms of $A(\tilde{C}_n)$, there is no action of $A(\tilde{C}_n)_\varphi$ on $\mathcal{C}_{\text{parab}}(A(\tilde{C}_n))$, so this approach does not yield a uniform proof of Theorem 1 as the one above.

REFERENCES

- [Bow08] Brian H. Bowditch, Tight geodesics in the curve complex, *Invent. Math.* 171 (2008), no. 2, 281–300.

- [BS72] Egbert Brieskorn, Kyoji Saito, Artin-Gruppen und Coxeter-Gruppen, *Invent. Math.* 17, (1972), 245–271.
- [CC21] Matthieu Calvez, Bruno Cisneros de la Cruz, Curve graphs for Artin–Tits groups of type B , \tilde{A} and \tilde{C} are hyperbolic. *Trans. London Math. Soc.* 8 (1), (2021), 151–173.
- [CC05] Ruth Charney, John Crisp, Automorphism groups of some affine and finite type Artin groups. *Math. Res. Lett.* 12 (2005), no. 2–3, 321–333.
- [CP02] John Crisp, Luis Paris, Artin groups of type B and D. Preprint (2002) arXiv:math/0210438v1.
- [CGGW19] María Cumplido, Volker Gebhardt, Juan González-Meneses and Bert Wiest, On parabolic subgroups of Artin–Tits groups of spherical type, *Adv. Math.* 352 (2019) 572–610.
- [CP22] María Cumplido, Luis Paris, Commensurability in Artin groups of spherical type. *Rev. Mat. Iberoam.* 38 (2022), no. 2, 503–526.
- [DGO21] Karel Dekimpe, Daciberg Lima Gonçalves, Oscar Ocampo, The R_∞ property for pure Artin braid groups. *Monatsh. Math.* 195 (2021), 15–33.
- [DS21] Karel Dekimpe, Peter Senden, The R_∞ -property for right-angled Artin groups. *Topology Appl.* 293 (2021), 107557.
- [FM12] Benson Farb, Dan Margalit, *A Primer on Mapping Class Groups*. Princeton Mathematical Series, 49. Princeton University Press, Princeton, NJ, 2012. xiv+472 pp.
- [Fel04] Alexander Fel’shtyn, The Reidemeister number of any automorphism of a Gromov-hyperbolic group is infinite, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 279 (2001), Geom. i Topol. 6, 229–240, 250; reprinted in *J. Math. Sci. (N.Y.)* 119 (2004), no. 1, 117–123.
- [FGD10] Alexander Fel’shtyn, Daciberg L. Gonçalves, Twisted conjugacy classes in symplectic groups, mapping class groups and braid groups. With an appendix written jointly with Francois Dahmani. *Geom. Dedicata* 146 (2010), 211–223.
- [FN16] Alexander Fel’shtyn, Timur Nasybullov, The R_∞ and S_∞ properties for linear algebraic groups. *J. Group Theory* 19 (2016), 901–921.
- [FT15] Alexander Fel’shtyn, Evgenij Troitsky, Aspects of the property R_∞ , *J. Group Theory* 18 (2015), 1021–1034.
- [Juh10] Arye Juhasz, Twisted conjugacy in certain Artin groups, *Ischia Group Theory 2010*, 175–195.
- [LL00] Gilbert Levitt, Martin Lustig, Most automorphisms of a hyperbolic group have very simple dynamics, *Ann. Sci. Ec. Norm. Sup.*, 33, 4, (2000), 507–517.
- [MM99] Howard Masur, Yair Minsky, Geometry of the complex of curves I: Hyperbolicity, *Invent. Math.* 138 (1999), 103–149.
- [McCS20] Jon McCammond, Robert Sulway, Artin groups of Euclidean type, *Invent. Math.* 210, (2017), 231–282.
- [MW21] Rose Morris-Wright, Parabolic subgroups in FC-type Artin groups, *J. Pure Appl. Algebra* 225 (2021), no. 1, 106468.
- [Osi16] Denis Osin, Acylindrically hyperbolic groups. *Trans. Amer. Math. Soc.* 368 (2016), no. 2, 851–888.
- [Rom11] V. Roman’kov, Twisted conjugacy classes in nilpotent groups, *J. Pure Appl. Algebra* 215 (2011), 664–671.
- [Sor20] Ignat Soroko, Linearity of some low-complexity mapping class groups. *Forum Math.* 32 (2020), no. 2, 279–286.
- [Sor21] Ignat Soroko, Artin groups of types F_4 and H_4 are not commensurable with that of type D_4 . *Topology Appl.* 300 (2021), 107770.
- [TW11] Jennifer Taback, Peter Wong, The geometry of twisted conjugacy classes in wreath products. *Geometry, rigidity, and group actions*, 561–587, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 2011.
- [Tro19] Evgenij Troitsky, Reidemeister classes in some weakly branch groups. *Russ. J. Math. Phys.*, 26 (2019), no. 1, 122–129.

IRISIB, RUE ROYALE 150, 1000 BRUXELLES, BELGIQUE

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD DE VALPARAÍSO, GRAN BRETAÑA 1091, 3ER PISO, PLAYA ANCHA, VALPARAÍSO, CHILE

Email address: calvez.matthieu@gmail.com

DEPARTMENT OF MATHEMATICS, FLORIDA STATE UNIVERSITY, TALLAHASSEE, FL 32306, USA

Email address: soroko@math.fsu.edu, ignat.soroko@gmail.com