

Riemannian proximal gradient methods and variants

Speaker: Wen Huang


Xiamen University

July 13, 2023

Chinese Academy of Sciences

Optimization on Manifolds with Structure:


$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x),$$



- \mathcal{M} is a finite-dimensional Riemannian manifold;
- f is smooth and may be nonconvex; and
- $h(x)$ is continuous and convex but may be nonsmooth;

Optimization on Manifolds with Structure:

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x),$$



- \mathcal{M} is a finite-dimensional Riemannian manifold;
- f is smooth and may be nonconvex; and
- $h(x)$ is continuous and convex but may be nonsmooth;

Applications: sparse PCA [ZHT06], compressed model [OLCO13], sparse partial least squares regression [CSG⁺18], sparse inverse covariance estimation [BESS19], sparse blind deconvolution [ZLK⁺17], and clustering [HWGVD22].

$F : \mathcal{M} \rightarrow \mathbb{R}$ is Lipschitz continuous

- [Huang \(2013\)](#), Gradient sampling method without convergence analysis.
- [Grohs and Hosseini \(2015\)](#), Two ϵ -subgradient-based optimization methods using line search strategy and trust region strategy, respectively. Any limit point is a critical point.
- [Hosseini and Uschmajew \(2017\)](#), Gradient sampling method and any limit point is a critical point.
- [Hosseini, Huang, and Yousefpour \(2018\)](#), Merge ϵ -subgradient-based and quasi-Newton ideas and show any limit point is a critical point.

Existing Nonsmooth Optimization on Manifolds

$F : \mathcal{M} \rightarrow \mathbb{R}$ is convex

- [Zhang and Sra \(2016\)](#), subgradient-based method and function value converges to the optimal $O(1/\sqrt{k})$.
- [Ferreira and Oliveira \(2002\)](#) proximal point method, convergence using convexity
[Bento, da Cruz Neto and Oliveira \(2011\)](#), convergence using Kurdyka-Łojasiewicz (KL); and
[Bento, Ferreira, and Melo \(2017\)](#), function value converges to the optimal $O(1/k)$ on Hadamard manifold using convexity

Existing Nonsmooth Optimization on Manifolds

$F = f + g$, where f is L-con, and g is non-smooth

- [Chen, Ma, So, and Zhang \(2018\)](#), A proximal gradient method with global convergence
- [Xiao, Liu, and Yuan \(2021\)](#), Infeasible approach over the Stiefel manifold
- [Zhou, Bao, and Ding \(2022\)](#), An augmented Lagrangian method on matrix manifolds
- [Huang and Wei \(2021-2023\)](#), A Riemannian proximal gradient method, an inexact Riemannian proximal gradient method, and a modified FISTA on embedded manifolds
- [Wang and Yang \(2023\)](#), A proximal quasi-Newton method on manifolds on the Stiefel manifold
- [Huang, Meng, Gallivan, and Van Dooren \(2023\)](#), An inexact proximal gradient method on embedded submanifolds
- [Beck and Rosset \(2023\)](#), A dynamic smoothing technique

Optimization with Structure:

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

- Proximal gradient methods
- Inexact proximal gradient methods
- A proximal Newton method

[HW2021]: W. Huang and K. Wei, Riemannian proximal gradient methods, Mathematics Programming, 194, 371-413, 2022.

[HW2023]: An inexact Riemannian proximal gradient method, Computational Optimization and Applications, 85, 1-32, 2023

[HWGV2023]: A Riemannian optimization approach to clustering problems, arxiv, 2023

[SAHJV2023]: A Riemannian proximal Newton method, arxiv, 2023

Optimization with Structure:

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

- Proximal gradient methods
 - Euclidean version
 - Riemannian version in [CMSZ20]
 - Riemannian version in [HW21a]
- Inexact proximal gradient methods
- A proximal Newton method

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping}^1\text{)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

1. The update rule: $x_{k+1} = \arg \min_x \langle \nabla f(x_k), x - x_k \rangle + \frac{L}{2} \|x - x_k\|^2 + h(x)$.

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;
- Proximal mapping: easy to compute;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;
- Proximal mapping: easy to compute;
- Any limit point is a critical point;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;
- Proximal mapping: easy to compute;
- Any limit point is a critical point;
- $O\left(\frac{1}{k}\right)$ sublinear convergence rate for convex f and h ;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;
- Proximal mapping: easy to compute;
- Any limit point is a critical point;
- $O\left(\frac{1}{k}\right)$ sublinear convergence rate for convex f and h ;
- Linear convergence rate for strongly convex f and convex h ;

Proximal Gradient Method

Euclidean version

Optimization with Structure: $\mathcal{M} = \mathbb{R}^n$

$$\min_{x \in \mathbb{R}^n} F(x) = f(x) + h(x).$$

initial iterate: x_0 ,

$$\begin{cases} d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p), & \text{(Proximal mapping)} \\ x_{k+1} = x_k + d_k. & \text{(Update iterates)} \end{cases}$$

- $h = 0$: reduce to steepest descent method;
- L : greater than the Lipschitz constant of ∇f ;
- Proximal mapping: easy to compute;
- Any limit point is a critical point;
- $O\left(\frac{1}{k}\right)$ sublinear convergence rate for convex f and h ;
- Linear convergence rate for strongly convex f and convex h ;
- Local convergence rate by KL property;

Proximal Gradient Method

Riemannian versions

Optimization with Structure: \mathcal{M}

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

Optimization with Structure: \mathcal{M}

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

Euclidean proximal mapping

$$d_k = \arg \min_{p \in \mathbb{R}^n} \langle \nabla f(x_k), p \rangle + \frac{L}{2} \|p\|_F^2 + h(x_k + p)$$

In the Riemannian setting:

- How to define the proximal mapping?
- Can be solved cheaply?
- Share the same convergence rate?

Proximal Gradient Method

Riemannian version in [CMSZ20]

A Riemannian proximal mapping [CMSZ20]

$$\textcircled{1} \quad \eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$$

- Only works for embedded submanifold;

[CMSZ18]: S. Chen, S. Ma, M. C. So, and T. Zhang, Proximal gradient method for nonsmooth optimization over the Stiefel manifold. SIAM Journal on Optimization, 30(1):210-239, 2020.

Proximal Gradient Method

Riemannian version in [CMSZ20]

A Riemannian proximal mapping [CMSZ20]

$$\textcircled{1} \quad \eta_k = \arg \min_{\eta \in T_{x_k}} \mathcal{M} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$$

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;

Proximal Gradient Method

Riemannian version in [CMSZ20]

A Riemannian proximal mapping [CMSZ20]

$$\textcircled{1} \quad \eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$$

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;
- **Convex programming**;

Proximal Gradient Method

Riemannian version in [CMSZ20]

[CMSZ20]

$$\textcircled{1} \quad \eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$$

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;
- Convex programming;
- Solved efficiently for the Stiefel manifold by a semi-smooth Newton algorithm [XLWZ18];

[XLWZ18]: X. Xiao, Y. Li, Z. Wen, and L. Zhang, A regularized semi-smooth Newton method with projection steps for composite convex programs. *Journal of Scientific Computing*, 76(1):364-389, 2018.


Proximal Gradient Method

Riemannian version in [CMSZ20]

ManPG [CMSZ20]

- ① $\eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$
- ② $x_{k+1} = R_{x_k}(\alpha_k \eta_k)$ with an appropriate step size α_k ;

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;
- Convex programming;
- Solved efficiently for the Stiefel manifold by a semi-smooth Newton algorithm [XLWZ18];
- Step size 1 is not necessary decreasing;




Proximal Gradient Method

Riemannian version in [CMSZ20]

ManPG [CMSZ20]

- ① $\eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$
- ② $x_{k+1} = R_{x_k}(\alpha_k \eta_k)$ with an appropriate step size α_k ;

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;
- Convex programming;
- Solved efficiently for the Stiefel manifold by a semi-smooth Newton algorithm [XLWZ18];
- Step size 1 is not necessary decreasing;
- **Convergence to a stationary point;**




Proximal Gradient Method

Riemannian version in [CMSZ20]

ManPG [CMSZ20]

- ① $\eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta);$
- ② $x_{k+1} = R_{x_k}(\alpha_k \eta_k)$ with an appropriate step size α_k ;

- Only works for embedded submanifold;
- Proximal mapping is defined in tangent space;
- Convex programming;
- Solved efficiently for the Stiefel manifold by a semi-smooth Newton algorithm [XLWZ18];
- Step size 1 is not necessary decreasing;
- Convergence to a stationary point;
- No convergence rate analysis;



Proximal Gradient Method

Riemannian version in [HW21a]

GOAL: Develop a Riemannian proximal gradient method with convergence rate analysis and good numerical performance for some instances

Proximal Gradient Method

Riemannian version in [HW21a]

GOAL: Develop a Riemannian proximal gradient method with convergence rate analysis and good numerical performance for some instances

A Riemannian Proximal Gradient Method (RPG)

$$\text{Let } \ell_{x_k}(\eta) = \underbrace{\langle \nabla f(x_k), \eta \rangle_{x_k}}_{\text{Riemannian metric}} + \frac{L}{2} \|\eta\|_{x_k}^2 + h(\underbrace{R_{x_k}(\eta)}_{\text{replace } x_k + \eta});$$

- ① $\eta_k \in T_{x_k} \mathcal{M}$ is a stationary point of $\ell_{x_k}(\eta)$, and $\ell_{x_k}(0) \geq \ell_k(\eta_k)$;
- ② $x_{k+1} = R_{x_k}(\eta_k)$;

- General framework for Riemannian optimization;

Proximal Gradient Method

Riemannian version in [HW21a]

GOAL: Develop a Riemannian proximal gradient method with convergence rate analysis and good numerical performance for some instances

A Riemannian Proximal Gradient Method (RPG)

$$\text{Let } \ell_{x_k}(\eta) = \underbrace{\langle \nabla f(x_k), \eta \rangle_{x_k}}_{\text{Riemannian metric}} + \frac{L}{2} \|\eta\|_{x_k}^2 + h(\underbrace{R_{x_k}(\eta)}_{\text{replace } x_k + \eta});$$

- ① $\eta_k \in T_{x_k} \mathcal{M}$ is a stationary point of $\ell_{x_k}(\eta)$, and $\ell_{x_k}(0) \geq \ell_k(\eta_k)$;
- ② $x_{k+1} = R_{x_k}(\eta_k)$;

- General framework for Riemannian optimization;
- Step size can be fixed to be 1;

Proximal Gradient Method

Riemannian version in [HW21a]

GOAL: Develop a Riemannian proximal gradient method with convergence rate analysis and good numerical performance for some instances

A Riemannian Proximal Gradient Method (RPG)

$$\text{Let } \ell_{x_k}(\eta) = \underbrace{\langle \nabla f(x_k), \eta \rangle_{x_k}}_{\text{Riemannian metric}} + \frac{L}{2} \|\eta\|_{x_k}^2 + h(\underbrace{R_{x_k}(\eta)}_{\text{replace } x_k + \eta});$$

- ① $\eta_k \in T_{x_k} \mathcal{M}$ is a stationary point of $\ell_{x_k}(\eta)$, and $\ell_{x_k}(0) \geq \ell_k(\eta_k)$;
- ② $x_{k+1} = R_{x_k}(\eta_k)$;

- General framework for Riemannian optimization;
- Step size can be fixed to be 1;
- Convergence rate results;

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- 1 The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;

This assumption hold if, for example, F is continuous and \mathcal{M} is compact.

$$\min_{X \in \text{St}(p, n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- 1 The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;
- 2 The function f is L -retraction-smooth with respect to the retraction R in the sublevel set Ω_{x_0} .

Definition

A function $h : \mathcal{M} \rightarrow \mathbb{R}$ is called L -retraction-smooth with respect to a retraction R in $\mathcal{N} \subseteq \mathcal{M}$ if for any $x \in \mathcal{N}$ and any $\mathcal{S}_x \subseteq T_x \mathcal{M}$ such that $R_x(\mathcal{S}_x) \subseteq \mathcal{N}$, we have that

$$h(R_x(\eta)) \leq h(x) + \langle \text{grad } h(x), \eta \rangle_x + \frac{L}{2} \|\eta\|_x^2, \quad \forall \eta \in \mathcal{S}_x.$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- ① The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;
 - ② The function f is L -retraction-smooth with respect to the retraction R in the sublevel set Ω_{x_0} .
-

If the following conditions hold, then f is L -retraction-smooth with respect to the retraction R in the manifold \mathcal{M} [BAC18, Lemma 2.7]

- \mathcal{M} is a compact Riemannian submanifold of a Euclidean space \mathbb{R}^n ;
- the retraction R is globally defined;
- $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is L -smooth in the convex hull of \mathcal{M} ;

$$\min_{X \in \text{St}(p, n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- ① The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;
 - ② The function f is L -retraction-smooth with respect to the retraction R in the sublevel set Ω_{x_0} .
-

Theoretical results:

- For any accumulation point x_* of $\{x_k\}$, x_* is a stationary point, i.e., $0 \in \partial F(x_*)$.

Proximal Gradient Method

Riemannian version in [HW21a]

Additional Assumptions:

- f and g are retraction-convex in $\Omega \supseteq \Omega_{x_0}$;
-

Definition

A function $h : \mathcal{M} \rightarrow \mathbb{R}$ is called retraction-convex with respect to a retraction R in $\mathcal{N} \subseteq \mathcal{M}$ if for any $x \in \mathcal{N}$ and any $\mathcal{S}_x \subseteq T_x \mathcal{M}$ such that $R_x(\mathcal{S}_x) \subseteq \mathcal{N}$, there exists a tangent vector $\zeta \in T_x \mathcal{M}$ such that $q_x = h \circ R_x$ satisfies

$$q_x(\eta) \geq q_x(\xi) + \langle \zeta, \eta - \xi \rangle_x \quad \forall \eta, \xi \in \mathcal{S}_x. \quad (1)$$

Note that $\zeta = \text{grad } q_x(\xi)$ if h is differentiable; otherwise, ζ is any subgradient of q_x at ξ .

Proximal Gradient Method

Riemannian version in [HW21a]

Additional Assumptions:

- f and g are retraction-convex in $\Omega \supseteq \Omega_{x_0}$;

Lemma

Given $x \in \mathcal{M}$ and a twice continuously differentiable function $h : \mathcal{M} \rightarrow \mathbb{R}$, if one of the following conditions holds:

- *Hess h is positive definite at x , and the retraction is second order;*
- *The manifold \mathcal{M} is an embedded submanifold of \mathbb{R}^n endowed with the Euclidean metric; \mathcal{W} is an open subset of \mathbb{R}^n ; $x \in \mathcal{W}$;
 $h : \mathcal{W} \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is a μ -strongly convex function in the Euclidean setting for a sufficient large μ ; the retraction is second order;*

then there exists a neighborhood of x , denoted by \mathcal{N}_x , such that the function $h : \mathcal{M} \rightarrow \mathbb{R}$ is retraction-convex in \mathcal{N}_x .

Proximal Gradient Method



Riemannian version in [HW21a]

Additional Assumptions:

- f and g are retraction-convex in $\Omega \supseteq \Omega_{x_0}$;

Nonsmooth? Example: $h(x) = \|x\|_1$ with exponential mapping

- unit sphere: $\{x \in \mathbb{R}^n \mid x^T x = 1\}$, $n = 100$
- Poincaré ball model [GBH18]: $\{x \in \mathbb{R}^n \mid x^T x < 1\}$, $n = 100$
- $h(\text{Exp}_x(t\eta_x))$ versus t



[GBH18] Ganea et al., Hyperbolic entailment cones for learning hierarchical embedding, ICML, 2018.

Proximal Gradient Method

Riemannian version in [HW21a]

Additional Assumptions:

- f and g are retraction-convex in $\Omega \supseteq \Omega_{x_0}$;
- Retraction approximately satisfies the triangle relation in Ω : for all $x, y, z \in \Omega$,

$$\left| \|\xi_x - \eta_x\|_x^2 - \|\zeta_y\|_y^2 \right| \leq \kappa \|\eta_x\|_x^2, \text{ for a constant } \kappa$$

where $\eta_x = R_x^{-1}(y)$, $\xi_x = R_x^{-1}(z)$, $\zeta_y = R_y^{-1}(z)$.

-
- In the Euclidean setting: $\eta_x = R_x^{-1}(y) = y - x$, $\xi_x = R_x^{-1}(z) = z - x$, $\zeta_y = R_y^{-1}(z) = z - y$:

$$\xi_x - \eta_x = (z - x) - (y - x) = z - y = \zeta_y.$$

- Holds for compact set $\overline{\Omega}$ with the exponential mapping;

Proximal Gradient Method

Riemannian version in [HW21a]

Additional Assumptions:

- f and g are retraction-convex in $\Omega \supseteq \Omega_{x_0}$;
- Retraction approximately satisfies the triangle relation in Ω : for all $x, y, z \in \Omega$,

$$\left| \|\xi_x - \eta_x\|_x^2 - \|\zeta_y\|_y^2 \right| \leq \kappa \|\eta_x\|_x^2, \text{ for a constant } \kappa$$

where $\eta_x = R_x^{-1}(y)$, $\xi_x = R_x^{-1}(z)$, $\zeta_y = R_y^{-1}(z)$.

Theoretical results:

- Convergence rate $O(1/k)$:

$$F(x_k) - F(x_*) \leq \frac{1}{k} \left(\frac{L}{2} \|R_{x_0}^{-1}(x_*)\|_{x_0}^2 + \frac{L\kappa C}{2} (F(x_0) - F(x_*)) \right).$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- 1 Assumptions for the global convergence

-
- 1 The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;
 - 2 The function f is L -retraction-smooth with respect to the retraction R in the sublevel set Ω_{x_0} .

$$\min_{X \in \text{St}(p,n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- 1 Assumptions for the global convergence
- 2 f is locally Lipschitz continuously differentiable

Definition ([AMS08, 7.4.3])

A function f on \mathcal{M} is Lipschitz continuously differentiable if it is differentiable and if there exists β_1 such that, for all x, y in \mathcal{M} with $\text{dist}(x, y) < i(\mathcal{M})$, it holds that

$$\|\mathcal{P}_\gamma^{0 \leftarrow 1} \text{grad } f(y) - \text{grad } f(x)\|_x \leq \beta_1 \text{dist}(x, y),$$

where γ is the unique minimizing geodesic with $\gamma(0) = x$ and $\gamma(1) = y$.

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- 1 Assumptions for the global convergence
- 2 f is locally Lipschitz continuously differentiable

If f is smooth and the manifold \mathcal{M} is compact, then the function f is Lipschitz continuously differentiable. [AMS08, Proposition 7.4.5 and Corollary 7.4.6].

$$\min_{X \in \text{St}(p,n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- ① Assumptions for the global convergence
- ② f is locally Lipschitz continuously differentiable
- ③ F satisfies the Riemannian KL property [BdCNO11]

Definition

A continuous function $f : \mathcal{M} \rightarrow \mathbb{R}$ is said to have the Riemannian KL property at $x \in \mathcal{M}$ if and only if there exists $\varepsilon \in (0, \infty]$, a neighborhood $U \subset \mathcal{M}$ of x , and a continuous concave function $\varsigma : [0, \varepsilon] \rightarrow [0, \infty)$ such that

- $\varsigma(0) = 0$, ς is C^1 on $(0, \varepsilon)$, and $\varsigma' > 0$ on $(0, \eta)$,
- For every $y \in U$ with $f(x) < f(y) < f(x) + \varepsilon$, we have

$$\varsigma'(f(y) - f(x)) \operatorname{dist}(0, \partial f(y)) \geq 1,$$

where $\operatorname{dist}(0, \partial f(y)) = \inf \{\|v\|_y : v \in \partial f(y)\}$ and ∂ denotes the Riemannian generalized subdifferential. The function ς is called the desingularising function.

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- ① Assumptions for the global convergence
 - ② f is locally Lipschitz continuously differentiable
 - ③ F satisfies the Riemannian KL property [BdCNO11]
-

Theoretical results:

- it holds that

$$\sum_{k=0}^{\infty} \text{dist}(\mathbf{x}_k, \mathbf{x}_{k+1}) < \infty.$$

Therefore, there exists only a unique accumulation point.

Proximal Gradient Method

Riemannian version in [HW21a]

Assumption:

- ① Assumptions for the global convergence
 - ② f is locally Lipschitz continuously differentiable
 - ③ F satisfies the Riemannian KL property [BdCNO11]
-

Theoretical results:

- If the desingularising function has the form $\varsigma(t) = \frac{C}{\theta} t^\theta$ for $C > 0$ and $\theta \in (0, 1]$ for all $x \in \Omega_{x_0}$, then
 - if $\theta = 1$, then the Riemannian proximal gradient method terminates in finite steps;
 - if $\theta \in [0.5, 1)$, then $\|x_k - x_*\| < C_1 d^k$ for $C_1 > 0$ and $d \in (0, 1)$;
 - if $\theta \in (0, 0.5)$, then $\|x_k - x_*\| < C_2 k^{\frac{-1}{1-2\theta}}$ for $C_2 > 0$;

Proximal Gradient Method

Numerical experiments

Sparse PCA problem

$$\min_{X \in \text{St}(p,n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

where $A \in \mathbb{R}^{m \times n}$ is a data matrix.

Proximal Gradient Method

Numerical experiments




Figure: Two typical runs of ManPG, RPG, A-ManPG, and A-RPG for the Sparse PCA problem. $n = 1024$, $p = 4$, $\lambda = 2$, $m = 20$.

Optimization with Structure:

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

- Proximal gradient methods
- Inexact proximal gradient methods
 - Inexact version of RPG [HW21a]
 - Inexact version of ManPG [HWGVD22]
- A proximal Newton method

Both ManPG and RPG require the Riemannian proximal mapping to be solved exactly

- Theoretically, but not practical numerically
- Can we relax this requirement and still preserve desired convergence properties?
- Inexact RPG
- Inexact ManPG

Inexact Proximal Gradient Method

Inexact RPG

Inexact RPG (IRPG)

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_x \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

- 2 $x_{k+1} = R_{x_k}(\eta_k)$;

Inexact Proximal Gradient Method

Inexact RPG

Inexact RPG (IRPG)

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_{x_k} \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

- 2 $x_{k+1} = R_{x_k}(\eta_k)$;

Four choices of q lead to different convergence results:

- 1) **Global** $q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \varepsilon_k$ with $\varepsilon_k \rightarrow 0$;
- 2) **Global** $q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \tilde{q}(\|\hat{\eta}_{x_k}\|)$ with $\tilde{q} : \mathbb{R} \rightarrow [0, \infty)$ a continuous function satisfying $\tilde{q}(0) = 0$;
- 3) **Unique** $q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \varepsilon_k^2$, with $\sum_{k=0}^{\infty} \varepsilon_k < \infty$; and
- 4) **Rate** $q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \min(\varepsilon_k^2, \delta_q \|\hat{\eta}_{x_k}\|^2)$ with a constant $\delta_q > 0$ and $\sum_{k=0}^{\infty} \varepsilon_k < \infty$.

Inexact Proximal Gradient Method

Inexact RPG

Inexact RPG (IRPG)

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- ① Find $\hat{\eta}_k \in T_x \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

- ② $x_{k+1} = R_{x_k}(\eta_k)$;

Not a Riemannian generalization of any of the existing
Euclidean inexact proximal gradient methods

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- [Com04]: Patrick L. Combettes. Solving monotone inclusions via compositions of nonexpansive averaged operators. *Optimization*, 53(5-6):475–504, 2004.
- [FP11]: J. M. Fadili, and G. Peyre, Total variation projection with first order schemes. *IEEE Transactions on Image Processing*, 20(3), 657-669, 2001.
- [SRB11]: M. Schmidt, N. Roux, and F. Bach. Convergence rates of inexact proximal-gradient methods for convex optimization. *NIPS*, 2001.
- [VSBV13]: S. Villa, S. Salzo, L. Baldassarre, and A. Verri. Accelerated and inexact forward-backward algorithms. *SIAM Journal on Optimization*, 23(3), 1607-1633, 2013
- [BPR20]: S. Bonettini, M. Prato, and S. Rebegoldi. Convergence of inexact forward-backward algorithms using the forward-backward envelope. *SIAM Journal on Optimization*, 30(4), 3069-3097, 2020

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- $z = \text{Prox}_{\lambda g}(y) = \operatorname{argmin}_x \Phi_\lambda(x) := \lambda h(x) + \frac{1}{2} \|x - y\|^2;$

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- $z = \text{Prox}_{\lambda g}(y) = \operatorname{argmin}_x \Phi_\lambda(x) := \lambda h(x) + \frac{1}{2}\|x - y\|^2$;
- z satisfies

$$(y - z)/\lambda \in \partial^E h(z) \text{ and } \operatorname{dist}(0, \partial^E \Phi_\lambda(z)) = 0.$$

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- $z = \text{Prox}_{\lambda g}(y) = \operatorname{argmin}_x \Phi_\lambda(x) := \lambda h(x) + \frac{1}{2}\|x - y\|^2$;
- z satisfies

$$(y - z)/\lambda \in \partial^E h(z) \text{ and } \operatorname{dist}(0, \partial^E \Phi_\lambda(z)) = 0.$$

- Approximation \hat{z} satisfies any one of the following conditions:

$$\operatorname{dist}(0, \partial^E \Phi_\lambda(\hat{z})) \leq \frac{\varepsilon}{\lambda}, \quad \Phi_\lambda(\hat{z}) \leq \min \Phi_\lambda + \frac{\varepsilon^2}{2\lambda}, \text{ and } \frac{y - \hat{z}}{\lambda} \in \partial_{\frac{\varepsilon^2}{2\lambda}}^E h(\hat{z}),$$

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- $z = \text{Prox}_{\lambda g}(y) = \operatorname{argmin}_x \Phi_\lambda(x) := \lambda h(x) + \frac{1}{2}\|x - y\|^2$;
- z satisfies

$$(y - z)/\lambda \in \partial^E h(z) \text{ and } \operatorname{dist}(0, \partial^E \Phi_\lambda(z)) = 0.$$

- Approximation \hat{z} satisfies any one of the following conditions:

$$\operatorname{dist}(0, \partial^E \Phi_\lambda(\hat{z})) \leq \frac{\varepsilon}{\lambda}, \quad \Phi_\lambda(\hat{z}) \leq \min \Phi_\lambda + \frac{\varepsilon^2}{2\lambda}, \text{ and } \frac{y - \hat{z}}{\lambda} \in \partial_{\frac{\varepsilon^2}{2\lambda}}^E h(\hat{z}),$$

- Algorithms based on strong convexity of the Euclidean proximal mapping

Inexact Proximal Gradient Method

Inexact RPG

Inexact proximal gradient methods in the Euclidean setting:
[Com04, FP11, SRB11, VSBV13, BPR20]

- $z = \text{Prox}_{\lambda g}(y) = \operatorname{argmin}_x \Phi_\lambda(x) := \lambda h(x) + \frac{1}{2}\|x - y\|^2$;
- z satisfies

$$(y - z)/\lambda \in \partial^E h(z) \text{ and } \operatorname{dist}(0, \partial^E \Phi_\lambda(z)) = 0.$$

- Approximation \hat{z} satisfies any one of the following conditions:

$$\operatorname{dist}(0, \partial^E \Phi_\lambda(\hat{z})) \leq \frac{\varepsilon}{\lambda}, \quad \Phi_\lambda(\hat{z}) \leq \min \Phi_\lambda + \frac{\varepsilon^2}{2\lambda}, \text{ and } \frac{y - \hat{z}}{\lambda} \in \partial_{\frac{\varepsilon}{2\lambda}}^E h(\hat{z}),$$

- Algorithms based on strong convexity of the Euclidean proximal mapping
- Riemannian: may not be convex

$$\ell_{x_k}(\eta) = \langle \operatorname{grad} f(x_k), \eta \rangle_{x_k} + \frac{L}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$$

Inexact Proximal Gradient Method

Inexact RPG

Assumption (same as the RPG):

- 1 The function F is bounded from below and the sublevel set $\Omega_{x_0} = \{x \in \mathcal{M} \mid F(x) \leq F(x_0)\}$ is compact;
 - 2 The function f is L -retraction-smooth with respect to the retraction R in the sublevel set Ω_{x_0} .
-

Theoretical results:

- Suppose $\lim_{k \rightarrow \infty} q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = 0$, then for any accumulation point x_* of $\{x_k\}$, x_* is a stationary point, i.e., $0 \in \partial F(x_*)$.

Inexact Proximal Gradient Method

Inexact RPG

Assumption:

- 1 Assumptions for the global convergence
 - 2 f is locally Lipschitz continuously differentiable
 - 3 F satisfies the Riemannian KL property
-

Inexact Proximal Gradient Method

Inexact RPG

Assumption:

- ① Assumptions for the global convergence
 - ② f is locally Lipschitz continuously differentiable
 - ③ F satisfies the Riemannian KL property
 - ④ F is locally Lipschitz continuous with respect to the retraction R
-

Definition

A function $h : \mathcal{M} \rightarrow \mathbb{R}$ is called locally Lipschitz continuous with respect to a retraction R if for any compact subset \mathcal{N} of \mathcal{M} , there exists a constant L_h such that for any $x \in \mathcal{N}$ and $\xi_x, \eta_x \in T_x \mathcal{M}$ satisfying $R_x(\xi_x) \in \mathcal{N}$ and $R_x(\eta_x) \in \mathcal{N}$, it holds that $|h \circ R(\xi_x) - h \circ R(\eta_x)| \leq L_h \|\xi_x - \eta_x\|$.

Inexact Proximal Gradient Method

Inexact RPG

Assumption:

- ① Assumptions for the global convergence
 - ② f is locally Lipschitz continuously differentiable
 - ③ F satisfies the Riemannian KL property
 - ④ F is locally Lipschitz continuous with respect to the retraction R
-

If the manifold \mathcal{M} is an embedded submanifold and function F is locally Lipschitz in the embedding space, then the function is locally Lipschitz continuous with respect to any global defined retraction R .

Inexact Proximal Gradient Method

Inexact RPG

Assumption:

- ① Assumptions for the global convergence
 - ② f is locally Lipschitz continuously differentiable
 - ③ F satisfies the Riemannian KL property
 - ④ F is locally Lipschitz continuous with respect to the retraction R
-

Theoretical results:

- If $\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq \varepsilon_k^2$ for $\sum_{k=0}^{\infty} \varepsilon_k < \infty$ and $\varepsilon_k > 0$, then it holds that

$$\sum_{k=0}^{\infty} \text{dist}(x_k, x_{k+1}) < \infty.$$

Therefore, there exists only a unique accumulation point.

Inexact Proximal Gradient Method

Inexact RPG

Assumption:

- 1 Assumptions for the global convergence
 - 2 f is locally Lipschitz continuously differentiable
 - 3 F satisfies the Riemannian KL property
 - 4 F is locally Lipschitz continuous with respect to the retraction R
-

Theoretical results:

- If $\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq \min\left(\varepsilon_k^2, \frac{\beta}{2L_F} \|\hat{\eta}_{x_k}\|^2\right)$ for $\sum_{k=0}^{\infty} \varepsilon_k < \infty$ and $\varepsilon_k > 0$, and if the desingularising function has the form $\varsigma(t) = \frac{C}{\theta} t^\theta$ for $C > 0$ and $\theta \in (0, 1]$ for all $x \in \Omega_{x_0}$, then
 - if $\theta = 1$, then the Riemannian proximal gradient method terminates in finite steps;
 - if $\theta \in [0.5, 1)$, then $\|x_k - x_*\| < C_1 d^k$ for $C_1 > 0$ and $d \in (0, 1)$;
 - if $\theta \in (0, 0.5)$, then $\|x_k - x_*\| < C_2 k^{\frac{-1}{1-2\theta}}$ for $C_2 > 0$;

Inexact Proximal Gradient Method

Inexact RPG

IRPG

Let $\ell_{x_k}(\eta) = \langle \text{grad}f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_{x_k} \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

How to find $\hat{\eta}_k$ for different q ?

Inexact Proximal Gradient Method

Inexact RPG

IRPG

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_{x_k} \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

How to find $\hat{\eta}_k$ for different q ?

- Only consider manifolds with a linear ambient space;

Inexact Proximal Gradient Method

Inexact RPG

IRPG

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_{x_k} \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

How to find $\hat{\eta}_k$ for different q ?

- Only consider manifolds with a linear ambient space;
- Use the semi-smooth Newton method iteratively;

Inexact Proximal Gradient Method

Inexact RPG

IRPG

Let $\ell_{x_k}(\eta) = \langle \text{grad} f(x_k), \eta \rangle_{x_k} + \frac{\tilde{L}}{2} \|\eta\|_{x_k}^2 + h(R_{x_k}(\eta))$;

- 1 Find $\hat{\eta}_k \in T_{x_k} \mathcal{M}$ such that

$$\|\hat{\eta}_{x_k} - \eta_{x_k}^*\| \leq q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) \text{ and } \ell_{x_k}(0) \geq \ell_{x_k}(\hat{\eta}_{x_k}),$$

where $\varepsilon_k > 0$, and $q : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function;

How to find $\hat{\eta}_k$ for different q ?

- Only consider manifolds with a linear ambient space;
- Use the semi-smooth Newton method iteratively;
- For sufficiently large \tilde{L} , η_k from ManPG guarantees global convergence;

Inexact Proximal Gradient Method

Inexact RPG

ManPG [CMSZ20]

$$\eta_k = \arg \min_{\eta \in T_{x_k} \mathcal{M}} \langle \nabla f(x_k), \eta \rangle + \frac{L}{2} \|\eta\|_F^2 + h(x_k + \eta)$$

Above problem can be rewritten as

$$\arg \min_{B_x^T \eta = 0} \langle \xi_x, \eta \rangle + \frac{1}{2\mu} \|\eta\|_F^2 + h(x + \eta)$$

where $B_x^T \eta = (\langle b_1, \eta \rangle, \langle b_2, \eta \rangle, \dots, \langle b_m, \eta \rangle)^T$, and $\{b_1, \dots, b_m\}$ forms an orthonormal basis of $N_x \mathcal{M}$.

Inexact Proximal Gradient Method

Inexact RPG

The Lagrangian function:

$$\mathcal{L}(\eta, \Lambda) = \langle \xi_x, \eta \rangle + \frac{1}{2\mu} \langle \eta, \eta \rangle + h(X + \eta) - \langle \Lambda, B_x^T \eta \rangle.$$

Therefore

$$\text{KKT: } \begin{cases} \partial_{\eta} \mathcal{L}(\eta, \Lambda) = 0 \\ B_x^T \eta = 0 \end{cases} \implies \begin{cases} \eta = \text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x \\ B_x^T \eta = 0 \end{cases}$$

where $\text{Prox}_{\mu g}(z) = \operatorname{argmin}_{v \in \mathbb{R}^{n \times p}} \frac{1}{2} \|v - z\|_F^2 + \mu h(v)$.

Inexact Proximal Gradient Method

Inexact RPG

Semi-smooth Newton method finds the Λ such that

$$\begin{aligned}\Psi(\Lambda) &:= B_x^T (\text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x) = 0 \\ \eta_* &= \text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x\end{aligned}$$

- Ψ is not differentiable everywhere but semi-smooth for $h(\cdot) = \|\cdot\|_1$;
- Semi-smooth Newton:
 - 1 $J_\Psi(\Lambda_k)[d] = -\Psi(\Lambda_k)$, where J_Ψ is the generalized Jacobian of Ψ ;
 - 2 $\Lambda_{k+1} = \Lambda_k + d_k$

Inexact Proximal Gradient Method

Inexact RPG

Semi-smooth Newton method finds the Λ such that

$$\Psi(\Lambda) := B_x^T (\text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x) \approx 0$$

- Ψ is not differentiable everywhere but semi-smooth for $h(\cdot) = \|\cdot\|_1$;
- Semi-smooth Newton:
 - ① $J_\Psi(\Lambda_k)[d] = -\Psi(\Lambda_k)$, where J_Ψ is the generalized Jacobian of Ψ ;
 - ② $\Lambda_{k+1} = \Lambda_k + d_k$
- Solving the equation inexactly

Inexact Proximal Gradient Method

Inexact RPG

If $\Psi(\Lambda) = \epsilon$,

- $\eta_* = \text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x$ is not even in the tangent space $T_x \mathcal{M}$ in this case
- Use $\hat{\eta}_x := \hat{v}(\Lambda) = P_{T_x \mathcal{M}}(\text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x)$ instead
- How small does ϵ need to be?

Inexact Proximal Gradient Method

Inexact RPG

If $\Psi(\Lambda) = \epsilon$,

- $\eta_* = \text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x$ is not even in the tangent space $T_x \mathcal{M}$ in this case
- Use $\hat{\eta}_x := \hat{v}(\Lambda) = P_{T_x \mathcal{M}}(\text{Prox}_{\mu g}(x - \mu(\xi_x - B_x \Lambda)) - x)$ instead
- How small does ϵ need to be?

$$\|\epsilon\| \leq \min(\phi(\hat{v}(\Lambda)), 0.5),$$

with $\phi(0) = 0$ and ϕ is nondecreasing.

Inexact Proximal Gradient Method

Inexact RPG

The function q is:

$$q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \frac{2L_h\kappa_2}{\tilde{L} - 2L_h\kappa_2} \|\hat{\eta}_{x_k}\| + \sqrt{\frac{4L_h\kappa_2 - 4L_h^2\kappa_2^2}{(\tilde{L} - 2L_h\kappa)^2} \|\hat{\eta}_{x_k}\|^2 + \frac{4\vartheta}{\tilde{L} - 2L_h\kappa_2} \min(\phi(\|\hat{\eta}_{x_k}\|), 0.5)}$$

- ManPG can be viewed as an inexact RPG for sufficiently large \tilde{L} ;

Inexact Proximal Gradient Method

Inexact RPG

The function q is:

$$q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) = \frac{2L_h\kappa_2}{\tilde{L} - 2L_h\kappa_2} \|\hat{\eta}_{x_k}\| + \sqrt{\frac{4L_h\kappa_2 - 4L_h^2\kappa_2^2}{(\tilde{L} - 2L_h\kappa)^2} \|\hat{\eta}_{x_k}\|^2 + \frac{4\vartheta}{\tilde{L} - 2L_h\kappa_2} \min(\phi(\|\hat{\eta}_{x_k}\|), 0.5)}$$

- ManPG can be viewed as an inexact RPG for sufficiently large \tilde{L} ;
- This q may not guarantee local convergence results;

Inexact Proximal Gradient Method

Inexact RPG

The function q is:

$$q(\varepsilon_k, \|\hat{\eta}_{x_k}\|) =$$

$$\frac{2L_h\kappa_2}{\tilde{L} - 2L_h\kappa_2} \|\hat{\eta}_{x_k}\| + \sqrt{\frac{4L_h\kappa_2 - 4L_h^2\kappa_2^2}{(\tilde{L} - 2L_h\kappa_2)^2} \|\hat{\eta}_{x_k}\|^2 + \frac{4\vartheta}{\tilde{L} - 2L_h\kappa_2} \min(\phi(\|\hat{\eta}_{x_k}\|), 0.5)}$$

- ManPG can be viewed as an inexact RPG for sufficiently large \tilde{L} ;
- This q may not guarantee local convergence results;
- Improving accuracy is needed;

Inexact Proximal Gradient Method

Inexact RPG

$$\eta_x = \arg \min_{\eta \in T_x \mathcal{M}} \ell_x(\eta) := \langle \nabla f(x), \eta \rangle_x + \frac{L}{2} \|\eta\|_x^2 + h(R_x(\eta))$$

Solving the Riemannian Proximal Mapping [HW21a]

initial iterate: $\eta_0 \in T_x \mathcal{M}$, $\sigma \in (0, 1)$, $k = 0$;

- ① $y_k = R_x(\eta_k)$;
- ② Compute
$$\xi_k^* \approx \arg \min_{\xi \in T_{y_k} \mathcal{M}} \langle \mathcal{T}_{R_{\eta_k}}^{-\sharp}(\text{grad } f(x) + \tilde{L}\eta_k), \xi \rangle_x + \frac{\tilde{L}}{4} \|\xi\|_F^2 + h(y_k + \xi);$$
- ③ Find $\alpha > 0$ such that $\ell_x(\eta_k + \alpha \mathcal{T}_{R_{\eta_k}}^{-1} \xi_k^*) < \ell_x(\eta_k) - \sigma \alpha \|\xi_k^*\|_x^2$;
- ④ $\eta_{k+1} = \eta_k + \alpha \mathcal{T}_{R_{\eta_k}}^{-1} \xi_k^*$;
- ⑤ If $\|\xi_k^*\|$ is sufficiently small, then stop;
- ⑥ $k \leftarrow k + 1$ and goto Step 1;

Inexact RPG for global convergence (IRPG)

- 1 Approximately solve

$$\min_{\eta \in T_{x_k} \mathcal{M}} \langle \text{grad } f(x_k), \eta \rangle + \frac{\tilde{L}}{2} \|\eta\|_F^2 + h(x_k + \eta)$$

such that $\|\Psi_k(\Lambda)\|_F \leq \min(\phi(\hat{v}(\Lambda)), 0.5)$;

- 2 Let $\eta_k = \hat{v}(\Lambda)$;
- 3 $x_{k+1} = R_{x_k}(\eta_k)$;

-
- Global convergence requires a sufficient large of \tilde{L} ;
 - Step size one is used;

Inexact Proximal Gradient Method

Inexact ManPG

Inexact RPG for global convergence (IRPG)

- 1 Approximately solve

$$\min_{\eta \in T_{x_k} \mathcal{M}} \langle \text{grad } f(x_k), \eta \rangle + \frac{\tilde{L}}{2} \|\eta\|_F^2 + h(x_k + \eta)$$

such that $\|\Psi_k(\Lambda)\|_F \leq \min(\phi(\hat{v}(\Lambda)), 0.5)$;

- 2 Let $\eta_k = \hat{v}(\Lambda)$;
- 3 $x_{k+1} = R_{x_k}(\eta_k)$;

-
- Global convergence requires a sufficient large of \tilde{L} ;
 - Step size one is used;
 - Is $\hat{\eta}_x$ a descent direction for any positive \tilde{L} ?

Inexact Proximal Gradient Method

Inexact ManPG

Algorithm 1 ManPG without solving the subproblem exactly

- 1: Given x_0 , $\nu \in (0, 1)$, $\sigma \in (0, 1/(8\mu))$, $\mu > 0$;
- 2: **for** $k = 0, 1, \dots$ **do**
- 3: Approximately solve

$$\min_{\eta \in T_{x_k} \mathcal{M}} \langle \text{grad } f(x_k), \eta \rangle + \frac{1}{2\mu} \|\eta\|_F^2 + h(x_k + \eta)$$

such that $\|\Psi_k(\Lambda)\|_F \leq \sqrt{4\mu^2 L_h^2 + \|\hat{v}_k(\Lambda)\|_F^2/2} - 2\mu L_h$;

- 4: Set $\eta_k = \hat{v}_k(\Lambda)$ and set $\alpha = 1$;
 - 5: **while** $F(R_{x_k}(\alpha\eta_{x_k})) > F(x_k) - \sigma\alpha\|\eta_{x_k}\|_F^2$ **do**
 - 6: $\alpha = \nu\alpha$;
 - 7: **end while**
 - 8: $x_{k+1} = R_{x_k}(\alpha\eta_{x_k})$;
 - 9: **end for**
-

Inexact Proximal Gradient Method

Inexact ManPG

Assumption

The function f is Lipschitz continuously differentiable on \mathcal{M} and h is Lipschitz continuous with constant L_h .

Theorem

Suppose the assumption holds. Then for any $\mu > 0$, there exists a constant $\bar{\alpha} \in (0, 1]$ such that for any $0 < \alpha < \bar{\alpha}$, the sequence $\{x_k\}$ generated by Algorithm 1 satisfies

$$F(R_{x_k}(\alpha\eta_{x_k})) - F(x_k) \leq -\frac{\alpha}{8\mu} \|\eta_{x_k}\|_F^2.$$

Moreover, the step size $\alpha > \rho\bar{\alpha}$ for all k .

Inexact Proximal Gradient Method

Inexact ManPG

Assumption

The function f is Lipschitz continuously differentiable on \mathcal{M} and h is Lipschitz continuous with constant L_h .

Theorem

Suppose the assumption holds. Then any accumulation point of the sequence $\{x_k\}$ generated by Algorithm 1 is a stationary point, i.e., if x_ is an accumulation point of the above sequence, then $0 \in P_{T_{x_*} \mathcal{M}} \partial F(x_*)$.*

Inexact Proximal Gradient Method

Numerical experiments

Sparse PCA problem

$$\min_{X \in \text{St}(p,n)} -\text{trace}(X^T A^T A X) + \lambda \|X\|_1,$$

where $A \in \mathbb{R}^{m \times n}$ is a data matrix.

Inexact Proximal Gradient Method

Numerical experiments




Figure: Average of 10 random runs, $p = 4$, $m = 20$, $\lambda = 2$;

- IRPG-G: an inexact version of ManPG
- IRPG-U: $\psi = \varepsilon_k^2$
- IRPG-L: $\psi = \min(\varepsilon_k^2, \varrho \|\hat{\eta}_{x_k}\|^2)$

Inexact Proximal Gradient Method

Numerical experiments

Community detection:

$$\min_{X \in \mathcal{F}_{\mathbf{1}_n}} -\text{trace}(X^T M X) + \lambda \|X\|_1,$$

where $\mathcal{F}_{\mathbf{1}_n} = \{X \in \mathbb{R}^{n \times k} : X^T X = I_k, \mathbf{1}_n \in \text{span}(X)\}$

Inexact Proximal Gradient Method

Numerical experiments

Comparing efficiency of I-AManPG and E-AManPG

	I-A	E-A	I-A	E-A	I-A	E-A	I-A	E-A
(n, q)	(5000, 10)	(5000, 10)	(5000, 20)	(5000, 20)	(10000, 10)	(10000, 10)	(10000, 20)	(10000, 20)
iter	63	58	47	50	55	55	73	51
SSNiter	34	311	32	381	52	330	146	376
nf	140	128	105	112	123	122	161	113
ng	81	72	60	62	71	68	92	64
nR	139	127	104	111	122	121	160	112
nSG	4	13	2	5	3	10	3	13
F	-2.84_2	-2.84_2	-6.55_2	-6.56_2	-2.51_2	-2.51_2	-6.11_2	-6.14_2
$\ \eta_{z_k}\ $	6.31_{-4}	5.82_{-4}	5.32_{-4}	7.54_{-4}	5.22_{-4}	6.86_{-4}	4.02_{-4}	6.60_{-4}
time	0.84	3.04	1.51	9.81	1.54	5.19	9.48	19.21

- AManPG: add acceleration [HW21b]
- I-AManPG: Inexact version
- E-AManPG: Exact version, i.e., $\epsilon = 10^{-10}$
- An average of 10 random runs

Inexact Proximal Gradient Method

Numerical experiments

Comparing efficiency of I-AManPG and E-AManPG

	I-A	E-A	I-A	E-A	I-A	E-A	I-A	E-A
(n, q)	(5000, 10)		(5000, 20)		(10000, 10)		(10000, 20)	
iter	63	58	47	50	55	55	73	51
SSNiter	34	311	32	381	52	330	146	376
nf	140	128	105	112	123	122	161	113
ng	81	72	60	62	71	68	92	64
nR	139	127	104	111	122	121	160	112
nSG	4	13	2	5	3	10	3	13
F	-2.84_2	-2.84_2	-6.55_2	-6.56_2	-2.51_2	-2.51_2	-6.11_2	-6.14_2
$\frac{\ \eta_{z_k}\ }{\ \eta_{z_0}\ }$	6.31_{-4}	5.82_{-4}	5.32_{-4}	7.54_{-4}	5.22_{-4}	6.86_{-4}	4.02_{-4}	6.60_{-4}
time	0.84	3.04	1.51	9.81	1.54	5.19	9.48	19.21

Less computational time, same effectiveness

Optimization with Structure:

$$\min_{x \in \mathcal{M}} F(x) = f(x) + h(x).$$

- Proximal gradient methods
- Inexact proximal gradient methods
- A proximal Newton method
 - Euclidean inexact proximal Newton methods
 - A naive Riemannian proximal Newton method
 - A proposed Riemannian proximal Newton method

Inexact Proximal Newton Methods

Euclidean version

Given x_0 ;

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, H_k p \rangle + h(x_k + p) \\ x_{k+1} = x_k + t_k d_k, \text{ for a step size } t_k \end{cases}$$

Inexact Proximal Newton Methods

Euclidean version

Given x_0 ;

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, H_k p \rangle + h(x_k + p) \\ x_{k+1} = x_k + t_k d_k, \text{ for a step size } t_k \end{cases}$$

-
- H_k is Hessian or a positive definite approximation to Hessian [LSS14, MYZZ22];

[LSS14] Jason D Lee, Yuekai Sun, and Michael A Saunders. Proximal newton-type methods for minimizing composite functions. *SIAM Journal on Optimization*, 24(3):1420-1443, 2014.

[MYZZ22] Boris S Mordukhovich, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang. A globally convergent proximal newton-type method in nonsmooth convex optimization. *Mathematical Programming*, pages 1-38, 2022.

Inexact Proximal Newton Methods

Euclidean version

Given x_0 ;

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, H_k p \rangle + h(x_k + p) \\ x_{k+1} = x_k + t_k d_k, \text{ for a step size } t_k \end{cases}$$

-
- H_k is Hessian or a positive definite approximation to Hessian [LLS14, MYZZ22];
 - t_k is one for sufficiently large k ;

[LLS14] Jason D Lee, Yuekai Sun, and Michael A Saunders. Proximal newton-type methods for minimizing composite functions. *SIAM Journal on Optimization*, 24(3):1420-1443, 2014.

[MYZZ22] Boris S Mordukhovich, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang. A globally convergent proximal newton-type method in nonsmooth convex optimization. *Mathematical Programming*, pages 1-38, 2022.

Inexact Proximal Newton Methods

Euclidean version

Given x_0 ;

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, H_k p \rangle + h(x_k + p) \\ x_{k+1} = x_k + t_k d_k, \text{ for a step size } t_k \end{cases}$$

-
- H_k is Hessian or a positive definite approximation to Hessian [LSS14, MYZZ22];
 - t_k is one for sufficiently large k ;
 - Quadratic/Superlinear convergence rate for strongly convex f and convex h ;

[LLS14] Jason D Lee, Yuekai Sun, and Michael A Saunders. Proximal newton-type methods for minimizing composite functions. *SIAM Journal on Optimization*, 24(3):1420-1443, 2014.

[MYZZ22] Boris S Mordukhovich, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang. A globally convergent proximal newton-type method in nonsmooth convex optimization. *Mathematical Programming*, pages 1-38, 2022.

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Focus on embedded submanifolds

Euclidean version:

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, \nabla^2 f(x_k) p \rangle + h(x_k + p) \\ x_{k+1} = x_k + d_k \end{cases}$$

A native generalization by replacing the Euclidean gradient and Hessian by the Riemannian gradient and Hessian:

$$\begin{cases} \eta_k = \operatorname{argmin}_{\eta \in T_{x_k} \mathcal{M}} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Focus on embedded submanifolds

Euclidean version:

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, \nabla^2 f(x_k) p \rangle + h(x_k + p) \\ x_{k+1} = x_k + d_k \end{cases}$$

A native generalization by replacing the Euclidean gradient and Hessian by the Riemannian gradient and Hessian:

$$\begin{cases} \eta_k = \operatorname{argmin}_{\eta \in T_{x_k} \mathcal{M}} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$

Does it converge superlinearly locally?

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Focus on embedded submanifolds

Euclidean version:

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, \nabla^2 f(x_k) p \rangle + h(x_k + p) \\ x_{k+1} = x_k + d_k \end{cases}$$

A native generalization by replacing the Euclidean gradient and Hessian by the Riemannian gradient and Hessian:

$$\begin{cases} \eta_k = \operatorname{argmin}_{\eta \in T_{x_k} \mathcal{M}} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$

Does it converge superlinearly locally?

Not necessarily!

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Consider the Sparse PCA over sphere:

$$\min_{x \in \mathbb{S}^{n-1}} -x^T A^T A x + \mu \|x\|_1,$$

where $f(x) = -x^T A^T A x$, $h(x) = \mu \|x\|_1$.




Figure: Comparisons of native generalization (RPN-N) and the proximal gradient method (ManPG) in [CMSZ20].

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Euclidean version:

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, \nabla^2 f(x_k) p \rangle + h(x_k + p) \\ x_{k+1} = x_k + d_k \end{cases}$$

A native generalization by replacing the Euclidean gradient and Hessian by the Riemannian gradient and Hessian:

$$\begin{cases} \eta_k = \operatorname{argmin}_{\eta \in T_{x_k}} \mathcal{M} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$

- $x_k + \eta$ in h is only a first order approximation;

Inexact Proximal Newton Methods

Riemannian version: a naive generalization

Euclidean version:

$$\begin{cases} d_k = \operatorname{argmin}_p \langle \nabla f(x_k), p \rangle + \frac{1}{2} \langle p, \nabla^2 f(x_k) p \rangle + h(x_k + p) \\ x_{k+1} = x_k + d_k \end{cases}$$

A native generalization by replacing the Euclidean gradient and Hessian by the Riemannian gradient and Hessian:

$$\begin{cases} \eta_k = \operatorname{arg min}_{\eta \in T_{x_k}} \mathcal{M} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$
$$\begin{cases} \eta_k = \operatorname{arg min}_{\eta \in T_{x_k}} \mathcal{M} \langle \operatorname{grad} f(x_k), \eta \rangle + \frac{1}{2} \langle \eta, \operatorname{Hess} f(x_k) \eta \rangle + h(x_k + \eta + \frac{1}{2} \Pi(\eta, \eta)) \\ x_{k+1} = R_{x_k}(\eta_k) \end{cases}$$

- $x_k + \eta$ in h is only a first order approximation;
- If a second order approximation is used, then the subproblem is difficult to solve;

Inexact Proximal Newton Methods

Riemannian version

A Riemannian proximal Newton method (RPN)

- 1 Compute

$$v(x_k) = \operatorname{argmin}_{v \in T_{x_k} \mathcal{M}} f(x_k) + \langle \nabla f(x_k), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x_k + v);$$

- 2 Find $u(x_k) \in T_{x_k} \mathcal{M}$ by solving

$$J(x_k)[u(x_k)] = -v(x_k),$$

where $J(x_k) = -[I_n - \Lambda_{x_k} + t\Lambda_{x_k}(\nabla^2 f(x_k) - \mathcal{L}_{x_k})]$, Λ_{x_k} and \mathcal{L}_{x_k} are defined later ;

- 3 $x_{k+1} = R_{x_k}(u(x_k));$

Inexact Proximal Newton Methods

Riemannian version

A Riemannian proximal Newton method (RPN)

① Compute

$$v(x_k) = \operatorname{argmin}_{v \in T_{x_k} \mathcal{M}} f(x_k) + \langle \nabla f(x_k), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x_k + v);$$

② Find $u(x_k) \in T_{x_k} \mathcal{M}$ by solving

$$J(x_k)[u(x_k)] = -v(x_k),$$

where $J(x_k) = -[I_n - \Lambda_{x_k} + t\Lambda_{x_k}(\nabla^2 f(x_k) - \mathcal{L}_{x_k})]$, Λ_{x_k} and \mathcal{L}_{x_k} are defined later ;

③ $x_{k+1} = R_{x_k}(u(x_k));$

① Step 1: compute a Riemannian proximal gradient direction (ManPG)

Inexact Proximal Newton Methods

Riemannian version

A Riemannian proximal Newton method (RPN)

- 1 Compute

$$v(x_k) = \operatorname{argmin}_{v \in T_{x_k} \mathcal{M}} f(x_k) + \langle \nabla f(x_k), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x_k + v);$$

- 2 Find $u(x_k) \in T_{x_k} \mathcal{M}$ by solving

$$J(x_k)[u(x_k)] = -v(x_k),$$

where $J(x_k) = -[I_n - \Lambda_{x_k} + t\Lambda_{x_k}(\nabla^2 f(x_k) - \mathcal{L}_{x_k})]$, Λ_{x_k} and \mathcal{L}_{x_k} are defined later ;

- 3 $x_{k+1} = R_{x_k}(u(x_k));$

- 1 Step 1: compute a Riemannian proximal gradient direction (ManPG)
- 2 Step 2: compute the Riemannian proximal Newton direction, where $J(x_k)$ is from a generalized Jacobi of $v(x_k)$;

Inexact Proximal Newton Methods

Riemannian version

A Riemannian proximal Newton method (RPN)

- 1 Compute

$$v(x_k) = \operatorname{argmin}_{v \in T_{x_k} \mathcal{M}} f(x_k) + \langle \nabla f(x_k), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x_k + v);$$

- 2 Find $u(x_k) \in T_{x_k} \mathcal{M}$ by solving

$$J(x_k)[u(x_k)] = -v(x_k),$$

where $J(x_k) = -[I_n - \Lambda_{x_k} + t\Lambda_{x_k}(\nabla^2 f(x_k) - \mathcal{L}_{x_k})]$, Λ_{x_k} and \mathcal{L}_{x_k} are defined later ;

- 3 $x_{k+1} = R_{x_k}(u(x_k))$;

- 1 Step 1: compute a Riemannian proximal gradient direction (ManPG)
- 2 Step 2: compute the Riemannian proximal Newton direction, where $J(x_k)$ is from a generalized Jacobi of $v(x_k)$;
- 3 Step 3: Update iterate by a retraction;

Inexact Proximal Newton Methods

Riemannian version

A Riemannian proximal Newton method (RPN)

- 1 Compute

$$v(x_k) = \operatorname{argmin}_{v \in T_{x_k} \mathcal{M}} f(x_k) + \langle \nabla f(x_k), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x_k + v);$$

- 2 Find $u(x_k) \in T_{x_k} \mathcal{M}$ by solving

$$J(x_k)[u(x_k)] = -v(x_k),$$

where $J(x_k) = -[I_n - \Lambda_{x_k} + t\Lambda_{x_k}(\nabla^2 f(x_k) - \mathcal{L}_{x_k})]$, Λ_{x_k} and \mathcal{L}_{x_k} are defined later ;

- 3 $x_{k+1} = R_{x_k}(u(x_k));$

Next, we will show:

- 1 G-semismoothness of $v(x_k)$ and its generalized Jacobi;
- 2 Superlinear convergence rate;

Inexact Proximal Newton Methods

Riemannian version

Definition (G-Semismoothness [Gow04])

Let $F : \mathcal{D} \rightarrow \mathbb{R}^m$ where $\mathcal{D} \subset \mathbb{R}^n$ be an open set, $\mathcal{K} : \mathcal{D} \rightrightarrows \mathbb{R}^{m \times n}$ be a nonempty set-valued mapping. We say that F is G-semismooth at $x \in \mathcal{D}$ with respect to \mathcal{K} if for any $J \in \mathcal{K}(x + d)$,

$$F(x + d) - F(x) - Jd = o(\|d\|) \text{ as } d \rightarrow 0.$$

If F is G-semismooth at any $x \in \mathcal{D}$ with respect to \mathcal{K} , then F is called a G-semismooth function with respect to \mathcal{K} .

The standard definition of semismoothness additional requires:

- \mathcal{K} is compact valued, upper semicontinuous set-valued mapping;
- F is a locally Lipschitz continuous function;
- F is directionally differentiable at x ;

[Gow04] M Seetharama Gowda. Inverse and implicit function theorems for h-differentiable and semismooth functions. Optimization Methods and Software, 19(5):443-461, 2004.

Inexact Proximal Newton Methods

Riemannian version

$v(x)$ (dropping the subscript for simplicity)

$$v(x) = \operatorname{argmin}_{v \in T_x \mathcal{M}} f(x) + \langle \nabla f(x), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x + v);$$

Inexact Proximal Newton Methods

Riemannian version

$v(x)$ (dropping the subscript for simplicity)

$$v(x) = \operatorname{argmin}_{v \in T_x \mathcal{M}} f(x) + \langle \nabla f(x), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x + v);$$

Above problem can be rewritten as

$$\arg \min_{B_x^T v = 0} \langle \xi_x, v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x + v)$$

where $B_x^T v = (\langle b_1, v \rangle, \langle b_2, v \rangle, \dots, \langle b_m, v \rangle)^T$, and $\{b_1, \dots, b_m\}$ forms an orthonormal basis of $T_x^\perp \mathcal{M}$.

Inexact Proximal Newton Methods

Riemannian version

The Lagrangian function:

$$\mathcal{L}(v, \lambda) = \langle \xi_x, v \rangle + \frac{1}{2t} \langle v, v \rangle + h(X + v) - \langle \lambda, B_x^T v \rangle.$$

Therefore

$$\text{KKT: } \begin{cases} \partial_v \mathcal{L}(v, \lambda) = 0 \\ B_x^T v = 0 \end{cases} \implies \begin{cases} v = \text{Prox}_{th}(x - t(\xi_x - B_x \lambda)) - x \\ B_x^T v = 0 \end{cases}$$

where $\text{Prox}_{tg}(z) = \operatorname{argmin}_{v \in \mathbb{R}^{n \times p}} \frac{1}{2} \|v - z\|_F^2 + th(v)$.

Define

$$\mathcal{F} : \mathbb{R}^n \times \mathbb{R}^{n+d} \mapsto \mathbb{R}^{n+d} : (x; v, \lambda) \mapsto \begin{pmatrix} v + x - \text{Prox}_{th}(x - t[\nabla f(x) + B_x \lambda]) \\ B_x^T v \end{pmatrix}.$$

$v(x)$ is the solution of the system $\mathcal{F}(x, v(x), \lambda(x)) = 0$;

Inexact Proximal Newton Methods

Riemannian version

Define

$$\mathcal{F} : \mathbb{R}^n \times \mathbb{R}^{n+d} \mapsto \mathbb{R}^{n+d} : (x; v, \lambda) \mapsto \begin{pmatrix} v + x - \text{Prox}_{th} \left(x - t[\nabla f(x) + B_x \lambda] \right) \\ B_x^T v \end{pmatrix}.$$

-
- \mathcal{F} is semismooth;
 - $v(x)$ is G-semismooth by the G-semismooth Implicit Function Theorem in [Gow04, PSS03];

[Gow04] M Seetharama Gowda. Inverse and implicit function theorems for h-differentiable and semismooth functions. Optimization Methods and Software, 19(5):443-461, 2004.

[PSS03] Jong-Shi Pang, Defeng Sun, and Jie Sun. Semismooth homeomorphisms and strong stability of semidefinite and Lorentz complementarity problems. Mathematics of Operations Research, 28(1):39-63, 2003.

Inexact Proximal Newton Methods

Riemannian version

Lemma (Semismooth Implicit Function Theorem)

Suppose that $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a **semismooth** function with respect to $\partial_B F$ in an open neighborhood of (x^0, y^0) with $F(x^0, y^0) = 0$. Let $H(y) = F(x^0, y)$, if every matrix in $\partial_C H(y^0)$ is nonsingular, then there exists an open set $\mathcal{V} \subset \mathbb{R}^n$ containing x^0 , a set-valued function $\mathcal{K} : \mathcal{V} \rightarrow \mathbb{R}^{m \times n}$, and a G -semismooth function $f : \mathcal{V} \rightarrow \mathbb{R}^m$ with respect to \mathcal{K} satisfying $f(x^0) = y^0$, for every $x \in \mathcal{V}$,

$$F(x, f(x)) = 0,$$

and the set-valued function \mathcal{K} is

$$\mathcal{K} : x \mapsto \{-(A_y)^{-1}A_x : [A_x \ A_y] \in \partial_B F(x, f(x))\},$$

where the map $x \mapsto \mathcal{K}(x)$ is **compact valued and upper semicontinuous**.

Not new but an arrangement of existing results.

Inexact Proximal Newton Methods

Riemannian version

Without loss of generality, we assume that the nonzero entries of x_* are in the first part, i.e., $x_* = [\bar{x}_*^T, 0^T]^T$

Assumption

Let $B_{x_}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank.*

Inexact Proximal Newton Methods

Riemannian version

Without loss of generality, we assume that the nonzero entries of x_* are in the first part, i.e., $x_* = [\bar{x}_*^T, 0^T]^T$

Assumption

Let $B_{x_*}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank.

$v(x)$ is a G-semismooth function of x in a neighborhood of x_*

Under the above Assumption, there exists a neighborhood \mathcal{U} of x_* such that $v : \mathcal{U} \rightarrow \mathbb{R}^n : x \mapsto v(x)$ is a G-semismooth function with respect to \mathcal{K}_v , where

$$\mathcal{K}_v : x \mapsto \left\{ -[I_n, 0]B^{-1}A : [A \ B] \in \partial_B \mathcal{F}(x, v(x), \lambda(x)) \right\}.$$

For $x \in \mathcal{U}$, any element of $\mathcal{K}_v(x)$ is called a **generalized Jacobi of v at x** .

Here, the semismooth implicit function theorem is used

Inexact Proximal Newton Methods

Riemannian version

The generalized Jacobi of v at x is

$$\left\{ \mathcal{J}_x \mid \mathcal{J}_x[\omega] = - \left[I_n - \Lambda_x + t\Lambda_x(\nabla^2 f(x) - \mathcal{L}_x) \right] \omega - M_x B_x H_x (DB_x^T[\omega])v, \forall \omega \right. \\ \left. M_x \in \partial_{\mathcal{C}\text{prox}_{th}}(x) \right\},$$

where $\Lambda_x = M_x - M_x B_x H_x B_x^T M_x$, $H_x = (B_x^T M_x B_x)^{-1}$, $\mathcal{L}_x(\cdot) = \mathcal{W}_x(\cdot, B_x \lambda(x))$, and \mathcal{W}_x denotes the Weingarten map;

- $v(x_*) = 0$;
- Set $J(x) = I_n - \Lambda_x + t\Lambda_x(\nabla^2 f(x) - \mathcal{L}_x)$;
- The Riemannian proximal Newton direction: $J(x)u(x) = -v(x)$;
- Let $u(x) = (\bar{u}(x); \hat{u}(x))$, then

$$\hat{u}(x) = \hat{v} \text{ and } \bar{J}(x)\bar{u}(x) = -\bar{v}(x)$$

Inexact Proximal Newton Methods

Riemannian version

Assumption:

- ① Let $B_{x_*}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank;
-

Inexact Proximal Newton Methods

Riemannian version

Assumption:

- ① Let $B_{x_*}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank;
 - ② There exists a neighborhood \mathcal{U} of $x_* = [\bar{x}_*^T, 0^T]^T$ on \mathcal{M} such that for any $x = [\bar{x}^T, \tilde{x}^T]^T \in \mathcal{U}$, it holds that $\bar{x} + \bar{v} \neq 0$ and $\hat{x} + \hat{v} = 0$.
-

$$v(x) = \operatorname{argmin}_{v \in T_x \mathcal{M}} f(x) + \langle \nabla f(x), v \rangle + \frac{1}{2t} \|v\|_F^2 + h(x + v)$$

Inexact Proximal Newton Methods

Riemannian version

Assumption:

- 1 Let $B_{x_*}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank;
- 2 There exists a neighborhood \mathcal{U} of $x_* = [\bar{x}_*^T, 0^T]^T$ on \mathcal{M} such that for any $x = [\bar{x}^T, \tilde{x}^T]^T \in \mathcal{U}$, it holds that $\bar{x} + \tilde{v} \neq 0$ and $\hat{x} + \hat{v} = 0$.

Theorem

Suppose that x_ be a local optimal minimizer. Under the above Assumptions, assume that $J(x_*)$ is nonsingular. Then there exists a neighborhood \mathcal{U} of x_* on \mathcal{M} such that for any $x_0 \in \mathcal{U}$, RPN Algorithm generates the sequence $\{x_k\}$ converging superlinearly to x_* .*

Inexact Proximal Newton Methods

Riemannian version

Assumption:

- ① Let $B_{x_*}^T = [\bar{B}_{x_*}^T, \hat{B}_{x_*}^T]$, where $\bar{B}_{x_*} \in \mathbb{R}^{j \times d}$ and $\hat{B}_{x_*} \in \mathbb{R}^{(n-j) \times d}$. It is assumed that $j \geq d$ and \bar{B}_{x_*} is full column rank;
- ② There exists a neighborhood \mathcal{U} of $x_* = [\bar{x}_*^T, 0^T]^T$ on \mathcal{M} such that for any $x = [\bar{x}^T, \tilde{x}^T]^T \in \mathcal{U}$, it holds that $\bar{x} + \tilde{v} \neq 0$ and $\hat{x} + \hat{v} = 0$.

Theorem

Suppose that x_* be a local optimal minimizer. Under the above Assumptions, assume that $J(x_*)$ is nonsingular. Then there exists a neighborhood \mathcal{U} of x_* on \mathcal{M} such that for any $x_0 \in \mathcal{U}$, RPN Algorithm generates the sequence $\{x_k\}$ converging superlinearly to x_* .

If the intersection of manifold and sparsity constraints forms an embedded manifold around x_* , then $\nabla^2 \bar{f}(x_*) - \bar{\mathcal{L}} \succeq 0$. If $\nabla^2 \bar{f}(x_*) - \bar{\mathcal{L}} \succ 0$, then $J(x_*)$ is nonsingular.

Inexact Proximal Newton Methods

Riemannian version

Smooth case: $\min_{x \in \mathcal{M}} f(x)$

- KKT conditions:

$$\nabla f(x) + \frac{1}{t}v + B_x \lambda = 0, \text{ and } B_x^T v = 0;$$

- Closed form solutions:

$$\lambda(x) = -B_x^T \nabla f(x), \quad v = -t \operatorname{grad} f(x);$$

- Action of $J(x)$: for $\omega \in T_x \mathcal{M}$

$$J(x)[\omega] = -t P_{T_x \mathcal{M}}(\nabla^2 f(x) - \mathcal{L}_x) P_{T_x \mathcal{M}} \omega = -t \operatorname{Hess} f(x)[\omega]$$

- $J(x)u(x) = -v(x) \implies \operatorname{Hess} f(x)[u(x)] = -\operatorname{grad} f(x);$
- It is the Riemannian Newton method;

Inexact Proximal Newton Methods

Numerical Experiments

Sparse PCA problem

$$\min_{X \in \text{St}(r,n)} -\text{trace}(X^T A^T A X) + \mu \|X\|_1,$$

where $A \in \mathbb{R}^{m \times n}$ is a data matrix and

$\text{St}(r, n) = \{X \in \mathbb{R}^{n \times r} \mid X^T X = I_r\}$ is the compact Stiefel manifold.

- $R_x(\eta_x) = (x + \eta_x)(I + \eta_x^T \eta_x)^{-1/2}$;
- $t = 1/(2\|A\|_2^2)$;
- Run ManPG until $\|v\|$ reaches 10^{-4} , i.e., it reduces by a factor of 10^3 . The resulting x as the input of RPN;

Inexact Proximal Newton Methods

Numerical Experiments




Figure: Random data. Left: different $n = \{100, 200, 300, 400\}$ with $r = 5$ and $\mu = 0.6$; Right: different $r = \{2, 4, 6, 8\}$ with $n = 300$ and $\mu = 0.8$

Summary and Future Work

Summary:

- A non-exhaustive review of nonsmooth optimization on manifolds;
- Euclidean/Riemannian proximal gradient methods;
- Inexact versions;
- Euclidean/Riemannian proximal Newton methods;

Future work:

- Accelerated version: $O(1/k^2)$ convergence rate analysis;
- Globalization for Riemannian Newton method;
- Design a Riemannian quasi-Newton method with superlinear local convergence rate;
- Generalize those methods to generic manifolds;

Thank you

Thank you!

References I



P.-A. Absil, R. Mahony, and R. Sepulchre.
Optimization algorithms on matrix manifolds.
Princeton University Press, Princeton, NJ, 2008.



Nicolas Boumal, P-A Absil, and Coralia Cartis.
Global rates of convergence for nonconvex optimization on manifolds.
IMA Journal of Numerical Analysis, 39(1):1–33, 02 2018.



G. C. Bento, J. X. de Cruz Neto, and P. R. Oliveira.
Convergence of inexact descent methods for nonconvex optimization on Riemannian manifold.
arXiv preprint arXiv:1103.4828, 2011.



Matthias Bollh ofer, Aryan Eftekhari, Simon Scheidegger, and Olaf Schenk.
Large-scale sparse inverse covariance matrix estimation.
SIAM Journal on Scientific Computing, 41(1):A380–A401, 2019.



S. Bonettini, M. Prato, and S. Rebegoldi.
Convergence of inexact forward–backward algorithms using the forward–backward envelope.
SIAM Journal on Optimization, 30(4):3069–3097, 2020.



Shixiang Chen, Shiqian Ma, Anthony Man-Cho So, and Tong Zhang.
Proximal gradient method for nonsmooth optimization over the Stiefel manifold.
SIAM Journal on Optimization, 30(1):210–239, 2020.



Patrick L. Combettes.
Solving monotone inclusions via compositions of nonexpansive averaged operators.
Optimization, 53(5-6):475–504, 2004.



Haoran Chen, Yanfeng Sun, Junbin Gao, Yongli Hu, and Baocai Yin.
Fast optimization algorithm on riemannian manifolds and its application in low-rank learning.
Neurocomputing, 291:59 – 70, 2018.

References II



Jalal M. Fadili and Gabriel Peyré.

Total variation projection with first order schemes.
IEEE Transactions on Image Processing, 20(3):657–669, 2011.



Octavian Eugen Ganea, Gary Becigneul, and Thomas Hofmann.

Hyperbolic entailment cones for learning hierarchical embeddings.
35th International Conference on Machine Learning, ICML 2018, 4:2661–2673, 2018.



M Seetharama Gowda.

Inverse and implicit function theorems for h-differentiable and semismooth functions.
Optimization Methods and Software, 19(5):443–461, 2004.



W. Huang and K. Wei.

Riemannian proximal gradient methods.
Mathematical Programming, 2021.
published online, DOI:10.1007/s10107-021-01632-3.



Wen Huang and Ke Wei.

An extension of fast iterative shrinkage-thresholding algorithm to Riemannian optimization for sparse principal component analysis.
Numerical Linear Algebra with Applications, page e2409, 2021.



Wen Huang, Meng Wei, Kyle A. Gallivan, and Paul Van Dooren.

A Riemannian Optimization Approach to Clustering Problems, 2022.



Jason D Lee, Yuekai Sun, and Michael A Saunders.

Proximal newton-type methods for minimizing composite functions.
SIAM Journal on Optimization, 24(3):1420–1443, 2014.



Boris S Mordukhovich, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang.

A globally convergent proximal newton-type method in nonsmooth convex optimization.
Mathematical Programming, pages 1–38, 2022.

References III



Vidvuds Ozolinš, Rongjie Lai, Russel Caflisch, and Stanley Osher.
Compressed modes for variational problems in mathematics and physics.
Proceedings of the National Academy of Sciences, 110(46):18368–18373, 2013.



Jong-Shi Pang, Defeng Sun, and Jie Sun.
Semismooth homeomorphisms and strong stability of semidefinite and lorentz complementarity problems.
Mathematics of Operations Research, 28(1):39–63, 2003.



Mark Schmidt, Nicolas Roux, and Francis Bach.
Convergence rates of inexact proximal-gradient methods for convex optimization.
In J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems*, volume 24. Curran Associates, Inc., 2011.



Silvia Villa, Saverio Salzo, Luca Baldassarre, and Alessandro Verri.
Accelerated and inexact forward-backward algorithms.
SIAM Journal on Optimization, 23(3):1607–1633, 2013.



Xiantao Xiao, Yongfeng Li, Zaiwen Wen, and Liwei Zhang.
A regularized semi-smooth newton method with projection steps for composite convex programs.
Journal of Scientific Computing, 76(1):364–389, Jul 2018.



Hui Zou, Trevor Hastie, and Robert Tibshirani.
Sparse principal component analysis.
Journal of Computational and Graphical Statistics, 15(2):265–286, 2006.



Y. Zhang, Y. Lau, H.-W. Kuo, S. Cheung, A. Pasupathy, and J. Wright.
On the global geometry of sphere-constrained sparse blind deconvolution.
In *Proceedings of IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2017.